

The Effect of Some Sudanese Medicinal Plant Extracts on Some Clinically Isolated Pulmonary Tuberculosis Bacteria

Elmutasim Osman Ibnouf¹*, Awatif Abdelbagi Elegami², Hisham Ali Waggiallah³

¹Department of Pharmaceutics, College of Pharmacy, Prince Sattam bin Abdulaziz University, Alkharj 11942, Saudi Arabia.

²Medicinal and Aromatic Plants Research Institute, National Centre for Research, Khartoum, Sudan. ³Department of Medical Laboratory Science, College of Applied Medical Science, Prince Sattam bin Abdulaziz University, Alkharj, Saudi Arabia.

ABSTRACT

Plants are an abundant source of biologically active compounds that have been shown to be effective antimicrobial agents. Many plants have traditionally been used to treat Mycobacterium infections. The goal of this research was to evaluate the effect of local medicinal plants on *Mycobacterium tuberculosis* isolates. The investigation was carried out on 130 pulmonary tuberculosis specimens from humans, obtained from Abo Anga and Al-Shaab hospitals. There were 103 infected males and 27 infected females, all of whom were between the ages of 20 and 30. The specimens were smeared, fixed, and stained directly with Ziehl-Neelsen. The acid-fast bacilli (AFB) were visible as red, straight, or slightly curved rods, singly or in small groups against a blue background.

The seven isolates were tested against extracts of Sudanese medicinal plants that had previously been shown to inhibit the growth of other microorganisms. The most active extracts were those of six plants particularly four extracts, belonging to two families that demonstrated activity against clinical isolates, and the minimum inhibitory concentrations were determined. Phytochemical screening was performed on the plants that demonstrated high anti-TB activity. It was concluded that specific tannins, saponins, and flavonoids play a significant role in anti-TB activity.

Key Words: Sudanese medicinal plants, Tuberculosis, Isolates, Phytochemical screening

eIJPPR 2021; 11(2):31-37

HOW TO CITE THIS ARTICLE: Ibnouf EO, Elegami AA, Waggiallah HA. The Effect of Some Sudanese Medicinal Plant Extracts on Some Clinically Isolated Pulmonary Tuberculosis Bacteria. Int J Pharm Phytopharmacol Res. 2021;11(2):31-7. https://doi.org/10.51847/wv0dR4Dgl5

INTRODUCTION

More than half of patients do not recover, the organisms become resistant to the medicines used, and since patients typically live in close quarters, drug-resistant organisms may be transmitted to others. The primary aim of anti-TB

Tuberculosis (TB) is still a major public health concern in various parts of the world, particularly in developing countries [1]. It is a contagious disease that is spreading in some parts of the world. It is estimated that in developing countries, 30-60% of adults have *Mycobacterium tuberculosis* [2]. Each year, about 8-10 million people develop clinical TB, and 3 million people die as a result of the disease [3].

Corresponding author: Elmutasim Osman Ibnouf Address: Department of Pharmaceutics, College of Pharmacy, Prince Sattam bin Abdulaziz University, Alkharj 11942, Saudi Arabia. E-mail: 🖂 elmutasim.ibnouf@Yahoo.com Received: 08 February 2021; Revised: 06 March 2021; Accepted: 04 April 2021 treatments are to cure each patient and thus reduce disease mortality and morbidity. The burden of TB is almost entirely carried by those living in low-income countries, with the disease accounting for a quarter of all avoidable deaths in these countries [4]. TB primarily affects young adults who are the economically productive segment of society [5]. The worldwide emergence of MDRTB is a major threat to TB control that is defined as TB caused by bacteria resistant to at least the 2 most important first-line medicines, isoniazid and rifampicin [6]. It is well understood that the emergence of drug resistance is linked to poor treatment practices, particularly erratic drug intake or monotherapy in the early months of treatment. Patients' noncompliance with treatment, especially failure to adhere

This is an **open access** journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

to multi-drug regimens and repeated interruptions can result in the selection of drug-resistant mutants of tuberculosis bacteria in such cases [7].

The Combretaceae family contains about 20 genera and 600 species. This plant is found primarily in tropical and subtropical regions such as Brazil and Africa [8]. This family is extensively used as traditional medicine [9]. Most African communities use these plants to treat snake and scorpion bites, as well as pneumonia, chest coughs associated with tuberculosis, fever, colds and sore throats, mental problems, and venereal diseases such as syphilis [10]. Phytochemical constituents found in the genus include alkaloids, saponins, tannins, and cardiac glycosides [11, 12]. To date, over 27000 alkaloid-based compounds have been recorded in the Dictionary of Natural Products [13].

This study aimed at evaluating the effect of local medicinal plants on *Mycobacterium tuberculosis* isolates.

MATERIALS AND METHODS

Study area and sample collection

One hundred and thirty sputum specimens were collected from patients of various ages and genders at Khartoum State's Abo-Anga and Al-Shaab hospitals who were clinically suspected of being infected with *M. tuberculosis*. The specimens were transferred to the microbiology lab for staining by Ziehl-Neelsen technique and culturing in Lowenstein-Jensen (L.J) medium [14, 15].

Chloroform and methanol extraction

Before testing for anti-TB activity, 50 grams of each plant sample were powdered and extracted. To extract substances, chloroform was added and heated at 35°C for 24 hours. The dry samples were placed in a soxhlet containing methanol (96%). To extract substances, samples were packed into soxhlet with methanol (96%) and heated at 40°C for 24 hours.

Testing of extracts for anti-TB activity

Using serial dilution, six different concentrations were created. Before solidification, 3 ml of each concentration

were added to seven bottles of L.J. medium. Finally, six different concentrations of the seven extracts were tested on each isolate.

Preliminary phytochemical screening of samples having antibacterial activity

Phytochemical analysis of the active morphological sample is extremely informative in terms of the nature of the constituents of each plant sample. It was thought essential to correlate the nature of chemical constituents and antibacterial activity tests for the detection of the various chemicals.

Preparation of the extracts

For 4 hours, 10 grams of powdered plant material were refluxed with 100ml of 80% ethanol. The cool solution was filtered, and 100 ml of 80% ethanol was passed through the filtrate. The extract was used in a variety of tests, including Unsaturated Sterols, Triterpenes, Alkaloids, Flavonoids, Saponins, Cyanogenic glycoside, Anthraquinone glycoside, Coumarins, and Anthraquinone glycoside.

RESULTS AND DISCUSSION

Clinical isolates

One hundred thirty sputum isolates were obtained from patients suspected of having *Mycobacterium tuberculosis* based on their symptoms.

Microscopical examination

Microscopically, 90 (69.23%) specimens showed positive Acid-Fast Bacilli (AFB) by Ziehl Neelsen stain, that appeared as red, straight or slightly curved, and 40 (30.77%) showed negative result.

Age and gender

The age group ranged between 10-60 years. Most of them were males, 62 (68.89%) males and 28 (31.11%) females (**Table 1**). Some of them were treated for more than two months.

Table 1. Gender and age rates of TB Patients									
Less than 10 Years	11-20	21-30	51-60) Over 60 Years 7					
(n)	3	50	40	17	12	8	130		
(%)	2.31%	38.46%	30.77%	13.08%	9.23%	6.15%	100%		
Gender	Male/Female 3	Male/Female	Male/Female	Male/Female	Male/Female	Male/Female	Male/Female		
	0	38 12	32 8	13 4	10 2	7 1	103 27		

Cultural examination

The cultivation of the obtained specimens on slopes of L.J. medium resulted in pure positive growth for M.

tuberculosis in 15 (16.67%) and contamination in 32 (35.56%), and 43 (47.78%) were considered negative after the optimal period of incubation.

Thermal, photo, and biochemical activities of the isolates According to the tests, seven isolates (46.67%) were identified as M. tuberculosis and tested against antibacterial agents (antibiotics and medicinal plant extracts).

Susceptibility testing to commonly used antibiotics

The in vitro studies of the susceptibility to anti-tuberculosis drugs were performed on L.J. medium and the inoculums of the isolated species were prepared. The resistance of the isolates to Streptomycin (SM), Rifampicin (RMP), Ethambutol (EMP), and Isoniazid (INH) was 3 (42.86%), 3 (42.86%), 4 (57.14%), and 2 (28.57%), respectively (Table 2). Multi-drug resistant (MDR) strains were 5 (71.43%), and those resistant to two or three drugs were 4 (57.14%) and 1 (14.29%), respectively. None of them showed resistance to all drugs (Table 3).

Table 2. Drug resistance Patterns in TB patients									
Drug	Streptor	nycin I	Rifampicin	Ethambutol	Isoniazid				
Seven Isolates	42.86 %		42.86 %	57.14 %	28.57 %				
	T	able 3. Drug resist	ance rate in TB pa	atients					
R- rate	\mathbf{R}_1	\mathbf{R}_2	R 3	R 4	S				
Strains	1	4	1	0	1				
Suams	4 4 8 8 8 4			0					

14.29 %

57.14 % Key: R₁: Resistant to one drug, R₂: Resistant to two drugs, R₃: Resistant to three Drug, R₄: Resistant to four Drug, S: Sensitive

Screening for the anti-TB activity of some Sudanese plants In the preliminary screening for the anti-TB activity of six Sudanese medicinal plants belonging to two families, the total number of extracts examined against the tested

14.29 %

organisms was seven. All of them were methanol extracts and exhibited inhibitory effects against one or more of the isolates (Table 4).

0

14.29 %

Table 4. Preliminary screening for the anti-TB activity of some Sudanese plants

Family/botanical name/vernacular name	Folkloric uses (Local uses)	Area of collection	Part used	Solvent used	Yield %
Combretaceae <i>C.aculeatum</i> Ver. Shiheit	For wounds healing	W.S	R	CHCL ₃ MeOH	1.82 10.68
Combretaceae		W.S	L	CHCL ₃	4.6
C.adenogonium				MeOH	16.12
Combretaceae	For hepatic disease	W.S	L	CHCL ₃	2.94
C.glutinosum				MeOH	22.52
Combretaceae	For jaundice	W.S	L	CHCL ₃	9.8
C.hartmannianum				MeOH	10.6
Ver. Habiel		-	S	CHCL ₃	6.6
				MeOH	8.6
Combretaceae	For eye-wash	W.S	L	CHCL ₃	3.24
T.laxiflora				MeOH	18.6
Ver. Daroot					
Loranthaceae	For wound healing	C.S	L	CHCL ₃	7.8
P.acaciae				MeOH	9.0
Ver.Inab elnabag					

Key: R: Root, L: Leaf, S: Seed, Ver.: Vernacular name, W.S: Western Sudan, C.S: Central Sudan

Determination of minimum inhibitory concentration (MIC) The MIC of seven extracts belonging to six medicinal plants, which showed antibacterial activity, was determined against the clinical isolates of M. tuberculosis. The results were summarized in Table 5 as mg/ml of crude extract.

International Journal of Pharmaceutical and Phytopharmacological Research (eIJPPR) | April 2021 | Volume 11 | Issue 2 | Page 31-37 Elmutasim Osman Ibnouf, The Effect of Some Sudanese Medicinal Plant Extracts on Some Clinically Isolated Pulmonary Tuberculosis Bacteria

Plant	Part used	Solvent used	Clinical isolates						
C.aculeatum	R	MeOH	75.0	37.5	37.5	75.0	75.0	75.0	75.0
C.adenogonium	L	MeOH	18.75	37.5	18.75	37.5	18.75	9.38	18.75
C.glutinosum	L	MeOH	150.0	150.0	150.0	75.0	150.0	150.0	75.0
C hartmannianna	L	MeOH	18.75	18.75	37.5	37.5	18.75	18.75	9.38
C.nartmannianum	S	MeOH	75.0	75.0	37.5	75.0	75.0	37.5	75.0
T.laxiflora	L	MeOH	37.5	37.5	37.5	37.5	75.0	37.5	9.38
P.acaciae	L	MeOH	9.38	37.5	9.38	9.38	9.38	37.5	9.38
Key: P. Poot I. Leaf	S. Sood								

Table 5. Minimum inhibitory concentration (mg/ml) of the crude extracts against the clinical isolates

Key: R: Root, L: Leaf, S: Seed

Preliminary phytochemical screening of selected Sudanese medicinal plants with a relatively high anti-TB activity Seven samples of different parts of six plants belonging to two families that proved to have a high anti-TB activity were examined phytochemically. The presence or absence of various types of secondary metabolites (alkaloids, flavonoids, terpenoids, saponins, anthraquinones, and cyanogenic glycosides) was tested. The results were summarized in **Table 6**. The susceptibility of the isolates to the extracts of the medicinal plants is shown in **Table 7**.

Table 6. Preliminary	v Phytochemica	1 Screening of	the most active	plants sample
Lable 0. I femininary	y i nytoenennea	i bereening of	the most detive	plants sample

Family	Botanical name	Part used	Alkaloids	Flavonoids	Coumarins	Saponins	Tannins	Anthra- quinones	Cyanogenic- glycosides	Sterols and/or triterpenes
Combretaceae	C. aculeatum	R	_	+	+	_	+	+	+	+
	C. adenogon-ium	L	_	+	+	+	+	_	+	+
	C. glutinosum	L	_	+	+	+	+	+	_	+
	C hartmann janum	L	+	+	+	+	+	+	+	+
	C. hartmann-lanum	S	_	+	+	+	+	+	_	+
	T. laxiflora	L	_	<u>+</u>	+	+	++	+	+	++
Loranthaceae	P. acaciae	L	_	+	+	+	+++	+	_	<u>+</u>

Key: L: Leaf (-): Not detectable, R: Root (±): Traces, S: Seed (+): Low concentration, (++): Medium concentration (+++): High concentration

Table 7. Susceptibility of the isolates to extracts of the medicinal plants

М	edicinal –	MIC of Medicinal Plant Extracts (mg/ml)									
Isolates		C. ac (R)	C. ad (L)	C. gl (L)	C. h (L)	C. h (S)	T. lax (L)	P. ac (L)			
TB 1		75.0	18.75	150.0	18.75	75.0	37.5	09.38			
TB 2		37.5	37.5	150.0	18.75	75.0	37.5	37.5			
TB 3		37.5	18.75	150.0	37.5	37.5	37.5	09.38			
TB 4		75.0	37.5	75.0	37.5	75.0	37.5	09.38			
TB 5		75.0	18.75	150.0	18.75	75.0	75.5	09.38			
TB 6		75.0	09.83	150.0	18.75	37.5	37.5	37.5			
TB 7		75.0	18.75	75.0	09.38	75.0	09.38	09.38			

Key: MIC: Minimum Inhibitory Concentration, C. ac: Combretum aculeatum, C. ad: Combretum adenogonium, C. gl: Combretum glutinosum, C. h: Combretum hartmannianum, T. lax: Terminalia laxiflora, P. ac: Plicosepalus acaciae, R: Root, L: Leaf, S: Seed

Given that *Mycobacterium tuberculosis* is the cause of tuberculosis for more than 100 years and that efficient

chemotherapy against the disease has been available for more than 50 years, it is a threat against humanity that this disease is still one of the world's most serious public health issues today [16]. TB treatment has been reshaped, and current therapy regimens are based on multidrug therapy, with 3-4 anti-tuberculosis drugs typically used. Even so, multidrug-resistant tubercle bacilli are becoming a problem for a variety of drugs, including isoniazid, ethambutol, rifampin, and streptomycin [17].

The drug resistance patterns of SM, RMP, EMP and INH were: (42.86%), (42.86%), (57.14%), (28.57%), respectively.

In the present study, the in vitro drug susceptibility testing showed that out of seven isolates, 1(14.29%) was fully sensitive to all the four drugs under investigation. However, 6 (85.71%) were resistant to one or more drugs. One (14.29%) isolate was resistant to a single drug, while 5 (71.43%) were multi-drug resistant (MDR), 4 (57.14%) were resistant to two drugs, and 1(14.29%) was resistant to three drugs. In another study, it was reported that primary and secondary drug resistance strains were 2.4% and 14.3%, respectively [18].

Drug-resistant tuberculosis is extremely difficult to treat, which necessitates the use of more and different medications over a longer period of time. Surgery is often required to remove areas of demolished lungs that are strongly infected with mycobacteria and are inaccessible to drugs [19]. Because mycobacteria are becoming increasingly resistant to traditional anti-tuberculosis drugs, there is an opportunity for new anti-tuberculosis agents. According to a recent WHO report, 2% of all tuberculosis cases worldwide are multi-drug resistant by description, meaning they are resistant to isoniazid + rifampicin (plus/minus other resistances). These cases can be cured in the United States and other high-resource countries, but at a high cost and with long courses of rather toxic medicines, posing a serious compliance dilemma [20]. It is critical to developing new anti-tuberculosis agents, preferably those that can be easily and cheaply manufactured from local sources. The use of antimicrobials derived from natural plants has a significant effect on human health care in developing countries.

Local healers have used herbal remedies in rural areas for centuries, and it has been developed in industrialized countries. Medicinal plant research has yielded a number of substances used in advanced medication to cure serious diseases [21]. In the situation of *C. aculeatum*, the root extract had the same effectivity as the seeds extract of *C. hartmannianum*, and both *C. hartmannianum and C. aculeatum* rank fifth in effectivity against the seven clinical isolates when compared to other plant extracts. In the case of *C. glutinosum*, the leaves extract had the lowest efficacy against clinical isolates when compared to all other extracts. C. aculeatum aerial part water decoction has antimycobacterial activity [22]. Martini *et al.* 2004 investigated the antibacterial activity of *C. erythrophyllum*

and discovered that seven antibacterial flavonoids had antimicrobial activity against Gram-negative and -positive strains in another study [23].

In the case of C. hartmannianum, the leaves extract was more effective than the Plicosepalus acaciae extract against the seven clinical isolates, with a lower MIC comparable to C. hartmannianum seeds extract. C. hartmannianum leaves extract exhibited lower MIC in five clinical isolates out of seven samples, while C. hartmannianum seeds extract showed lower MIC in two clinical isolates. In the case of C. adenogonium, the leaves extract demonstrated lower MIC in one clinical isolate and moderate affectivity in the remaining six isolates. T. laxiflora leaves extracts exhibited moderate effectivity in five clinical isolates, higher effectivity in one, and lower effectivity in the last. The leaves extract of *P.acaciae* demonstrated high activity against six isolates in the current study. El-Shafeia et al. (2017) analyzed P.acaciae for antibacterial activity and discovered that the plant contains a high concentration of flavonoids, tannins, and alkaloids [24].

El- Shafeia *et al.* (2017) evaluated the antimicrobial activity of chloroform, methanol, and aqueous extracts of *P. acaciae* leaves and stem, and discovered that the leaf methanol extract exhibited the highest level of activity against a variety of Gram-positive and Gram-negative clinical isolates from Sudanese patients [24].

Except for *C. adenogonium*, all of these plant extracts contained anthraquinones. Except for the leaves of *C. hartmannianum*, none of them were found to contain alkaloids. Saponins, which have been linked to antibacterial activity in some plants, were found in all plant extracts except *C. aculeatum*. Tannins were abundant in *T. laxiflora* and *P. acaciae*. *P. acaciae* leaves, on the other hand, were high in sterols and triterpenes. All plant extracts contained flavonoids and coumarins, and four plant samples contained cyanogenic glycosides [25].

These chemical groups may vary even within the same plant species' morphological parts. At this point, it is impossible to say whether the activity of these samples was caused by one or more of these chemical groups, or by other groups of compounds that were not tested. Further research is required to identify the compounds responsible for the activity in each sample.

CONCLUSION

The most effective extracts (7) from six plants (7 parts) belonging to two families demonstrated activity against clinical isolates, particularly four extracts, and the minimum inhibitory concentrations were determined. Phytochemical screening was performed on the plants that demonstrated high anti-TB activity. It can be stated that

specific tannins, saponins, and flavonoids play a significant role in the anti-TB activity.

Acknowledgments: This publication was supported by the Deanship of scientific research at Prince Sattam bin Abdulaziz University.

Conflict of interest: None

Financial support: None

Ethics statement: This study was approved by College of Pharmacy, University of Khartoum ethical board, and consent was also obtained from every participant in this study before the samples were obtained.

REFERENCES

- Waggiallah HA, Eltayeb LB, BinShaya AS, Elmahi OM. A saga of Hepcidin anti-microbial Effectiveness as Iron Acquisitor and Anemia Initiator in Mycobacterium Tuberculosis Infection. J Biochem Tech. 2020;11(2):128-34.
- [2] Bahmanjeh A, Kachooei SA, Ghasemi MF, Mosavari N, Hassanzadeh SM. Study on differentiation of pathogen-nonpathogen Mycobacterial infections using ESAT6-CFP10 in ELISA system. Arch Pharm Pract. 2020;11(2):28-36.
- [3] Mangwani N, Singh PK, Kumar V. Medicinal plants: Adjunct treatment to tuberculosis chemotherapy to prevent hepatic damage. J Ayurveda Integr Med. 2020;11(4):522-8.
- [4] Gemechu A, Giday M, Worku A, Ameni G. In vitro Anti-mycobacterial activity of selected medicinal plants against Mycobacterium tuberculosis and Mycobacterium bovis Strains. BMC Complement Altern Med. 2013;13(1):1-6.
- [5] Duarte R, Lönnroth K, Carvalho C, Lima F, Carvalho A, Mu noz-Torricoh M, et al. Tuberculosis, social determinants and co-morbidities (including HIV). Pulmono J. 2018;24(2):115-9.
- [6] Kurz SG, Furin JJ, Bark CM. Drug-Resistant Tuberculosis: Challenges and Progress. Infect Dis Clin North Am. 2016;30(2):509-22.
- [7] Ekor M. The growing use of herbal medicines: issues relating to adverse reactions and challenges in monitoring safety. Front Pharmacol. 2014;4:177.
- [8] Burman S, Bhattacharya K, Mukherjee D, Chandra G. Antibacterial Efficacy of Leaf Extracts of Combretum Album Pers. against some Pathogenic Bacteria. BMC Complement Altern Med. 2018;18(1):213.
- [9] Cock IE. The medicinal properties and phytochemistry of plants of the genus Terminalia

(Combretaceae). 2015;23(5):203-29.

Inflammopharmacol.

- [10] Masengu C, Zimba F, Mangoyi R, Mukanganyama S. Inhibitory Activity of Combretum zeyheri and its S9 Metabolites against Escherichia coli, Bacillus subtilis and Candida albicans. J Microbial Biochem Technol. 2014;6(4):228-35.
- [11] Zhang XR, Kaunda JS, Zhu HT, Wang D, Yang CR, Zhang YJ. The Genus Terminalia (Combretaceae): An Ethnopharmacological, Phytochemical and Pharmacological Review. Nat Prod Bioprospect. 2019;9(6):357-92.
- [12] Nyambuya T, Mautsa R, Mukanganyama S. Alkaloid extracts from Combretum zeyheri inhibit the growth of Mycobacterium smegmatis. BMC Complement Altern Med. 2017;17(1):124.
- [13] Amirkia V, Heinrich M. Alkaloids as drug-leads- A predictive structural and biodiversity-based analysis. Phytochem Lett. 2014;10:1874-3900.
- [14] Palucci I, Delogu G. Host directed therapies for tuberculosis: futures strategies for an ancient disease. Chemotherapy. 2018; 63(3):172-80.
- [15] Gygli SM, Borrell S, Trauner A, Gagneux S. Antimicrobial resistance in Mycobacterium tuberculosis: mechanistic and evolutionary perspectives. FEMS Microbiol Rev. 2017;41(3):354-73.
- [16] Ben Ayed H, Koubaa M, Marrakchi C, Rekik K, Hammami F, Smaoui F. Extrapulmonary Tuberculosis: Update on the Epidemiology, Risk Factors and Prevention Strategies. Int J Trop Dis. 2018;1(006).
- [17] Palange P, Narang R, Kandi V. Evaluation of Culture Media for Isolation of Mycobacterium Species from Human Clinical Specimens. Cureus. 2016;8(8):e757.
- [18] Hamusse SD, Teshome D, Hussen MS, Demissie M, Lindtjørn B. Primary and secondary anti-tuberculosis drug resistance in Hitossa District of Arsi zone, Oromia regional state, Central Ethiopia. BMC public health. 2016;16(1):593.
- [19] Subotic D, Yablonskiy P, Sulis G, Cordos I, Petrov D, Centis R, et al. Surgery and pleuro-pulmonary tuberculosis: a scientific literature review. J Thorac Dis. 2016;8(7):474-85.
- [20] Méda ZC, Huang CC, Sombié I, Konaté L, Somda PK, Djibougou AD, et al. Tuberculosis in developing countries: conditions for successful use of a decentralized approach in a rural health district. Pan Afr Med J. 2014;17:198.
- [21] Yuan H, Ma Q, Ye L, Piao G. The Traditional Medicine and Modern Medicine from Natural Products. Molecules. 2016;21(5):559.
- [22] Queiroz EF, Marcourt L, Kicka S, Rudaz S, Diop T, Soldati T, et al. Antimycobacterial activity in a single-

cell infection assay of ellagitannins from Combretum aculeatum and their bioavailable metabolites. J Ethnopharmacol. 2019;238:111832.

- [23] Martini ND, Katerere DR, Eloff JN. Biological activity of five antibacterial flavonoids from Combretum erythrophyllum (Combretaceae). J Ethnopharmacol. 2004;93(2-3):207-12.
- [24] El-Shafei G, Al-Hazmi B, Marghelani A, Al-Moalem D, Badr JM, Moneib NA. Antimicrobial activity of

different extracts of Plicosepalus acacia. Rec Pharm Biomed Sci. 2017;1(1):47-51.

[25] Albagouri A, Elegami A, Koko W, Osman E, Dahab M. In Vitro Anticercarial Activities of some Sudanese Medicinal Plants of the Family Combretaceae. J For Prod Ind. 2014;3(2):93-9.