



Developing a Green UV-Spectroscopic Method for Quercetin and Celecoxib Estimation Using AGREE Assessment Tool

Ujwala Desai^{1*}, Smita Pimple¹, Divya Dhamankar¹, Ganesh Desai¹, Vishal Vare¹

¹PES' Modern College of Pharmacy, Department of Pharmaceutics, Faculty of Pharmacy, Savitribai Phule Pune University, Pune, Maharashtra, India.

ABSTRACT

The Simultaneous Estimation of quercetin & celecoxib in original form was achieved by the development and validation of a innovative, straightforward, accurate, repeatable, and efficient simultaneous equation technique. The technique relied on measuring the absorbance of quercetin and celecoxib at 2 wavelengths (373 nm and 251 nm) and determining their λ_{max} . The conc. ranges of 4–12 $\mu\text{g/mL}$ for quercetin and 1.6–4.8 $\mu\text{g/mL}$ for celecoxib were found to have linear calibration curves. It was discovered that the calibration curves for quercetin and celecoxib had linear conc. ranges of 4–12 $\mu\text{g/mL}$ and 1.6–4.8 $\mu\text{g/mL}$, accordingly, with values for the correlation coefficient (R^2) of 0.999. A percentage RSD value was confirmed to be within limits ($RSD < 2\%$) in the precision study. In this work, the greenness of a binary combination of quercetin and celecoxib is measured using spectrophotometric techniques. Within the analytical community, the use of green analytical chemistry has become widely accepted. The Analytical GREENness calculator was used to analyze the assessment's greenness. The simultaneous equation methodology is applied in the technique for Quercetin at 373 nm and Celecoxib at 251 nm, respectively. The established procedures underwent validation in compliance with ICH recommendations. Quercetin and celecoxib can be successfully estimated together in pharmaceutical and pure dose forms using the suggested method.

Key Words: Quercetin, Celecoxib, Green analytical chemistry, Simultaneous estimation and UV spectroscopic technique

eIJPPR 2024; 14(4):9-16

HOW TO CITE THIS ARTICLE: Desai U, Pimple S, Dhamankar D, Desai G, Vare V. Developing a Green UV-Spectroscopic Method for Quercetin and Celecoxib Estimation Using AGREE Assessment Tool. Int J Pharm Phytopharmacol Res. 2024;14(4):9-16. <https://doi.org/10.51847/G4jD6FEZ11>

INTRODUCTION

Plant flavonol quercetin belongs to the flavonoid class of polyphenols. 2-(3,4-dihydroxyphenyl)-3,5,7-trihydroxy-4H-chromen-4-one is the chemical name for quercetin. Its potential therapeutic effects in several forms of arthritis, such as osteoarthritis (OA) and rheumatoid arthritis (RA), have been thoroughly researched. The chemical name of Celecoxib 4-[5-(4-methylphenyl)-3-(trifluoromethyl)pyrazol-1-yl] benzene sulfonamide is. Celecoxib is a nonsteroidal anti-inflammatory medication and a COX-2 inhibitor [1]. The FDA recommends celecoxib as a first-line analgesic for patients with osteoarthritis and rheumatoid arthritis. Compared to other NSAIDs like

Aspirin, Etodolac, Nabumethone and Salsalate which are prescribed for OA/RA dose of celecoxib is low (100 mg) thus proving better efficiency than other NSAIDs [2]. There aren't many documented UV spectroscopic techniques for figuring out Celecoxib and Quercetin separately or in various matrices with other medications [3-12]. These methods are selective and sensitive, but they frequently involve the use of dangerous solvents. Therefore, to have a greater positive impact on the environment and workforce, ecologically friendly solutions must be developed. Limiting reagent usage, favoring sustainable and minimally hazardous reagents, decreasing waste output as well as minimizing elements

Corresponding author: Ujwala Desai

Address: PES' Modern College of Pharmacy, Department of Pharmaceutics, Faculty of Pharmacy, Savitribai Phule Pune University, Pune, Maharashtra, India.

E-mail: ujudesai@gmail.com

Received: 27 April 2024; **Revised:** 08 July 2024; **Accepted:** 03 August 2024

This is an **open access** journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.



that may negatively affect the analyst are all important components of the evaluation of naturalness [13, 14]. Evaluated the greenness Analytical GREENness calculator (AGREE), [15-17]. To determine Quercetin and Celecoxib in an experimental formulation or synthetic combination selectively, green spectroscopic methods were developed and validated per the recommendations of the ICH [18-20].

MATERIALS AND METHODS

Apparatus

UV probe software and a UV 1800 double-beam spectrophotometer were used by Shimadzu. A set of matching quartz cells measuring one centimeter was used to record absorbance values.

Chemicals and reagents

Quercetin as well as celecoxib were kindly provided by Sisco Research Laboratories Pvt. Ltd Taloja Maharashtra, India and Lupin Pharmaceuticals Pvt. Ltd was used.

Greenness evaluation software

The Analytical GREENness calculator (AGREE), version 0.5, was used to measure and analyze the method performance's greenness.

Method

EQ approach A calibrated curve for quercetin and celecoxib was created by building the regression equation for each medication and plotting the absorbances at 373 nm and 251 nm, respectively, against the appropriate conc [21].

Assessment methods for green analytical chemistry

Agree

An analytical technique's greenness is indicated by the 12 parameters in the analytical Greenness tool being checked [22] as given in **Figure 1**. This software provides details on reagents, procedures, reagent toxicity, power usage, and other topics [23, 24].

Analytical Greenness report sheet

29/04/2024 12:28:36



Figure 1. Analytical Greenness Report Sheet

Preparations of standard stock solutions

2.50 mg of quercetin was dissolved in 10 mL of ethanol to create the standard stock solution, then the resulting mixture underwent ten minutes of sonication. For working standard solution with a concentration of 25 μ g/mL, 1 mL of the standard stock solution was extracted and diluted with 10 mL of the identical solution. Similarly, 1 mg of celecoxib was dissolved in 10 mL of ethanol to create the standard stock solution. After 10 minutes of sonication, the resultant solution's final volume was adjusted to 10 mL using ethanol. After additional dilution of this solution, a workable standard solution containing 10 mcg/mL was obtained and the graph are given in **Figure 2** [25, 26].

Diagram of overlain spectra




Figure 2. UV Spectra of Quercetin and Celecoxib

Simultaneous equation method development

The 200–400 nm UV range was scanned using the working solutions of both drugs. Both drug’s overlay spectra were captured [27]. Using the simultaneous equation approach, wavelengths 373 nm (the λ_{max} of Quercetin) and 251 nm (the λ_{max} of Celecoxib) were picked from overlain spectra for investigation of both drugs (λ_1 -373 nm for Quercetin and λ_2 -251 nm for Celecoxib) mentioned in **Figure 2** [28]. As a result, it might be able to identify the two drugs using the simultaneous equation approach or the from method [29-32].

Five standard solutions were prepared with a ratio of 2.5:1 (Quercetin: Celecoxib) with conc. of 4, 6, 8, 10, and 12 mcg/mL for Quercetin and 1.6, 2.4, 3.2, 4, and 4.8 μ g/mL for Celecoxib. Both 373 nm and 251 nm were used to estimate the comparable absorbance [33-35]. Using the SE method and the following formula, the concentrations of drug x (quercetin) and y (celecoxib) in the test solutions have been calculated:

$$C_x = \frac{A_2 a y_1 - A_1 a y_2}{a x_2 a y_1} - a x_1 a y_2 \quad (1)$$

$$C_y = \frac{A_1 a x_2 - A_2 a x_1}{a x_2 a y_1} - a x_1 a y_2 \quad (2)$$

The concentrations of Quercetin & Celecoxib are denoted by Cx and Cy, respectively. At 373 nm and 251 nm, the sample solution's absorbance is measured. Quercetin's absorptivity is measured at 373 nm and 251 nm, while Celecoxib's is measured at 373 nm and 251 nm, respectively [36].

Determination of absorptivity value

Table 1 display the absorptivity values of celecoxib & quercetin from every solution, which were determined using the formula [37].

$$Absorotivity = \frac{Absorbance}{Concentration} (gm/100mL) \quad (3)$$

As directed by ICH, [38] the developed approach was validated.

RESULTS AND DISCUSSION

Table 1. Absorptivity value for Quercetin and Celecoxib

Absorptivity value for Quercetin				
Concentration (mcg/mL)	Absorbance	Absorptivity	Absorbance	Absorptivity
	λ_1 -373 nm	λ_1 -373 nm	λ_2 -251 nm	λ_2 -251 nm
4	0.282	0.0705	0.310	0.0775
6	0.418	0.0696	0.595	0.0991
8	0.473	0.0591	0.645	0.0806
10	0.616	0.0616	0.718	0.0718
12	0.673	0.0560	0.890	0.0741
	Absorptivity for λ_1	0.06336	Absorptivity for λ_2	0.08062
Absorptivity value for Celecoxib				
Concentration (μ g/mL)	Absorbance	Absorptivity	Absorbance	Absorptivity
	λ_1 -373 nm	λ_1 -373 nm	λ_2 -251 nm	λ_2 -251 nm
1.6	0.031	0.019	0.202	0.0126
2.4	0.068	0.028	0.416	0.0173
3.2	0.108	0.033	0.719	0.0224
4	0.250	0.0625	0.829	0.0207
4.8	0.420	0.0875	0.978	0.0203
	Absorptivity for λ_1	0.046	Absorptivity for λ_2	0.0187

Agree

The criteria that have been presented align with the twelve SIGNIFICANCE principles, which can be adjusted in weight to offer a certain level of flexibility. Every one of the twelve input variables is converted to a normal 0–1 scale. The ultimate evaluation result is the sum of the assessment results for each premise. This leads to the creation of a graph that resembles a clock, with the overall score and color representation displayed in the middle [39].

The greenness evaluation report for the experimental procedures is shown and shown in **Figure 1**.

Specificity


By measuring the absorbance of Quercetin and Celecoxib separately at 373 nm and 251 nm in respect to the blank and synthesized excipients and comparing their absorbance with the blank and synthesized excipients, the specificity of the procedure was assessed [40-43]. The fact that there




was no interference at 251 and 373 nm suggests that the procedure is specific.

Linearity

Graphing the absorbance against the conc. ranges of 4, 6, 8, 10, and 12 mcg/mL for quercetin and 1.6, 2.4, 3.2, 4, and 4.8 mcg/mL for celecoxib allowed for the construction of the calibration curves (**Figure 3**). In these concentration ranges, the correlation coefficient values of the calibration curves were found to be linear. The sample's conc. and absorbance showed an acute relation, according to the results [44].



a) linearity of Quercetin in Simultaneous Estimation



b) Linearity of Celecoxib in Simultaneous Estimation

Figure 3. linearity of Quercetin and Celecoxib in Simultaneous Estimation

Accuracy

By using the recommended process on simulated excipients containing known levels of each medicine equal to 50%, 75%, 100%, 125%, and 150% of the claims made

on the labels of Quercetin and Celecoxib, the accuracy of the method was assessed. label claims [45, 46] **Table 2** presents the results of calculating the amount of Quercetin and Celecoxib recovered at each level.

Precision

By evaluating the sample solution's absorbance without altering the test protocol, a precision study was established, and the outcomes are shown in **Table 3** [47].

Repeatability

At least three duplicate measurements of the sample solution's absorbance were made for this investigation on the same day at 0 hours, 8 hours, and 16 hours shown in **Table 3**.

Intermediate precision

By measuring the sample solution's absorbance on three separate days using three distinct equipment and analysts, intermediate precision was achieved shown in **Table 3**.

Reproducibility

Two laboratories verified the method's reproducibility, and the outcomes were compared for celecoxib and quercetin shown in **Table 3**. The low percentage RSD (<2%) suggested that the procedure is accurate.

Robustness

By varying the wavelength from 373 nm to 251 nm by ± 1 nm, the robustness of the approach was assessed, and the findings are presented in **Table 4**.

Limit of quantification (LOQ) and limit of detection (LOD)

The calibration graph's slope and the response standard deviation were used to calculate LOD and LOQ. The LOD & LOQ of celecoxib were determined to be 13.075 ng/mL & 4.315 ng/mL, respectively. while quercetin were 0.884 ng/mL & 2.681 ng/mL [48, 49].

Stability

For three days at room temperature, the stability of the standard and sample solutions was examined, and daily absorbance measurements were made. Three days pass at room temperature before the sample solution begins to deteriorate. according to the findings of a calculation made on the amount of medication present [50].

Table 2. Recovery result for quercetin and celecoxib

Concentration (%)	Quercetin				Celecoxib				
	Quercetin/Celecoxib	Amt added (mg)	Recovered amt (mg)	Recovered amt (%)	RSD (%)	Amt added (mg)	Recovered amt (mg)	Recovered amt (%)	RSD (%)
50		1.94	1.95	100.51±0.32	0.32	0.72	0.71	98.61±0.76	0.76

75	2.19	2.21	100.91±0.17	0.17	0.89	0.81	100.09±0.51	0.51
100	2.50	2.52	100.8±0.15	0.15	1.0	0.9	99.50±0.38	0.38
125	2.69	2.68	99.62±0.18	0.18	1.90	1.86	99.15±0.61	0.61
150	2.88	2.91	100.69±0.07	0.07	2.21	2.11	98.47±0.50	0.50

The range of percentage recovery for Quercetin and Celecoxib was 99.62 to 100.51 and 98.47 to 100.09, respectively, suggesting that the method's accuracy was satisfactory.

Table 3. Precision result for quercetin and celecoxib

Specifications	Time of sampling	QUERCETIN			CELECOXIB		
		Amt present (mg)	Amt present (%)	RSD (%)	Amt present (mg)	Amt present (%)	RSD (%)
Repeatability (n=30)	0 hrs	1.9	100.46±0.87	0.48	0.6	100.88±0.98	0.97
	8th hrs	2.3	100.46±0.49	0.46	0.7	100.86±0.20	1.20
	16th hrs	2.2	100.46±0.25	0.25	0.8	100.08±0.76	0.36
Intermediate precision (n=6)	1st day	2.10	100.93 ±0.10	0.46	0.7	100.62±0.75	0.74
	2nd day	2.35	100.52 ±0.12	0.51	0.6	100.14±0.59	0.58
	3rd day	1.90	100.10 ±0.17	0.12	0.8	100.88±0.29	0.30
	Analyst-1	2.30	100.12 ±0.33	0.33	0.9	100.51±0.26	0.46
	Analyst-2	2.25	100.24±0.20	0.03	0.7	100.12 ±0.98	0.58
	Instrument-1	2.20	100.34±0.51	0.22	0.8	100.62 ±0.46	0.97
	Instrument-2	2.22	100.17± 0.25	0.25	0.6	100.34 ±0.98	0.37
	Reproducibility (n=6)	Lab-1	2.34	100.12± 0.20	0.12	0.6	100.23 ±0.20
			100.34± 0.12	0.20	0.8	100.66 ±0.38	0.37
Lab-2		2.22	100.12± 0.20	0.12	0.6	100.23 ±0.20	0.36
			100.34± 0.12	0.20	0.8	100.66 ±0.38	0.37

The method's precision was indicated by the low percentage RSD (<2%) for both celecoxib and quercetin.

Table 4. Result observed by changing the wavelength ±1 nm.

Quercetin				Celecoxib			
Wavelength (nm)	Amt present (mg)	Amt present (%)	RSD (%)	Wavelength (nm)	Amt present (mg)	Amt present (%)	RSD (%)
372	2.35	99.87±0.10	0.67	250	0.89	99.67±0.10	0.70
374	2.27	100 ± 0.24	0.45	252	0.86	99.23±0.24	0.39

The computed percent RSD value for PAR and FLU in the robustness investigation was less than 2%, indicating the robustness of the approach.

CONCLUSION

The simultaneous equation method that was created is straightforward, accurate, precise, and specific. The simultaneous estimate of quercetin and celecoxib in pure and pharmaceutical dosage forms without interference from excipients was demonstrated to be a reliable and selective process through statistical analysis. These medications can be routinely estimated using this new, straightforward method. To assess quercetin and celecoxib concurrently in their pure forms, synthetic mixtures, and dosage forms created in laboratories with varying ratios of active ingredients, new, accurate, and dependable spectrophotometric techniques have been developed.

Through the use of AGREE, the ecological impact of the techniques was investigated, and it was concluded that the suggested processes are environmentally beneficial. The fact that these methods are simple and don't require expensive or complex equipment is another advantage.

Acknowledgments: The authors thank P.E Society's Modern College of Pharmacy, Nigdi, Pune for providing the facilities required for carrying out research work.

Conflict of interest: None

Financial support: None

Ethics statement: None

REFERENCES

- [1] Guan F, Wang Q, Bao Y, Chao Y. Anti-rheumatic effect of quercetin and recent developments in nano formulation. *RSC Adv.* 2021;11(13):7280-93.
- [2] Patel BH, Teraiya NH. Simultaneous estimation and stability indicating method of vildagliptin and dapagliflozin by rp-hplc in pharmaceutical dosage form. *Res Sq.* 2024;2-10. doi:10.21203/rs.3.rs-4080300/v1
- [3] Darji H, Dedania Z, Dedania R, Jain V. Eco-friendly based HPLC and UV-spectrophotometric methods for simultaneous estimation of efonidipine hydrochloride ethanolate and chlorthalidone in their dosage form. *Green Anal Chem.* 2024;8:100092.
- [4] Mevada S, Patel H, Shukla S. Simultaneous equation method for the estimation of gallic acid and apigenin by UV-visible spectrophotometry. *Accredit Qual Assur.* 2024;29(1):11-7.
- [5] Sebaiy M, El-Adl S, Elbaramawi S, Abdel-Raheem S, Nafie A. Developing a highly validated and sensitive HPLC method for simultaneous estimation of cefotaxime and paracetamol in pure and pharmaceutical preparations. *Curr Chem Lett.* 2024;13(2):435-44.
- [6] Prasanth SS, Panchami GS, Reeba KR, Thomas B, Shibila NT, Niyas Sh. Method development and validation for the simultaneous estimation of amiodarone and propranolol by using UV spectroscopy. *Int J Pharm Sci Rev Res.* 2023;79(2):73-7. doi:10.47583/ijpsrr.2023.v79i02.013
- [7] Akabari AH, Mistry P, Patel SK, Surati J, Patel SP, Shah U. Simultaneous estimation of fimasartan potassium trihydrate and atorvastatin calcium with greenness assessment using HPLC and UV spectrophotometric methods. *Green Anal Chem.* 2023;6:100067. doi:10.1016/j.greeac.2023.100067
- [8] Kayali Z, Obaydo RH, Sakur AA. Spider diagram and sustainability evaluation of UV-methods strategy for quantification of aspirin and sildenafil citrate in the presence of salicylic acid in their bulk and formulation. *Heliyon.* 2023;9(4). doi:10.1016/j.heliyon.2023.e15260
- [9] Darji H, Dedania Z. Simultaneous estimation of azelmidipine and metoprolol succinate with greenness assessment using HPLC and UV-spectrophotometric methods. *Green Anal Chem.* 2023;7:100079. doi:10.1016/j.greeac.2023.100079
- [10] Anwar MS, Khan A, Khan I, Khan SA, Ahmad L, Kaleem WA, et al. Evaluation of marketed herbal medicines for the simultaneous estimation of steroidal adulterants using FTIR and RP-HPLC-UV. *Microchem J.* 2023;190:108745.
- [11] Afify KK, Ali R, El-Dosoky MA, Nassar MW. HPLC/UV approach method for the first simultaneous estimation of molnupiravir and ertapenem as a binary mixture in human plasma and dosage form as a regimen for COVID-19 treatments. *BMC Chem.* 2023;17(1):121.
- [12] Bherje S, Patel M. A green perspective on simultaneous HPLC and UV spectrophotometric estimation of Udenafil and Dapoxetine hydrochloride in pharmaceutical formulations. *Green Anal Chem.* 2024;9:100106.
- [13] Arji SR, Eranki SS, Kadimi A, Sarella PN, Mangam VT. Development and validation of an HPLC method for the simultaneous estimation of salbutamol, theophylline and ambroxol in tablet dosage form. *Int J Sci Res Arch.* 2023;10(2):634-45.
- [14] Shi M, Zheng X, Zhang N, Guo Y, Liu M, Yin L. Overview of sixteen green analytical chemistry metrics for evaluation of the greenness of analytical methods. *TrAC Trends Anal Chem.* 2023:117211.
- [15] Elshahed MS, Toubar SS, Ashour AA, El-Eryan RT. Novel sensing probe using Terbium-sensitized luminescence and 8-hydroxyquinoline for determination of prucalopride succinate: Green assessment with Complex-GAPI and analytical Eco-Scale. *BMC Chem.* 2022;16(1):80. doi:10.1186/s13065-022-00876-0
- [16] Pena-Pereira F, Wojnowski W, Tobiszewski M. AGREE—Analytical GREENness metric approach and software. *Anal Chem.* 2020;92(14):10076-82. doi:10.1021/acs.analchem.0c01887
- [17] Nowak PM, Kościelniak P. What color is your method? Adaptation of the RGB additive color model to analytical method evaluation. *Anal Chem.* 2019;91(16):10343-52.
- [18] Gowdak LH. The role of ivabradine in managing symptomatic patients with chronic coronary syndromes: A clinically oriented approach. *Cardiol Ther.* 2022;11(1):163-74.
- [19] Hasan N, Imran M, Jain D, Shamim A, Beg S, Kesharwani P, et al. Rapid analytical method development and validation for the simultaneous estimation of 5-Fluorouracil and cannabidiol in plasma and lipid-based nanoformulations. *Curr Anal Chem.* 2022;18(7):798-808.
- [20] Babar SA, Padwal SL, Bachute MT. Qbd based RP-HPLC method development and validation for simultaneous estimation of amlodipine besylate and lisinopril dihydrate in bulk and pharmaceutical dosage form. *J Pharm Res Int.* 2021;33(43A):143-64.
- [21] Awasthi A, Kumar A, Kumar R, Vishwas S, Khursheed R, Kaur J, et al. RP-HPLC method development and validation for simultaneous estimation of mesalamine and curcumin in bulk form

- as well as nanostructured lipid carriers. *S Afr J Bot.* 2022;151:529-37.
- [22] Mortada WI, Zedan HE, Khalifa ME. Spectrophotometric determination of trace vanadium in fresh fruit juice samples by ion pair-based surfactant-assisted microextraction procedure with solidification of floating organic drop. *Spectrochim Acta A Mol Biomol Spectrosc.* 2023;302:123107. doi:10.1016/j.saa.2023.123107
- [23] Wojnowski W, Tobiszewski M, Pena-Pereira F, Psillakis E. AGREEprep-analytical greenness metric for sample preparation. *TrAC Trends Anal Chem.* 2022;149:116553. doi:10.1016/j.trac.2022.116553
- [24] Piponski M, Balkanov T, Logoyda L. Development and validation of a fast and simple HPLC method for the simultaneous determination of bisoprolol and enalapril in dosage form. *Pharmacia.* 2021;68:69-77.
- [25] Ali SN, Mobina L, Mehfuza M, Seema P, Ahmed A, Khan GJ. Analytical method development and validation and forced degradation stability-indicating studies of Favipiravir by RP-HPLC and UV in bulk and pharmaceutical dosage form. *J Pharm Res Int.* 2021;33(48B):254-71.
- [26] Zaki MM, Abdelwahab NS, Ali AA, Sharkawi SM. Simultaneous determination of bisoprolol fumarate and rosuvastatin calcium in a new combined formulation by validated RP-HPLC. *Eur J Chem.* 2019;10(1):52-6.
- [27] Foch C, Allignol A, Hohenberger T, Boutmy E, Schaefer S, Hostalek U. Effectiveness of bisoprolol versus other β -blockers and other antihypertensive classes: A cohort study in the clinical practice research datalink. *J Comp Eff Res.* 2022;11(6):423-36.
- [28] Mehmood T, Hanif S, Azhar F, Ali I, Alafnan A, Hussain T, et al. HPLC method validation for the estimation of lignocaine HCl, ketoprofen and hydrocortisone: Greenness analysis using AGREE score. *Int J Mol Sci.* 2022;24(1):440. doi:10.3390/ijms24010440
- [29] Attimarad M, Venugopala KN, Shafi S, Balgoname AA, Altaysan AI. Smart spectrophotometric method development for simultaneous estimation of antidiabetic drugs in formulations. *Indian J Pharm Educ Res.* 2022;56(1).
- [30] El-Zahry MR. Electrochemical sensor based on molybdenum-doped graphene oxide nanorods anchored carbon spheres/vanadium pentoxide nanocomposites for simultaneous determination of diclofenac sodium and posaconazole. *Microchem J.* 2024:111593.
- [31] Agrawal S, Gurjar P, Katheriya B. Analytical method development and validation for simultaneous estimation of Trimetazidine hydrochloride and Metoprolol succinate using HPTLC. *Curr Pharm Anal.* 2019;15(3):243-50.
- [32] Andal P, Kshatriya AG, Mhaske A, Naik MS. A new validated analytical method for simultaneous estimation of trimetazidine dihydrochloride and bisoprolol fumarate from dosage forms using HPLC. *J Pharm Negative Res.* 2022:7398-407.
- [33] Mabrouk M, Abdel Hamid M, Michael M. Green simultaneous determination of amlodipine besylate and celecoxib by dual wavelength and simultaneous equation spectrophotometric methods. *J Adv Med Pharm Res.* 2021;2(1):16-23.
- [34] Palur K, Archakam SC, Koganti B. Chemometric assisted UV spectrophotometric and RP-HPLC methods for simultaneous determination of paracetamol, diphenhydramine, caffeine and phenylephrine in tablet dosage form. *Spectrochim Acta A Mol Biomol Spectrosc.* 2020;243:118801
- [35] Suman SS, Chaubey R. Method development of metformin, saxagliptin and dapagliflozin in marketed formulation by HPLC. *Int J Health Sci.* 2022;6(S8):5927-36.
- [36] Kharat AR, Pakhare MK, Kharat KR. Stability indicating uv spectrophotometric method development and validation for simultaneous estimation of pioglitazone hydrochloride and glimepiride in bulk drug and pharmaceutical formulation. *Rasayan J Chem.* 2024;17(3).
- [37] Kannaiah KP, Sugumaran A. Environmental benign AQbD based estimation of ketoconazole and beclomethasone by RP-HPLC and multi-analytical UV spectrophotometric method. *Microchem J.* 2022;172:106968.
- [38] Gandhi NK, Ezhava SB. Stability indicating analytical method development using quality by design (QbD) approach for simultaneous estimation of ivabradine and metoprolol. *Res J Pharm Technol.* 2021;14(11):5937-44.
- [39] Ma D, Xu T, Cai G, Wu X, Lei Z, Liu X, et al. Effects of ivabradine hydrochloride combined with trimetazidine on myocardial fibrosis in rats with chronic heart failure. *Exp Ther Med.* 2019;18(3):1639-44.
- [40] Yosrey E, Elmansi H, Sheribah ZA, Metwally ME. Micellar-emphasized simultaneous determination of ivabradine hydrochloride and felodipine using synchronous spectrofluorimetry. *Luminescence.* 2022;37(4):569-76.
- [41] Attimarad M, Elgorashe RE, Subramaniam R, Islam MM, Venugopala KN, Nagaraja S, et al. Development and validation of rapid RP-HPLC and green second-derivative UV spectroscopic methods for simultaneous quantification of metformin and

- remogliflozin in formulation using experimental design. *Separations*. 2020;7(4):59.
- [42] Veeshma A, Priyanka S, Kumar PK, Sirisha K. Simultaneous estimation of ciprofloxacin and metronidazole in bulk and tablet formulation by UV-spectrophotometry. *Int J Pharm Sci Res*. 2021;12(4):2247-56.
- [43] Aayushi MS, Surati JS. UV-spectrophotometric method development and validation for simultaneous estimation of hydroxychloroquine sulfate and nitazoxanide in synthetic mixture. *Int J Pharm Sci Res*. 2022;13(3):1374-82.
- [44] Polker A, Padmavathi Y, Kumar R, Babu NR. Quantitative bioanalytical and analytical methods for estimation of ivabradine hydrochloride in pure and pharmaceutical dosage form. *J Rep Pharm Sci*. 2020;9(1):1-0.
- [45] Gandhi S, Manikandan A, Rao SV. Novel stability indicating RP-UPLC method for simultaneous determination of ivabradine and metoprolol drug materials in bulk and their pharmaceutical dosage forms. *Res J Pharm Technol*. 2020;13(1):250-4.
- [46] Deshmukh M, Bhusari S, Wakte P. Development and validation of q-absorbance ratio spectrophotometric method for the simultaneous estimation of ciprofloxacin and ornidazole; in combined pharmaceutical dosage form. *J Drug Deliv Biother*. 2024;1(02):01-15.
- [47] Almaqtari MA, Al-Odaini N, Alarbagi FA, Alattab BM, Al-Maydama HM. Development and validation of a new spectrophotometric method for simultaneous determination of spiramycin and metronidazole in tablet pharmaceutical dosage forms using chemometrics technique in comparison with HPLC. *JAST*. 2023;1(1).
- [48] Patel D, Dobariya J, Pradhan P, Patel G, Meshram D. Development and validation of UV Spectrophotometric methods for simultaneous estimation of Lobeglitazone Sulfate and Glimperide in combined dosage form. *Drug Anal Res*. 2024;8(1):62-9.
- [49] Qasim FO, Mohammed NM. A nanodrop spectrophotometric method and stability indicating for determination of amlodipine besylate in pharmaceutical formulations of Kurdistan of Iraq. *Sci J Univ Zakho*. 2021;9(1):25-9.
- [50] Kowtharapu LP, Katari NK, Muchakayala SK, Pydimarry SP, Rekulapally VK, Sandoval CA. QbD green analytical procedure for novel study of a genotoxic and carcinogenic compound trace determination in physiological solution compatibility. *Sustain Chem Pharm*. 2023;33:101079.