



Antibiotic Resistance Trend of Uropathogens in a Tertiary Teaching Hospital in Rasht, Iran, a Longitudinal Study

Sonbol Taramian¹, lida Mahfoozi^{2*}, Mostafa Soodmand³

¹ Infectious Diseases Department, School of Medicine, Guilan University of Medical Sciences, Rasht, Iran.

² Assistant professor of Infectious Diseases, Guilan University of Medical Sciences, Rasht, Iran.

³ MSc Student of Medical-surgical Nursing, Student Research Committee, School of Nursing and Midwifery, Guilan University of Medical sciences, Rasht, Iran.

ABSTRACT

Background: The present study was conducted to determine the most prevalent isolates in urine cultures and resistance profile of the isolated pathogens and trend of changing resistance to common antibiotics used for the treatment of urinary tract infections. **Materials and methods:** In this longitudinal - descriptive study, the data of 2060 positive urine cultures and the susceptibility testing results of inpatients and outpatients' urine specimens from 2007 to 2014 were studied. The susceptibility testing using Kirby-Bauer disk diffusion method according to CLSE standards was done on isolated uropathogens for t ceftriaxone, ceftazidime, gentamicin, amikacine, ciprofloxacin and cotrimoxazole, piperacillin, nitrofurantoin and imipenem. **Results:** The most prevalent pathogen which was isolated was E.coli, 1141 (55.4%). The resistance trend of E.coli isolates for antibiotics showed increased resistance for: ceftazidim (67% to 81%), ciprofloxacin (57%-77%), piperacillin (20%-87%); stable resistance rate for : ceftriaxone(73%-70%), cotrimoxazole (79%-74%) and decreased resistance for : nitrofurantoin (32%-8%), gentamicine (49%-37%) and amikacine (21%-13%). 50% of E.coli isolates were co-resistant to ceftriaxone and ciprofloxacin and trimetoprim-sulfamethoxazole. **Conclusion:** Antibiotic resistance of uropathogens for most of the important antibiotics empirically used for UTI was very high and increasing in this hospital.

Key Words: Antibiotic resistance, urinary tract infection, E.coli, multidrug resistant

eIJPPR 2019; 9(2):98-102

HOW TO CITE THIS ARTICLE: Sonbol Taramian, lida Mahfoozi, Mostafa Soodmand (2019). "Antibiotic Resistance Trend of Uropathogens in a Tertiary Teaching Hospital in Rasht, Iran, a Longitudinal Study", International Journal of Pharmaceutical and Phytopharmacological Research, 9(1), pp.98-102.

INTRODUCTION

Globally, it has been estimated that there are 150 million urinary tract infections every year [1]. Patient morbidity, costs of treatment, rates of hospitalization, and use of broad-spectrum agents are remarkably increased by antimicrobial resistance [2]. There are a limited and predictable spectrum of organisms which lead to urinary tract infections. Considering both outpatients and inpatients, Escherichia coli has been the main urinary tract pathogen, causing 75 to 90% of the uncomplicated urinary tract infection isolates. Staphylococcus

saprophyticus, Klebsiella spp., Proteus spp., Enterococcus spp., and Enterobacter spp. have been the pathogens which were less commonly isolated from outpatients [3]. Antibiotic resistance development has been a major issue for managing infections both in community and hospital settings [4]. The Infectious Diseases Society of America recommends that physicians obtain information on local resistance rates and that ongoing surveillance be carried out to check the changes in susceptibility of uropathogens [5]. The present study was conducted to determine the most prevalent isolates in urine cultures of inpatients and institutional in vitro susceptibility and resistance profile of the isolated pathogens and the trend of changing

Corresponding author: lida Mahfoozi

Address: School of Medicine, Guilan University of Medical sciences, Rasht, Iran.

E-mail: drlidadmahfoozi@yahoo.com

Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 25 October 2018; **Revised:** 18 February 2019; **Accepted:** 23 February 2019



resistance to common antibiotics used for the treatment of urinary tract infections.

MATERIALS AND METHODS

Setting

This longitudinal - descriptive study was conducted in teaching hospitals, affiliated with Guilan University of Medical Sciences in Rasht from 2007 to 2014. For this purpose, the data of positive urine cultures and susceptibility testing results of inpatients and outpatients urine specimens from microbiology laboratory of Razi hospital (a tertiary teaching general hospital of Guilan university of medical sciences in Rasht) from 2007 to 2014 was gathered. During this 7- year period, 2060 positive urine culture results were obtained.

Data Collection

Susceptibility testing using Kirby-Bauer disk diffusion method according to CLSE standards was done on isolated uropathogens for ceftriaxone, ceftazidime, gentamicin, amikacine, ciprofloxacin and cotrimoxazole on all isolates. The susceptibility testing for piperacillin, nitrofurantoin and imipenem in some periods according to the availability of disks was done.

Statistical Analysis

The obtained information was inserted into SPSS-20, and descriptive statistics (percentage, frequency) were utilized.

RESULTS

During this 7- year period (2007-2014), 2060 positive urine culture results were obtained from the hospital's microbiological laboratory. The most prevalent pathogen which was isolated was E.coli, 1141 (55.4%). 60% of specimens were obtained from women. Other pathogens

in order of prevalence were: citrobacter spp (20%), enterobacter spp (9.8%), klebsiella spp (4%), enterococcus (2.8%), pseudomonas spp (2%), Staphylococcus saprophyticus (1%) (table 1).

The resistance trend of E.coli isolates for antibiotics showed the increased resistance for : ceftazidim(67% to 81%), ciprofloxacin (57%-77%), piperacillin (20%-87%); the stable resistance rate for: ceftriaxone (73%-70%), cotrimoxazole (79%-74%); and the decreased resistance for: nitrofurantoin (32%-8%), gentamicine (49%-37%) and amikacine (21%-13%)(table 2).

33% of E.coli isolates were co- resistant to ceftriaxone, ciprofloxacin and gentamicine or amikacin. 50% of E.coli isolates were co-resistant to ceftriaxone and ciprofloxacin and trimetoprim-sulfamethoxazole (MDR isolates). Citrobacter and enterobacter had very similar resistance rates: 77% and 81% for ceftriaxone, 42% and 42% for gentamicine, 32% and 39% for amikacine, 50% and 52% for ciprofloxacin, 84% and 85% for piperacilline , 78% and 83% for cotrimoxazole and 56% and 71% for nitrofurantoin; respectively. klebsiella.spp and pseudomonas aerogenoza showed significantly higher resistance rates than citrobacter and enterobacter (table 3).

For pseudomonas, the resistance rates for antibiotics ceftazidime and cefepime: two third generation of cephalosporine was highly active against pseudomonas aeruginosa : 90% and 86%; respectively, not shown in table 3. The resistance rates of enterococcus spp, a gram positive microorganism occasionally causing UTI also revealed very high rates: 57% for gentamicine, 67% for amikacine, 64% for ciprofloxacin. Also the resistance rates for doxycycline was (50%), for penicillin was (85%), and for ampicillin was (69%) ; the antibiotics sometimes used for infections caused by gram positive cocci, were very high (Not shown in table 3).

Table 1: microorganisms isolated from Urine cultures in Razi hospital: 2007-2014

| Y | N | E.coli N (%) | Citro N (%) | Entro N (%) | Pseudo N (%) | Klebsiella N (%) | Staph sapro N (%) | Enterococcus N (%) | Others N (%) |
|-------|------|-----------------|----------------|----------------|-----------------|---------------------|----------------------|-----------------------|-----------------|
| 2007 | 183 | 103(56) | 22(12) | 29(16) | 7(4) | 3(1.6) | 1(0.5) | 5(2.7) | 13(7) |
| 2008 | 188 | 107(57) | 31(17) | 24(13) | 6(3) | 0(0) | 2(1) | 6(3) | 12(6.3) |
| 2009 | 208 | 110(53) | 46(22) | 21(10) | 4(1.9) | 2(0.9) | 5(2.4) | 5(2.4) | 15(7.2) |
| 2010 | 300 | 181(60) | 65(22) | 15(5) | 8(3) | 4(1.3) | 4(1.3) | 7(2.3) | 16(5.3) |
| 2011 | 191 | 114((60) | 49(25) | 22(19) | 6(3) | 3(1.6) | 2(1) | 5(2.6) | 14(7.3) |
| 2012 | 235 | 113(48) | 63(27) | 29(12) | 5(2) | 4(1.7) | 3(1.2) | 6(2.5) | 12(5) |
| 2013 | 330 | 198(60) | 77(23) | 30(9) | 0(0) | 0(0) | 2(0.6) | 8(2.4) | 15(4.5) |
| 2014 | 425 | 215(50) | 65(15) | 32(8) | 11(3) | 67(16) | 3(0.7) | 15(4) | 17(4) |
| Total | 2060 | 1141(55) | 418(20) | 202(10) | 47(2.3) | 83(4) | 22(1) | 57(2.7) | 114(5.5) |

Table 2: Antibiotic resistance percentage of E.coli isolated from urine cultures in Razi hospital: 2007-2014

| Y | N | cefta | genta | amika | cipro | pipra | ceftria | imip | cotri | nitro |
|------|-----|-------|-------|-------|-------|-------|---------|------|-------|-------|
| 2007 | 103 | 67% | 49% | 21% | 57% | - | 73% | - | 79% | 32% |
| 2008 | 107 | 66% | 53% | 18% | 61% | - | 63% | - | - | - |
| 2009 | 110 | 70% | 39% | 13% | 59% | - | 65% | - | - | - |
| 2010 | 181 | 78% | 54% | 19% | 77% | - | 80% | - | - | - |
| 2011 | 212 | 57% | 42% | 11% | 72% | 20% | 68% | 73% | 73% | 12% |
| 2012 | 113 | 82% | 46% | 23% | 76% | 50% | 74% | 79% | 82% | 15% |
| 2013 | 198 | 81% | 47% | 22% | 70% | 39% | 72% | - | 66% | 11% |
| 2014 | 215 | - | 37% | 13% | 77% | 87% | 70% | - | 74% | 8% |

Table 3: antibiotic resistance percentage of other uropathogens isolated from urine cultures in Razi Hospital: 2007-2014

| pathogen | N | ceftria | genta | amika | cipro | pipra | imip | cotri | nitro |
|--------------|-----|---------|-------|-------|-------|-------|------|-------|-------|
| citrobacter | 418 | 77% | 42% | 32% | 50% | 84% | - | 78% | 56% |
| entrobacter | 202 | 81% | 42% | 39% | 52% | 85% | - | 83% | 71% |
| pseudomonas | 47 | - | 90% | 73% | 100% | 91% | 50% | - | - |
| klebsiella | 83 | 90% | 75% | 66% | 84% | 92% | - | 91% | 75% |
| enterococcus | 57 | - | 57% | 67% | 64% | - | - | - | 30% |

DISCUSSION

In this study, the most prevalent microorganism isolated from urine cultures of patients admitted to this hospital for UTI or other diagnoses, was E.coli (mean =55% over 7-year period :range 48%-60%). As in other studies, E.coli has been consistently the most prevalent microorganism causing urinary tract infection over the world [6-11].

In this study, in comparison to other similar studies worldwide, as follows, different resistance patterns were considered according to the local study results. The Infectious Diseases Society of America has advocated trimethoprim-sulfamethoxazole (SXT) as initial therapy for females with acute uncomplicated bacterial cystitis in settings where the prevalence of SXT resistance does not exceed 10 to 20%. The susceptibility testing data from The Surveillance Network (TSN) Database-USA (n = 286,187) from 1995 to 2001 showed the resistance rates among E. coli isolates to SXT (range, 14.8 to 17.0%), ciprofloxacin (range, 0.7 to 2.5%), and nitrofurantoin (range, 0.4 to 0.8%) [12]. In this study, the resistance of E.coli to cotrimoxazole (79%-74%) and ciprofloxacin (57%-77%) nitrofurantoin (32%-8%), was much higher, so other antibiotics should be recommended for UTI in this region. The comparison of the current study to a similar study from India indicated more similar resistance pattern to this study: ceftriaxone (60.5%), ceftazidime (57.1%), gentamicin (59.6%), nitrofurantoin (26.9%), meropenem (9.8%), ciprofloxacin (73%) and amikacin (23%) [13]. In the present study, the most important and the most prevalent co resistance phenotypes were to ceftriaxone, ciprofloxacin and trimetoprim-

sulfamethoxazole (50% of isolates) then to ceftriaxone and ciprofloxacin and (aminoglycosides: gentamicine or amikacin) (33% of isolates). In the same study from USA in 2001, it was indicated that the most frequent co resistant phenotypes were resistant to ampicillin and SXT (12.0% of all isolates; 82.3% of coresistant isolates) and to ampicillin, ciprofloxacin, and SXT (1.4% of all isolates; 9.9% of coresistant isolates) [12].

The Study for Monitoring Antimicrobial Resistance Trends program collected 1643 E.coli isolates in 2009–2010 from urinary tract infection (UTI) specimens of hospitalized patients in countries around the world. Ertapenem and imipenem were the most active agents tested, inhibiting >98% of all E. coli phenotypes [14]. The data of resistance to imipenem for E.coli isolates were collected for two years (2011-2012) that showed a high resistance rate (73%-79%). In that study, Amikacin and piperacillin–tazobactam achieved 90% inhibition on E.coli isolates while in the current study, the resistance rate to amikacin (21%-13%) and piperacillin (20%-87%) was very high [15]. FQ resistance in UTI pathogens also increased globally, with the SMART study reporting the resistance rates between 6% and 75% in Gram-negative urinary pathogens isolated from hospitalized patients [14]. In this study, the resistance rate of E.coli and other uropathogens isolated from urine samples of patients for ciprofloxacin, as one of the most prescribed antibiotics for UTI in Iran , was very high and increasing (57%-77%).

Comparing this study with other similar studies from Iran, the following results were obtained. In one study, the susceptibility testing on 50 E. coli isolates showed (74%) resistance to ciprofloxacin, approximately the same rate

obtained in this study, and (36%) resistance to gentamicin which was significantly lower than the rate obtained in this study [16]. In a study conducted in northwest of Iran which took over a 14-month period, the results of antimicrobial susceptibility analysis for *E. coli*, as the most prevalent cause of UTI and commonly used antibiotics were: amikacin (97.8%), gentamicin (97%), ciprofloxacin (94%), nitrofurantoin (87.1%), nalidixic acid (93.7%), trimethoprim-sulfamethoxazole (48.2%), that were very different from the results of this study [7]. The resistance pattern of pathogens other than *E. coli* in the current study was also very high, and approximately at the same rate of resistance of *E. coli* except for nitrofurantoin that the resistance of *E. coli* was much lower.

CONCLUSION

Antibiotic resistance of uropathogens for most of the important antibiotics used empirically for UTI was very high and increasing in this hospital compared to the other hospitals Iran and also the other countries. Continuing this trend might confer a crisis in the treatment of UTI patients especially those with complicated UTIs. The results of this study indicated that more attention to antibiotic prescription and antibiotic stewardship in hospitals is urgently needed to prevent the increasing trend of antibiotic resistance and preserving the invaluable present resources of antibiotics.

Conflict of interest

It should be noted that there was no conflict of interest in this article.

ACKNOWLEDGMENT

The present paper was extracted from Information of Infection Control Committee of Razi Therapeutic and Educational Center (Guilan University of Medical Sciences; www.gums.ac.ir). Hereby, the authors would like to thank all the staff who helped in collecting data.

REFERENCES

- [1] Stamm WE, and S. R. Norrby. Urinary tract infections: disease panorama and challenges. . J Infect Dis. 2001 183(Suppl. 1):S1-S4.
- [2] Hooton, TM. Besser, RM. Foxman, B. Fritsche, TR. & Nicolle, LE. Acute uncomplicated cystitis in an era of increasing antibiotic resistance: a proposed approach to empirical therapy. Clin Infect Dis. 2004;39 (1):75-80.
- [3] Gupta, K. Hooton, TM. Stamm, WE. K. Increasing antimicrobial resistance and the management of uncomplicated community-

acquired urinary tract infections. Ann Intern Med. 2001;135 (1):41-50.

- [4] Javier Garau DPN, Björn Wullt, Matteo Bassetti. Antibiotic stewardship challenges in the management of community-acquired infections for prevention of escalating antibiotic resistance. Journal of Global Antimicrobial Resistance. 2014;2(4):245-53.
- [5] Warren, JW. Abrutyn, E. Hebel, JR. Johnson, JR. Schaeffer, AJ. Stamm, WE. Guidelines for antimicrobial treatment of uncomplicated acute bacterial cystitis and acute pyelonephritis in women. Clin Infect Dis. 1999, 29 (4):745-58.
- [6] Kashef GED, N. & Shahbazi, S. Antimicrobial susceptibility patterns of community-acquired uropathogens in Tehran, Iran The Journal of Infection in Developing Countries 2014;4 (4):202-6.
- [7] Farajnia S, Alikhani MY, Ghotaslou R, Naghili B, Nakhilband A. Causative agents and antimicrobial susceptibilities of urinary tract infections in the northwest of Iran. International Journal of Infectious Diseases. 2009;13(2):140-4.
- [8] Lu P-L, Liu Y-C, Toh H-S, Lee Y-L, Liu Y-M, Ho C-M, et al. Epidemiology and antimicrobial susceptibility profiles of Gram-negative bacteria causing urinary tract infections in the Asia-Pacific region: 2009-2010 results from the Study for Monitoring Antimicrobial Resistance Trends (SMART). International Journal of Antimicrobial Agents. 2012;40, Supplement 1(0):S37-S43.
- [9] Mirzarazi M, Rezatofighi SE, Pourmahdi M, Mohajeri MR. Antibiotic Resistance of Isolated Gram Negative Bacteria From Urinary Tract Infections (UTIs) in Isfahan. Jundishapur J Microbiol. 2013;6(8):e6883.
- [10] Schito GC, Naber KG, Botto H, Palou J, Mazzei T, Gualco L, et al. The ARES study: an international survey on the antimicrobial resistance of pathogens involved in uncomplicated urinary tract infections. International Journal of Antimicrobial Agents. 2009;34(5):407-13.
- [11] Zhanel GG, Hisanaga TL, Laing NM, DeCorby MR, Nichol KA, Weshnoweski B, et al. Antibiotic resistance in *Escherichia coli* outpatient urinary isolates: final results from the North American Urinary Tract Infection Collaborative Alliance (NAUTICA). International Journal of Antimicrobial Agents. 2006;27(6):468-75.

- [12] Karlowski j, Kelly l, Thornsberry k, Jones m, Sahm d. Trends in Antimicrobial Resistance among Urinary Tract Infection Isolates of Escherichia coli from Female Outpatients in the United States. *Antimicrob Agents Chemother.* 2002;46 (8):2540-5.
- [13] Jharna Mandal NSA, D. Buddhapriya & Subhash Chandra Parija. Antibiotic resistance pattern among common bacterial uropathogens with a special reference to ciprofloxacin resistant Escherichia coli. *Indian J Med Res.* 2012;136 (5):pp 842-9.
- [14] Bouchillon S HD, Badal R, Hawser S. 2012;. Fluoroquinolone resistance among Gram-negative urinary tract pathogens: global SMART Program results, 2009–2010. . *Open Microbiol J.* 2012;6:74-8. doi: 10.2174/1874285801206010074
- [15] Hoban DJ, Nicolle LE, Hawser S, Bouchillon S, Badal R. Antimicrobial susceptibility of global inpatient urinary tract isolates of Escherichia coli: results from the Study for Monitoring Antimicrobial Resistance Trends (SMART) program: 2009–2010. *Diagnostic Microbiology and Infectious Disease.* 2011;70(4):507-11.
- [16] Kazemnia A, Ahmadi M, Dilmaghani M. Antibiotic Resistance Pattern of Different Escherichia coli Phylogenetic Groups Isolated from Human Urinary Tract Infection and Avian Colibacillosis. *Iranian Biomedical Journal.* 2014;18(4):219-24.