



International Journal of Pharmaceutical and Phytopharmacological Research (eIJPPR)

[Impact Factor – 0.852]

Journal Homepage: www.eijppr.com

Research Article

Development and Validation of UV-Visible Spectrophotometric Method for the Determination of Chloramphenicol in Pure and in its Dosage Form

P.Suguna*, B.Ramachandra and N.V.S.Naidu

Department of Chemistry, S.V.University, Tirupati-517502, A.P., India.

Article info

Article History:

Received 18 December 2014

Accepted 29 December 2014

Keywords:

Spectrophotometry, 1, 10-phenanthroline, Oxidative complexation.

Abstract

A simple, precise, rapid sensitive and accurate spectrophotometric methods have been developed for the estimation of Chloramphenicol UV in pure form and its pharmaceutical formulations based on oxidative complexation reaction UV with 1, 10-Phenanthroline reagent at pH - 4 which is extractable at 510 nm. Beer's law is obeyed in the concentration range 0.5-3 ml (5 to 30 $\mu\text{g ml}^{-1}$). The developed method was applied directly and easily for the analysis of the pharmaceutical formulations. RSD was found to be 0.0185% and recovery 99.73% respectively. The method was completely validated and proven to be rugged. The interferences of the ingredients and excipients were not observed. The repeatability and the performance of the proved method were established by point and internal hypothesis and through recovery studies.

1. INTRODUCTION

Several analytical methods have been reported for the determination of Chloramphenicol in various samples Chloramphenicol (CAP) is 2,2 dichloro- N-[(1R,2R)-2-hydroxy-1-hydroxymethyl-2-(4-nitrophenyl)ethyl]acetamide, C₁₁H₁₂Cl₂N₂O₅, whereas its chemical structure is: Its molecular weight is 323.1 g mol⁻¹. It is a white, greyish-white or yellowish-white, fine crystalline powder or fine crystals, needles or elongated plates, freely soluble in methanol, ethanol, butanol, ethyl acetate, acetone, and in propylene glycol, slightly soluble in water, and ether, insoluble in benzene, and petroleum ether, it melts at 150.5–151.5°C¹.

Chloramphenicol is a bacteriostatic antimicrobial. It is considered a prototypical broad-spectrum antibiotic, alongside the tetracyclines. Chloramphenicol is effective against a wide variety of Gram-positive and Gram-negative bacteria, including most anaerobic organisms. It is widely used because it is inexpensive and readily available². The most serious adverse effect associated with chloramphenicol treatment is bone marrow toxicity, which may occur in two distinct forms: bone marrow suppression, which is a direct toxic effect of the drug and is usually reversible, and aplastic anemia, which is idiosyncratic (rare, unpredictable, and unrelated to dose) and generally fatal. CAP is a non-irritant and is used by local application for the treatment of a variety of infections of the skin, ear and eye including trachoma³. Various methods have been reported for the determination of CAP in pharmaceutical preparations, including HPLC⁴, LC-Mass spectrometry⁵⁻⁷, Polarographic⁸, electrogenerated chemiluminescence⁹, Fluorescent¹⁰, enzymatic method¹¹, colorimetric and spectrophotometric methods¹²⁻¹⁹. The literature is still poor in analytical procedures based on kinetics, especially for drugs in pharmaceuticals or biological fluids. However, some specific advantages in the application of kinetic methods can be expected such as, selectivity due to the measurement of the evolution of the absorbance with the time of the reaction instead of the measurement of absorbance value. Potassium permanganate has been frequently utilized for kinetic measurements in the field of

pharmaceutical analysis. Many pharmaceutical compounds have been determined kinetically through this approach such as tetracycline hydrochloride²⁰, cephalosporins²¹. A norfloxacin²² was determined by its reaction with acetaldehyde and 2,3,5,6 - tetrachloro - 1, 4 -benzoquinone to give a colored product. Ketoprofen²³ was determined kinetically by oxidative coupling reaction of the drug with Hind S. Al-Ward2 3MBTH reagent in the presence of Ce (IV) in acidic medium. Ramipril has also determined kinetically based on the reaction of the carboxylic group of the drug with a mixture of potassium iodate and potassium iodide and the reaction was followed spectrophotometrically²⁴. The empirical formula for Ametocetradin is C₁₁H₁₂Cl₂N₂O₅ and the molecular weight is 323.13 grams. It has the following structure.




Figure-1: Chemical Structure of Ametocetradin

There is however no reported UV- Visible spectrophotometric method for the analysis of Chloramphenicol in its technical grade and formulations. In the present study an attempt has been made to develop simple UV- visible spectrophotometric method for the quantitative determination of Chloramphenicol. Functional group used for color development of Chloramphenicol was primary amine group. The results obtain in this method was based on oxidative coupling reaction with 1, 10-Phenanthroline. An attempt has been made to develop and validate the method to ensure their accuracy, precision, repeatability, reproducibility and other analytical method validation parameters as mentioned in the various guidelines.

2. MATERIALS AND METHODS

2.1 Materials

The pure sample was collected from CIPLA pharmaceuticals. Avalahalli, Vigro agar, Bangalore , 560049.

*Corresponding Author:

P. Suguna,
Department of Chemistry, S.V.University,
Tirupati-517502, A.P., India.
Email: pydalasuguna@gmail.com

2.2 Preparation of standard stock solution

Accurately weighed 100 mg of Chloramphenicol was dissolved in 40 ml of methanol in 100 ml volumetric flask and volume was made up to the mark with methanol. i.e. $1000 \mu\text{g ml}^{-1}$ (Stock solution A). From the above stock solution A 10 ml of solution was pipette out into 100 ml volumetric flask and the volume was made up to the mark with methanol to obtain the final concentration of $100 \mu\text{g ml}^{-1}$ (Stock solution B).

2.3 Preparation of calibration curve

Fresh aliquots of Chloramphenicol ranging from 0.5 to 3 ml were transferred into a series of 10ml volumetric flasks to provide final concentration range of 5 to $30 \mu\text{g ml}^{-1}$. To each flask 1ml of (0.01M) 1, 10-phenanthroline solution was added followed by 1ml of (0.2%) ferric chloride solution and resulting solution was heated and finally 1ml (0.2M) orthophosphoric acid solution was added. The solutions were cooled at room temperature and made up to mark with distilled water. The absorbance of orange red colored chromogen was measured at 510 nm against the reagent blank. The color species was stable for 24h. The amount of Chloramphenicol present in the sample solution was computed from its calibration curve.

2.4 Procedure for formulations

Twenty tablets containing Chloramphenicol were weighed and finely powdered. An accurately weighed portion of the powder equivalent to 100 mg of Chloramphenicol was dissolved in a 100 ml of methanol and mixed for about 5 min and then filtered. The methanol was evaporated to dryness. The remaining portion of solution was diluted in a 100 ml volumetric flask to the volume with methanol up to 100 ml to get the stock solution A. 10 ml of aliquots was pipette out into 100 ml volumetric flask and the volume was made up to the mark with methanol to obtain the final concentration of $100 \mu\text{g ml}^{-1}$ (Stock solution B). Subsequent dilutions of this solution were made with methanol to get concentration of 5-30 $\mu\text{g ml}^{-1}$ and were prepared as above and analyzed at the selected wavelength, 510 nm and the results were statistically validated.

2.5 Procedure for blood sample

After collection of blood sample, it will be centrifuged. For isolation of Chloramphenicol from plasma sample, Methanol was used for protein precipitation. Liquid- Liquid extraction was performed with plasma by alkalization with 1M NaOH, followed by extraction with 30% dichloromethane in Hexane. The upper organic layer was evaporated to dryness, and dry residue 100 mg was dissolved in 100 ml of Methanol ($1000 \mu\text{g ml}^{-1}$). From the above solution 10 ml is taken into a 100 ml of volumetric flask and made up to the mark with methanol ($100 \mu\text{g ml}^{-1}$).

From the above solution ranging from 1-6 ($10-60 \mu\text{g ml}^{-1}$) were transferred in to 10 ml volumetric flask and to the each flask 1ml of (0.01M) 1,10- Phenanthroline solution was added followed by 1ml of (0.2%) ferric chloride solution and made up to the mark with methanol. Then the resulting solution was heated for 15 min and finally 1ml (0.2M) orthophosphoric acid solution was added. The solutions were cooled at room temperature and made up to the mark with distilled water. The absorbance of orange red colored chromogen was measured at 510 nm against the reagent blank. The color species was stable for 24 h. The amount of Chloramphenicol present in the sample solution was computed from its calibration curve.

3. RESULTS AND DISCUSSIONS

3.1 Optical parameters

In order to ascertain the optimum wavelength of maximum absorption (λ_{max}) formed in UV-visible spectrophotometric method (Reference method – A) and of the colored species formed in each of the four visible spectrophotometric methods, specified amount of Chloramphenicol in final solution $10 \mu\text{g ml}^{-1}$ (method A), $5 \mu\text{g ml}^{-1}$ of this method were taken and the colors were developed following the above mentioned procedures individually. The absorption spectra were scanned on spectrophotometer in the wavelength region of 200-400nm (for method A) and 380-800 nm for this Method against corresponding reagent blanks. The reagent blank absorption spectrum of each method was also recorded against distilled water /methanol. The results are graphically represented in (fig- 2).




Fig 2: Absorption spectrum of Chloramphenicol with 1, 10-Phenanthroline/FeCl₃

3.2 Parameters fixation

In developing these methods, a systematic study of the effects of various relevant parameters in the methods concerned were under taken by verifying one parameter at a time and controlling all other parameter to get the maximum color development (for this method), reproducibility and reasonable period of stability of final colored species formed. The following studies were conducted.




Fig 3: Beer's law plot of Chloramphenicol with 1, 10-Phenanthroline/FeCl₃




Fig 4: Beer's law plot for Chloramphenicol in blood sample

3.3 Method

The results obtained in this method were based on oxidation followed by complex formation reaction of Chloramphenicol with 1,10-phenanthroline, ferric chloride and orthophosphoric acid to form an orange red colored chromogen that exhibited maximum absorption at 510 nm against the corresponding reagent blank. The functional group used for the color development for this method was primary amine group. A schematic reaction mechanism of Chloramphenicol with 1, 10-Phenanthroline reagent was shown in (fig-5). The effect of various parameters such as concentration and volume of 1, 10- Phenanthroline and strength of acid order of addition of reagents, solvent for final dilution were studied by means of control experiments varying one parameter at a time.

3.4 Optical characteristics

The reference method adhere to beer's law the absorbance at appropriate wave length of a set of solutions contains different amounts of Chloramphenicol and specified amount of reagents (as described in the recommended procedure) were noted against appropriate reagent blank. Least square regression analysis was carried out for the slope. Intercept and correlation coefficient, Beer's law limits, molar absorptivity and sandells sensitivity for Chloramphenicol with each of mentioned reagents was calculated. In order to test whether the colored species formed in the method adhere the beer's law the absorbance at appropriate wavelength of a set of solutions contain different amounts of Chloramphenicol and specified amount of reagents (as described in the recommended procedure) were noted against appropriate reagent blanks or distilled water. Least square regression analysis was carried out for the slope, intercept and correlation coefficient, beer's law limits molar absorptivity Sandells sensitivity for Chloramphenicol with each of mentioned reagents were calculated.




Fig 5: A Schematic reaction mechanism of Chloramphenicol with 1,10-Phenanthroline

Table 1: Optical characteristics and precision by (1, 10-PT)

Parameter	Visible method
Color	Orange red
Absorption maxima(nm)	510
Beer's law limits ($\mu\text{g ml}^{-1}$)	5-30
Molar absorptivity ($\text{l mol}^{-1} \text{cm}^{-1}$)	1.01486×10^4
Sandell's Sensitivity ($\mu\text{g cm}^{-2}$)	0.0318
Regression equation (Y^*)	
Slope (b)	0.0309
Intercept(a)	0.0019
Standard deviation(SD)	0.0001
Correlation coefficient (r^2)	0.9999
%RSD (Relative standard deviation)*	0.0185
Range of errors	
Confidence limits with 0.05 level	8.0015
Confidence limits with 0.01 level	0.0001
Limits of detection (LOD)($\mu\text{g ml}^{-1}$)	9.7087×10^{-3}
Limits of quantification (LOQ) ($\mu\text{g ml}^{-1}$)	0.03236

* RSD of six independent determinations.

3.5 Precision

The precision of each one among the five proposed spectrophotometric methods were ascertained separately from the absorbance values obtained by actual determination of a fixed amount of Chloramphenicol $10 \mu\text{g ml}^{-1}$ in final solution. The percent relative standard deviation and percent range of error (at 0.05 and 0.01 confidence limits) were calculated for the proposed methods and presented in Table-1.

3.6 Analysis of formulations

Commercial formulations of Chloramphenicol were successfully analyzed by the proposed methods. The values obtained from the proposed and reference methods were compared statistically by the t and F tests and were found that those proposed methods do not differ significantly from the reported methods and they were presented in Table -2. The proposed methods also applied for Biological Samples (Blood) for good recoveries are obtained which were recorded in Table - 7.

3.7 Accuracy

Recovery studies were carried by applying the method to drugs sample present in formulations to which known amount of Chloramphenicol of label claim was added (standard addition method). The recovery studies were carried by applying the method

to biological sample (Blood) to which known amount of Chloramphenicol corresponds to 2 mg formulations taken by the patient. By the follow of standard addition method 2 mg of label claim was added. After the addition of these standards the contents were transferred to 100 ml volumetric flask and dissolved in solvent. Finally the volume was made up to the mark with solvent. The solution was filtered through Whatman No. 41 filter paper. The mixed sample solutions were analyzed and their absorbance value was determined. At each level of recovery five determinations were performed and present in Table - 3. The results obtained were compared with expected results and were statistically validated in Table - 4.

3.8 Linearity and Range

The linearity of analytical method is its ability to elicit test results that are directly proportional to the concentration of analyze in sample with in a given range. The range of analytical method is the interval between the upper and lower levels of analyze that have been demonstrated within a suitable level of precision, accuracy and linearity.

3.9 Specificity and Selectivity

Specificity is a procedure to detect quantitatively the analyze in the presence of components that may be expected to present in the sample matrix. While selectivity is a procedure to detect the analyze qualitatively in presence of components that may be expected to present in the sample matrix. The excipient in formulations was spiked in a pre weighed quantity of drugs and then absorbance was measured and calculations were done to determine the quantity of the drugs.

3.10 Repeatability

Standard solutions of Chloramphenicol were prepared and absorbance was measured against the solvent as the blank. The observance of the same concentration solution was measure five times and standard deviation was calculated and presented in Tables - 7 and 9.

3.11 Interferences Studies

The effect of wide range of inactive, ingredients usually present in the formulations for the assay of Chloramphenicol under optimum conditions was investigated. None of them interfered in the proposed methods even when they are present in excess fold than anticipated in formulations.

3.12 Solution Stability

The stability of the solutions under study was established by keeping the solution at room temperature for 48 Hours. The results indicate no significant change in assay values indicating stability of Drug in the solvent used during analysis. The results are recorded in Table -6.

Table 2: Assay results of Chloramphenicol in formulations by UV-visible method

Name of the Formulation	Formulation in (mg)	Amount found by the proposed method(mg)	Amount found by the reference Method ^{40,41} (mg)	% Recovery
Ocupol-D	250	247.5 $t=0.0049^*$ $F=8.0606^*$	248.19	99.47
Phenicol	250	248.75 $t=0.0048^*$ $F=8.0505^*$	247.98	99.68

*t and F- values refer to comparison of the proposed method with reference method.

*Theoretical values at 95% confidence limits $t=0.0029$ and $F=6.5594$.

Table 3: Determination of accuracy of Chloramphenicol

Amount of CP in formulation (mg)	Amount of Standard CP added (mg)	Total amount found (mg)	% Recovery
248.66	200	447.58	99.46
247.5	200	445.5	99.00
245.66	200	442.18	98.26
249.33	250	498.66	99.73
248.75	250	497.5	99.5
247.83	250	495.66	99.13
249.44	300	548.76	99.77
248.95	300	547.69	99.58
248.19	300	546.01	99.27

Table 4: Statistical data for accuracy determination

Total amount found (mean)	Standard deviation	% RSD
247.27	1.5127	0.6117
248.63	0.7563	0.3041
248.86	0.6298	0.2530

The results are the mean of five readings at each level of recovered

Table 5: Repeatability data for CP at 510 nm

Conc. ($\mu\text{g ml}^{-1}$)	Abs 1	Abs2	Abs3	Mean	Std. deviation	(%) RSD*
5	0.154	0.155	0.153	0.154	0.0014	0.9090
10	0.309	0.308	0.307	0.308	0.001	0.324
15	0.458	0.457	0.456	0.457	0.001	0.2188
20	0.615	0.614	0.617	0.615	0.0015	0.2439
25	0.772	0.774	0.775	0.773	0.0015	0.194
30	0.927	0.928	0.926	0.927	0.001	0.1078

*RSD of six independent determinations.

Table 6: Color stability data for 1, 10-Phenanthroline Method

Conc. in $\mu\text{g ml}^{-1}$	Time in Hours							
	4	8	12	16	20	24	28	32
25	0.773	0.774	0.774	0.775	0.775	0.775	0.675	0.512

Table 7: Assay results of Chloramphenicol in Blood sample

Name of the Formulation	Formulation in (mg)	Amount found by the proposed method in (mg)	Amount found by the reference method ^{40,41} (mg)	% Recovery
Ocupol-D	5	3.98 $t=0.0029^*$ $F=9.9855^*$	3.88	97.42
Phenicol	5	3.97 $t=0.0028^*$ $F=9.9866^*$	3.87	97.41

* t and F values refer to comparison of the proposed method with reference method.

*Theoretical values at 95% confidence limits $t=0.00196$ and $F=9.7866$.

Table 8: Determination of accuracy of Chloramphenicol

Name of the Formulation in (mg)	Amount of Drug in Blood sample (mg)	Amount of Standard Drug added in (mg)	Total amount found (mg)	% Recovery
5	3.99	5	7.98	79.80
5	3.98	5	7.99	79.90

The results are the mean of five readings at each level of recovered

Table 9: Repeatability data for Chloramphenicol at 510nm

Concentration in ($\mu\text{g ml}^{-1}$)	Abs1	Abs2	Abs3	Mean	Std. Deviation	(%) RSD*
10	0.0246	0.0247	0.0246	0.0246	0.0001	0.4065
20	0.0493	0.0494	0.0492	0.0493	0.0001	0.2028
30	0.074	0.075	0.078	0.0756	0.0002	0.2645
40	0.0987	0.0986	0.0988	0.0987	0.0001	0.1013
50	0.1234	0.1235	0.1236	0.1235	0.0001	0.08097
60	0.1481	0.1483	0.1482	0.1482	0.0001	0.0674

*RSD of six independent determinations.

4. CONCLUSION

The method was found to be accurate and precise, as indicated by recovery studies close to 100 and % RSD is not more than 2. The summary of validation parameters of proposed UV- Visible method is given. The simple, accurate and precise UV- Visible method for the determination of Chloramphenicol as bulk, Commercial samples and Blood samples has been developed. The method may be recommended for routine and quality control analysis of the investigated pure in bulk and samples. The analytical solution is found to be stable up to 48 Hrs at room temperature. Hence, it is concluded that the analytical method is validated and can be used for routine analysis and for stability study.

ACKNOWLEDGEMENTS

The authors are grateful to S.V.University for providing the laboratory Facilities and the pure sample was collected from CIPLA pharmaceuticals.

REFERENCES

- "British Pharmacopoeia on CD-Rom" The Stationery Office on behalf of the Medicines and Healthcare products Regulatory Agency (MHRA). London. 5th., ed., 2007.
- Falagas, M. E.; Michalopoulos, A. A.; "Potential of old-generation antibiotics to address current need for new antibiotics"; *Expert Rev Anti Infect Ther.*, 6, 593–600, 2008.
- Wilson, A.; Schild, H. O.; Modell, W.; "Applied Pharmacology"; 11th Ed., Churchill Livingstone, London, 1975.
- Pan, Y.; Xu, Q.; Kang, X.; Zhang, J.; "Determination of chloramphenicol residues in milk by reversed-phase high performance liquid chromatography with fluorescence detection"; 23, 577-580, 2005.
- Storey, j.; Pfenning, A.; Turnipseed, S.; Nandrea, G.; Rebecca, L.; Burns, C.; Madson, M.; "Determination of Chloramphenicol Residues in Shrimp and Crab Tissues by Electrospray Triple Quadrupole LC/MS/MS"; 19, 1-16, 2003.
- Jiang, Y.; Zhong, X.; Zhong, T.; Shen, C.Y.; Ding, T.; Chen, H. L.; Wu, B.; Shen, W. J.; "Determination of chloramphenicol in royal jelly by liquid chromatography/ tandem mass spectrometry", *veterinary drug residues, Journal of AOAC International*; 47, 3464–3469, 2006.
- Teresa, K.; "Determination of Chloramphenicol in Feed Use of High - Performance Liquid Chromatographic - Mass Spectrometry", *Branch Institute of Animal Drugs Inspection*, 2003.
- Summa, A. F.; "Polarographic determination of chloramphenicol preparations", *J. Pharm. Sci.*; 54, 442-444, 1965.
- Lindino C. A.; Bulhões, L.O. S.; "Determination of chloramphenicol in tablets by electrogenerated chemiluminescence"; *J. Braz. Chem. Soc.*, 15, 876- 880, 2004.
- Haughland, P.; Kang, R.; Young, H.; Steven, L.; Melner, M.; "Fluorescent chloramphenicol derivatives for determination of chloramphenicol acetyltransferase activity", *Molecular Probes*, 18, 722- 730, 1991.

11. Morris, H.C.; Miller, J.; Campbell, L. S.; Hammond, P. M.; Berry, D. J.; Price, C. P.; "A rapid, enzymatic method for the determination of chloramphenicol in serum"; *Journal of Antimicrobial Chemotherapy*, 22, 935-944, 1988.
12. Chukwuenweniwe, J. E.; Johnson, S.; Adelusi, S. A.; "An alternative colorimetric method for the determination of chloramphenicol"; *Tropical Journal of Pharmaceutical Research*, 2, 215-221, 2003.
13. Wahbi, A. M.; Abdine, H.; Korany, M.A.;El-Yazbi, F. A.; "Spectrophotometric determination of chloramphenicol-sulphacetamide in eye drops"; *Pharmazie*, 33, 721-722, 1978.*Journal of Al-Nahrain University Vol.15 (4)*, December, 2012, pp.22-30
14. Naik, S.; Nagaraja, P.; Yathirajan, H.; Hemanthakumar; Mohan, H.; "New spectrophotometric methods for the quantitative determination of chloramphenicol in pharmaceuticals" , *Pharmaceutical Chemistry Journal*, 40, 576-581, 2006.
15. Shah, R. C.; Raman, P.V.; Shah, B. M.; "Spectrophotometric determination of chloramphenicol and tetracycline hydrochloride in mixtures" *Journal of Pharmaceutical Sciences*, 52, 167-168, 1963.
16. Freeman, F. M.; "the colorimetric determination of chloramphenicol", *J. Chem. Soc.*, 81, 298-299, 1956.
17. Mahrous, M. S.; Abdel-Khalek, M. M.; "Spectrophotometric determination of phenothiazines, tetracyclines and chloramphenicol with sodium cobaltinitrite", *Talanta*, 30, 792-794, 1983.
18. Moț, A.; Soponar, C.; Medvedovici, F.; Sarbu, A.; "Simultaneous Spectro-photometric Determination of Aspirin, Paracetamol, Caffeine, and Chlorphenamine from Pharmaceutical Formulations Using Multivariate Regression-Methods" *World Wide Science. org*, 2010.
19. Al-Sabha, T. N.; Rasheed, B. A.;" Spectrophotometric Method for Determination of Chloramphenicol in Pharmaceutical Preparations using 1,2-Naphthoquinone-4-Sulphonate as a Chromogenic Reagent", *JJC*, 5, 201-210, 2010.
20. Ahmidaa, N. H. S.; El-Hashemea, F.; El-Enany N.; Belal, F.; "Kinetic spectro-photometric method for the determination of tetracycline hydrochloride in pharmaceutical formulations", *Applied Science Research*, 1, 1-11, 2009.
21. Omar, M. A.; Abdelmageed, O. H.; Attia, T. Z.; "Kinatic spectrophotometric determination of certain cephalosporins in pharmaceutical formulations", *International Journal of Analytical Chemistry*, 5, 12-15, 2009.
22. Darwish, I. A.; Sultan, M. A.; Al-Arfaji, H. A.; "Novel selective kinetic spectrophotometric method for the determination of norfloxacin in its pharmaceuticals formulations", *Talanta*, 78, 1383-1388, 2009.
23. El-Brashy, A.; Eid, M.; Talaat, W.; "Kinetic spectrophotometric method for the determination of Ketoprofen in pharmaceuticals and biological fluids, *International Journal of Biomedical Science*, 2, 405-412, 2006.
24. Rahman, N.; Ahmad, Y.; Najmul S.; Azmi, H.; " Kinetic Spectrophotometric Method for the Determination of Ramipril in Pharmaceutical Formulations", *AAPS Pharm Sci Tech.*, 6, 543-551, 2006.