



International Journal of Pharmaceutical and Phytopharmacological Research (eIJPPR)

[Impact Factor – 0.852]

Journal Homepage: www.eijppr.com

Research Article

Article ID: 391

GC-MS analysis of the ethanol extract of *Biophytum sensitivum* (L.) DC (Oxalidaceae)

P. Devi Raja Rajeswari^{*1}, K. Aruna¹, K. Prabu², R. Chidambaram³ and S. Raja Sankar⁴

¹Department of Anatomy, Rajas Dental College, Kavalkinaru, Tirunelveli and ⁵Bharath University, Chennai, India.

² Sree Balaji Medical College & Hospital, Chennai, India.

³Sri Lakshmi Narayana Institute of Medical Sciences, Puducherry.

⁴Velammal Medical College Hospital & Research Institute, Madurai, India.

*Corresponding Author: Email: saran_gan@rediffmail.com

Article info

Article History:
Received 1 June 2015
Accepted 29 June 2015

Keywords:

Ruminal artery, Nitric oxide, sGC,
cGMP.

Abstract

The medicinal plant *Biophytum sensitivum* (L.) DC possess a wide spectrum of medicinal properties and are therefore used in the treatment of for asthma, snakebites, stomachalgia and phthisis, positive effects in inflammatory diseases, anti-tumor activities and antioxidant activity. In the present study, ethanol extracts from *Biophytum sensitivum* were subjected to GC-MS analysis to study the important phytochemical constituents responsible for the above reported pharmacological activities. The crude extracts of ethanol were obtained by soxlet method. The GC-MS analysis of ethanol extract from *Biophytum sensitivum* revealed the presence of eighteen phytocompounds. The compounds were identified by comparing their retention time and peak area with that of literature and by interpretation of mass spectra. The most prevailing major compounds were 9, 12-Octadecadienoic acid (Z,Z) (50.36%), n-Hexadecanoic acid (17.42%), 9,12-Octadecadienoic acid, methyl ester (E,E) (6.34%) and 2-Pentadecane,6,10,14-trimethyl- (3.31%).

1. INTRODUCTION

The use of local plants in folk medical practices has a long history. The resource base of the traditional medical practices prevalent in rural and tribal villages of India and abroad is mainly the plants. Medicinal plants are used to maintain and promote healthy life, prevent disease and cure ailments. It has been estimated that even today, 80% of the world population rely on herbal traditional medicine for their primary health care. Traditional knowledge of medicinal plants has always guided the search for new cures. In spite of the advent of modern high throughput drug discovery and screening techniques, traditional knowledge systems have given clues to the discovery of valuable drugs¹. Standardization is an essential measure of quality, purity and authenticity. The standardization of crude drug is an integral part for establishing its correct identity. Also, WHO has emphasized the need to ensure the quality of medicinal plants products using modern controlled technique and applying suitable standards². Gas chromatography-mass spectrometry (GC-MS) is the best technique to identify the bioactive constituents of long chain hydrocarbons, alcohols, acids, esters, alkaloids, steroids, amino acid and nitro compounds^{3, 4, 5}. Therefore, characterization of extracts of medicinal plants is necessary due to its numerous benefits to science and society. Review of literature divulges that information on the GC-MS analysis of *Biophytum sensitivum* (L.) DC belonging to the family Oxalidaceae is totally lacking. Hence, the objective of the present study is to identify the phytochemical constituents with the aid of GC-MS technique. This work will help to identify the compounds of therapeutic value.

Pharmacologically, the *Biophytum* plant has been investigated for its hypoglycemic⁶, hypocholesterolemic⁷, and anti-cancer effect⁸. It is a known traditional remedy for the treatment of diabetes⁹ and anti-tumor activities¹⁰.

1.1 Plant description

The little plant grows up to maximum of 20 cm and possess unbranched woody erect stem. Leaves abruptly pinnate, leaflets opposite, 6 to 12 pairs, and each leaflet is up to 1.5 cm long, the terminal pair is the largest. The flowers are many and crowded at the apices of the numerous peduncles, normally yellow, white, or orange with red streak in the center of each of the five petals. The sepals are subulate-lanceolate, striate, and about 7 mm long. Fruits are ellipsoid capsules which are shorter than the persistent calyx¹¹.

2. MATERIALS AND METHODS

2.1 Plant material

The medicinal plant *Biophytum sensitivum* (L.) DC was collected from Tirunelveli District, Tamil Nadu, India. The identified plant species was confirmed with Voucher specimen available in the Survey of Medicinal Plant Unit (SMP), Govt. Siddha Medical College, Palayamkottai, Tirunelveli, Tamil Nadu. The taxonomic features of the plant confirmed with the Flora of Presidency of Madras¹² and The Flora Tamil Nadu Carnatic¹³.

2.2 Soxhlet extraction

About 60 g dried sample was refluxed with 250 ml of the ethanol for 5 hour on a steam bath. The extract was collected and concentrated.

2.2.1 Procedure

The GC - MS analyses were carried out in a Shimadzu GC – MS - QP 2010 gas chromatograph fitted with a DB1 (methylphenylsiloxane, 30 m × 0.25 mm i.d.) capillary column. Carrier gas, helium with a flow rate of 1 ml/min; column oven temperature 70°C, 5 min in 180°C, 180-260°C at 3°C/min, 5 min in 260°C, 260-280°C at 0.2°C/min, and finally 5 min in 280°C; injector temperature, 280°C detector temperature, 290°C, volume injected, 2 µL: Split ratio, 10:1. The MS operating parameters were as follows: ionization potential 70 eV; ion source temperature 200°C; quadrupole 100°C, solvent delay 6.0 min, scan speed 2000 amu/s, total MS running time 36min. and scan range 30-600 amu, eV voltage 3000 volts.

The concentrated extract is injected into the GC/MS instrument (Hewlett Packard 5890 GC/MS with Mass Selective Detector with Turbo mass gold-perkin Elmer). The sample is volatilized at the injection port and eluted through a capillary column under increasing temperature. As the sample moves through the column, various components are separated due to their affinity for the stationary phase of the column and can be identified by retention time (the time it takes for a compound to pass through the column and gas chromatograph system). Each chemical component in a sample has a distinct retention time measured in minutes, shown in a peak on a graph which measures abundance on the ordinate against retention time on the abscissa. The integrated peak is correlated to the concentration of the chemical. A mass selective detector breaks up each chromatographic component into fragment ions, which are shown by their abundance, with each ion represented as a vertical line in increasing molecular weight. The height of each line corresponds to the abundance of that ion. The resulting mass spectrum is unique to that chemical. This mass spectrum forms a “fingerprint” that can identify the compound by a computer search of mass spectra. A computer search of the mass spectra corresponding to all the chromatographic peaks for a sample should yield a statistical match for nicotine at a 12.9 min retention time value if they were present two modes of GC/MS were possible with this instrumental method. First, there is a “Scan” mode which looks at all the constituents of a sample, listing whatever chemical components are present.

2.2.2 Compound Identification

Components of the methnolic extracts were identified by comparison of their mass spectra and retention indices with those published in the literature and contained in the NIST 2005 MS computer library (Wiley).


3. RESULTS

The bioactive compounds present in the ethanolic extract of leaf of *Biophytum sensitivum* were identified by GC-MS analysis (Figure 1). Eighteen compounds were detected in the ethanolic extract of *Biophytum sensitivum*. The active principles with their retention time, molecular formula, molecular weight and concentration (%) in the ethanol extracts of leaf of *B. sensitivum* are presented in Table 1, and the total running time was 36 min. The spectra of the compounds were matched with Wiley 9.0 and National Institute of Standards and Technology libraries. The most prevailing major compounds were 9,12- Octadecadienoic acid (Z,Z) (50.36%), n-Hexadecanoic acid (17.42%), 9,12-Octadecadienoic acid, methyl ester (E,E) (6.34%) (Fig: 2), 2-Pentadecane,6,10,14-trimethyl- (3.31%), n-hexadec Trans-3-Caren-2ol (3.16%), etc., and minor components were 2-Cyclopentene -1-undecanoic acid(+) (0.39%), Oleic acid (0.99%), and 9-Octadecenal (0.70 %). 9,12- Octadecadienoic acid is a unsaturated fatty acid, occurring widely in plant glycosides. It is an essential fatty acid in mammalian nutrition and is used in the biosynthesis of prostaglandins and cell membranes. It is commonly called as Linoleic acid. It is an essential fatty acid that must be consumed

for proper health. It can be used to show the antioxidant effect of natural phenols. Unsaturated fatty acids are important to every cell in the body for normal growth, especially of the blood vessels and nerves and to keep the skin and other tissues youthful and supple through their lubricating quality¹⁴. Linolenic acid possesses anti-inflammatory, insectifuge, hypocholesterolemic, cancer preventive, nematicide, hepatoprotective, antihistaminic, antieczemic, antiacne, 5-alpha reductase inhibitor, antiandrogenic, antiarthritic and anticoronary properties¹⁵. n-Hexadecanoic acid can be an antioxidant, hypocholesterolemic, nematicide, pesticide, lubricant activities and hemolytic 5-alpha is a reductase inhibitors^{16, 17}. *Euphorbia longan* leaves mainly contained n-hexadecanoic acid and 9, 12-Octadecadienoic acid. Hexadecanoic acid has earlier been reported as a component in alcohol extract of the leaves of *Kigelia pinnata*¹⁸ and *Melissa officinalis*¹⁹. 17 compounds with n-Hexadecanoic acid and Octadecanoic acid as the major compounds in the leaves of *Cleistanthus collinus*²⁰. GC-MS analysis of ethyl acetate extract of *Goniothalamus umbrosus* revealed the presence of n-Hexadecanoic acid²¹.

Table: 1 GC MS analysis of *Biophytum sensitivum*

No.	RT	Name of the compound	Molecular formula	Molecular Weight	Peak area %
1.	2.06	À-Pinene	C ₁₀ H ₁₆	136	1.72
2.	2.32	Trans-3-Carene-2ol	C ₁₀ H ₁₆ O	152	3.16
3.	2.52	1,4-Cyclohexadiene, 1-methyl-1-4-(1-methylethyl)-(q_ terpinen)	C ₁₀ H ₁₆	136	1.27
4.	11.05	7-Octen-1-ol, 3,7-dimethyl-(S) – (à-Citronellol)	C ₁₀ H ₂₀ O	156	2.05
5.	11.15	2-Pentadecane,6,10,14-trimethyl-	C ₁₈ H ₃₆ O	268	3.31
6.	11.37	E-7-Tetradecenol	C ₁₄ H ₂₈ O	201	0.54
7.	12.79	n-Hexadecanoic acid	C ₁₆ H ₃₂ O ₂	256	17.42
8.	14.02	2-Cyclopentene -1-undecanoic acid(+)	C ₁₆ H ₂₈ O ₂	252	0.39
9.	14.85	9,12-Octadecadienoic acid, methylester (E,E)	C ₁₉ H ₃₄ O ₂	294	6.34
10.	15.27	9,12- Octadecadienoic acid (Z,Z)	C ₁₈ H ₃₂ O ₂	280	50.36
11.	19.39	7-Hexadecenal(Z)-	C ₁₆ H ₃₀ O	238	1.11
12.	19.97	9-Octadecenal	C ₁₈ H ₃₄ O	266	0.70
13.	22.17	3-Hexadecyloxycarbonyl 1-5-(2-hydroxyethyl) -4-methylimidazolium ion	C ₂₄ H ₄₅ N ₂ O ₃	409	1.88
14.	23.71	2,6,10-Decacatrien-1-ol,3,7,1,-trimethyl,(E,E)-	C ₁₅ H ₂₆ O	222	2.14
15.	24.69	Pentanoic acid, 10-undecyryl ester	C ₁₆ H ₃₀ O ₂	254	1.42
16.	27.27	Oleic acid	C ₁₈ H ₃₄ O ₂	282	0.99
17.	27.40	1-Nonadecanol	C ₁₉ H ₄₀ O	284	2.77
18.	28.00	Vitamin E	C ₂₉ H ₅₀ O ₂	430	2.44

Fig: 1 Chromatogram of *Biophytum sensitivum* ethanol extract




Fig 2: Structure of major compounds

4. CONCLUSION

In the ethanol extracts of plant *Biophytum sensitivum* (Oxalidaceae) eighteen compounds have identified by (GC) and Mass Spectroscopy (MS) method. Components of the methnolic extracts were identified by comparison of their mass spectra and retention indices with those published in the literature and contained in the NIST '2005 MS computer library (Wiley). 9, 12, - Octadecatrienoic acid (Z,Z) and n-Hexadecanoic acid were the most abundant of fatty acid identified in the fatty acid fraction. Further experiments, are planned to establish the influence of the components of these mixtures on the pharmacological activity.

5. REFERENCES

1. Badugux LR. Phytochemical Screening, Quantitative Estimation Total Phenolics and Total Flavonoids, Anti-Microbial Evaluation of *Cyamopsis tetragonoloba*. International Journal of Research in Pharmaceutical and Biomedical Sciences 2012; 3(3):1139-1142. [\[Google Scholar\]](#)
2. Kaushik S, Sharma P, Jain A, Sikarwar SM. Preliminary phytochemical screening and HPTLC fingerprinting of *Nicotiana tabacum* leaf. J. Pharm. Res. 2010, 3:1144–1145.
3. Subramanian S, Ramakrishnan N. Chromatographic fingerprint analysis of *Naringi crenulata* by HPTLC technique. Asian Pac. J. Trop. Biomed. 2011, 1(Suppl. 2):S195–S198
4. Muthulakshmi A, Joshibhi-Margret R, Mohan VR. GC-MS analysis of bioactive components of *Feronia elephantum* Correa (Rutaceae). J. Appl. Pharm. Sci; 2012:2-69. [\[Google Scholar\]](#)
5. Yamunadevi M, Wesely EG, Johnson M. Chromatographic finger print analysis of steroids in *Aerva lanasa* L. by HPTLC technique. Asian Pac. J. Trop. Biomed. 2011, 1(6):428–433
6. Puri D, Baral N J Physiol Pharmacol. Hypoglycemic effect of *Biophytum sensitivum* in alloxan diabetic rabbits. Indian 1998; 42:401-6. [\[Google Scholar\]](#)
7. Puri D Pharm Biol. Hypocholesterolemic Effect of *Biophytum sensitivum* Leaf water extract. 2003; 41:253-8. [\[Google Scholar\]](#)
8. Guruvayoorappan C, Girija K J Med Res. Effect of *Biophytum sensitivum* on B16 - 10 melanoma cells induced metastasis. Indian 2005; 13(3):56-9. Available from: <http://toxnet.nlm.nih.gov/cgi-bin/sis/search2/r?dbs+hsdb:@term+@rn+298-93-1> PubMed PMID: 17552364. [\[Google Scholar\]](#)
9. Puri D, Baral N, Assoc UBM. Indigenous plant remedies in Nepal used in heart diseases. J Nepal 1997; 36:334-7. [\[Google Scholar\]](#)
10. Bhaskar VH, Rajyalakshmi V. Anti- tumor activity of aqueous extract of *Biophytum sensitivum* Linn. Annals of biological research. 2010; 1:76-80. [\[Google Scholar\]](#)
11. Kirtikar KR, Basu BD. Indian medicinal plants; Dehradun, India: International book distributors, 2005; p. 30
12. Gamble JS. *Oxalidaceae* In: Flora of the Presidency of Madras, Bishen Singh Mahandra Pal Singh Dehra Dun 1915 – 1921.
13. Matthew KM. The Flora of Tamil Nadu Carnatic Vol I, The Rapinat Herbarium, Tiruchirappali. Vol - I 1983. [\[Google Scholar\]](#)
14. Igwe OU. *Donatus Ebera Okwu*. GC-MS evaluation of bioactive compounds and antibacterial activity of the oil fraction from the seeds of *Brachystegia eurycoma*. HARMS) Asian Journal of Plant Science and Research; 2013:3-2. [\[Google Scholar\]](#)
15. Sermakkani M, Thangapandian V. GC-MS Analysis of *Cassia italica* leaf methanol extract Asian. Journal of Pharmaceutical and Clinical Research 2012; 5(2). [\[Google Scholar\]](#)
16. Praveen Kumar P, Kumaravel S, Lalitha C. Screening of antioxidant activity, total phenolics and GC-MS study of *Vitex negundo*. Afr. J. Biochemistry Res. 2010; 4(7):191-195. [\[Google Scholar\]](#)
17. Devi P, Nagarajan M, Christina AJM, Meera R, Merlin NJ. GC-MS analysis of *Euphorbia longan* leaves Int. J. of Pharmaceutical Res and Development 2009; 2009(8):1-4. [\[Google Scholar\]](#)
18. Grace OM, Light ME, Lindsey KL, Moholland DA, Staden JV, Jader AK. Antibacterial activity and isolation of antibacterial compounds from fruit of the traditional African medicinal plant, *Kigelia africana*. 2002; 68:220-222. [\[Google Scholar\]](#)

19. Sharafzadeh S, Khosh-Khui M, Javidnia K. Aroma profile of leaf and stem of Lemon Balm (*Melissa Officinalis* L.) grown under Greenhouse Conditions. *Advan. Environmental Biol.* 2011 2011; 5(4):547-550. [\[Google Scholar\]](#)
20. Parasuraman S, Raveendran R, Madhavrao C. GC-MS analysis of leaf extracts of *Cleistanthus collinus* Roxb. (Euphorbiaceae). *Int. J. Ph. Sci.* 2009. 1(2):284-286.
21. Siddiq Ibrahim A., Ahmad Bustamam A, Manal Mohammed E. Syam MI Mohamed Yousif M, Abdelbasit Adam, Alhaj NA, Rasedee Abdullah. GC-MS determination of bioactive components and antibacterial properties of *Goniothalamus umbrosus* extracts. *Afr. J. Biotech* 2009 2009; 8(14):3336-3340. [\[Google Scholar\]](#)