

ISSN (Online) 2249-6084 (Print) 2250-1029

International Journal of Pharmaceutical and Phytopharmacological Research (eIJPPR) [Impact Factor – 0.7826]

Journal Homepage: <u>www.eijppr.com</u>

Evaluation of Phytochemical Constituents and Anthelmintic Activity of Aerial Part of *Trichosanthes tricaspidata* Lour.

Brijesh Kumar Duvey

Research Article

Department of Pharmacy, Manav Bharti University, Solan (HP), India

Article info

Article History: Received 17 August 2013 Accepted 19 October 2013

Keywords: *Trichosanthes tricaspidata,* Albendazole, Anthelmintic activity, Chemical constituents.

Abstract

Recent times the trend of medicinal system of world is shifting from synthetic to herbal medicine, so we can say 'Come Back to Nature'. Medicinal plants known as millenaries and are highly respected all over the world as a rich source of therapeutic agents for the anticipation of diseases and ailments. Herbal drugs are well accepted among rural and urban community of Indian indigenous medicinal system (Ayurvedic, Sidha etc). *Trichosanthes tricaspidata* Lour. which is frequently known as Indrayan is a very little-exploited native plant of India has massive medicinal potential. The aim of this study was preliminary phytochemical investigation to find out the chemical constituents of the plant as well as evaluation of anthelmintic activity with respect to standard drug Albendazole. The result suggested that aerial part of this plant has several chemical constituents like carbohydrate, glycosides, protein, amino-acids, phenolic compounds, saponins and tannins. Ethanolic and aqueous extract of the plant also show significant value for anthelminitic activity in dose dependent manner. But the actually chemical constituents responsible for pharmacological action is yet to be investigate. So further, investigation is required for evaluation and isolation of the chemical constituent which is responsible for these pharmacological actions.

1. INTRODUCTION

India, with its great biodiversity, has a tremendous potential and advantage in the emerging field of herbal medicines. Medicinal plants as a group comprise approximately 7500 species and include representatives of about 17,000 species of higher flowering plants¹ While only three hundred species are using by 7800 medicinal drug manufacturing units in India² which consume about 2000 tons of herbs annually³. *Trichosanthes tricaspidata* Lour. (Fig.1) which is commonly known as Indrayan is a very little-exploited plant with immense medicinal potential⁴.



Fig.1: Trichosanthes tricaspidata Lour.

The vernacular names and scientific classification of *Trichosanthes tricaspidata* Lour. is represented in table-1. *T. tricuspidata* is vine which is native to East Asia. It is found at an elevation of 1200 to 2300 m. The plant is grown in forest edges and thickets in China,

*Corresponding Author Brijesh Kumar Duvey, Department of Pharmacy, Manav Bharti University, Solan (HP), India. Email: brijesh.duvey@gmail.com Japan, Mongolia, Vietnam, Malaysia and India. Indrayan is a large climber, 5-20 m long. It has strong, woody and grooved stem, with trailing branches. Tendrils are divided into 2-3 parts. Leaves are 6-12 cm across broadly heart-shaped but palmately 3-5 lobed. The leaf margin is toothed. Flowers occur either singly or in 5-10 flowered racemes, 3-4 cm long, white, with 5 wedge-shaped petals with frilly margin. Fruit is spherical initially green, 4-5 cm across, red when ripe, streaked with 10 orange streaks. Flowering May - April⁵. ⁶. The aim of this study was preliminary phytochemical investigation to find out the chemical constituents of the plant as well as evaluation of anthelmintic activity with respect to standard drug Albendazole.

2. MATERIALS AND METHODS

2.1 Collection and Identification of Plant material

Plant aerial parts were collected during the month of October from District Solan of Himachal Pradesh and was identified and authenticated as *Trichosantes palmate* Roxb. Synonyms *Trichosanthes tricuspidata* Lour., by Dr. Rajesh Raina, Sr. Scientist/ Prof. (Medicinal Plants) Department of Forest Products, Dr. Y. S. Parmar University of Horticulture and Forestry, Nouni, H.P. The herbarium sheet of the same plant sample was submitted in the Department with UHF-herbarium Field book No. 12397 dated 14/ 10/2011. Collected plant is then dried at room temperature and pulverized to optimum size.

2.2 Chemical and Reagents

Albendazole was used as standard drug for anthelmintic activity and various chemical and reagents of analytical grade used for phytochemical evaluation during this study.

3.3 Animals Used

Adult earthworms (Eisenia foetida) were used to evaluate anthelmintics activity using *in vitro* models. Earthworms were purchased from the Department of Agriculture, Solan, Himachal

VERNACULAR NAME			SCIENTIFIC CLASSIFICATION		
Hindi Marathi Tamil Malayalam Telugu Kannada Gujarati Sanskrit	: Mahakal, Indrayan : Kaundal : Ankorattai, Korattai : Kakkattonti : Avaduta : Avaguda-hannu : Ratanindrayan : Kakanasa, Shvetpushpi, Dhvamksanasa	Kingdom Family Genus Species Authority Synonyms	: Plantae : Cucurbitceae (Pumpkin family) : Trichosanthes : tricuspidata : Lour. : <i>T. palmata Roxb., T. bracteata Lamb.,</i> <i>T. pubera Blume or Modeccca bracteata</i> .		

Pradesh, vide bill No. 5511, Dated 11/04/2012. The average size of earthworm was 6 - 8 cm.

3.4 Preliminary Phytochemical Screening

The dried vein of the plant was mechanically crushed and powdered aerial part of plant (500g) was extracted by Soxhlet apparatus in ethanol and distilled water (5 liters each), at 60°C for 7 days. The extracts were filtered and concentrated to dryness using a rotary evaporator. Dry extracts were stored at 4°C till further use. The extracts obtained were subjected to qualitative chemical tests for identification of various plant constituents present in the crude drug. The extract was subjected to preliminary phytochemical investigation for detection of alkaloids, carbohydrates, glycosides, phenolic compounds, flavonoids, protein and amino acids, saponins, sterols, acidic compounds, gum, resins as per detection methods given in various books and literature⁷⁻¹³.

3.5 Anthelmintic Activities

The anthelmintic assay was carried out as per the method of Ajiyeoba¹⁴. The assay was performed in vitro using adult earthworm (*Eisenia foetida*) due to its anatomical and physiological resemblance with the intestinal roundworm parasites of human beings for preliminary evaluation of anthelmintic activity^{15, 16, 17}. Use

of (*Eisenia foetida*) species as a suitable model for screening of anthelmintic drug was advocated earlier^{18, 19}. Test samples of the extract was prepared at the concentrations 10, 20 and 40 mg/ml in distilled water and four worms of approximately equal size (same type) were placed in each nine cm Petri dish containing 40 ml of test solution of extracts. Albendazole (20 mg/ml) was used as standard and distilled water as control. All the test solution and standard drug solution were prepared freshly before starting the experiments^{19,-21}. Observations were made for the time taken for paralysis was noted when no movement of any sort could be observed except when the worms were shaken vigorously. Time for death of worms were shown in table and expressed as a mean \pm SEM of four worms in each group.

4. RESULTS AND DISCUSSION

Preliminary phytochemical screening of extract revealed the presence of glycosides, carbohydrate, saponins and tannins while the presence of phenolic compounds, mucilage and proteins was not established (Table-2).

Sr. No.	Test	Observation		Informa	
		Water extract	Ethanolic extract	Interence	
1	Test for Carbohydrate				
	Molish test	+	+		
	Fehling reagent test	+	+	Present	
	Bennedict's test	+	+		
2	Test for Glycosides				
	Keller Killani	-	-		
	Modified Borntrager test	+	+	Present	
	Borntrager test	+	+		
3	Test for Phenolic compound and Tannins				
	5% Fe Cl ₃	-	-		
	Dilute HNO3 test	+	+	Present	
	Lead Acetate test	+	+		
4	Test for Protein and Amino acid				
	Millons reagent	-	-	May be present	
	Ninihydrin test	+	+	May be present	
5	Test for Saponins				
	Foam test	+	+		
	Extract + NaHCO ₃	+	+	Present	
	Salkowski test	-	+		

Table 2: Results of the preliminary phytochemical screening

Both aqueous and ethanolic extract were evaluated for anthelmintic activity and observations made from the table is that, higher concentration of extract produced paralytic effect much earlier and the time of death was shorter for all worms. It was found that 40mg/ml ethanolic extract and aqueous extract of *Trichosantes tricaspidata Roxb*. show more significant action when compared with standard drug albendazole in concentration of 20mg/ml (Table-3). But when ethanolic extract and aqueous extract of *Trichosantes tricaspidata Roxb* were compared it was found that ethanolic extract show more potent action then aqueous extract.

Anthelmintic activity shown by both extract was dose-dependent manner.

From the above results, it is concluded that *Trichosantes palmate* (Roxb) used by tribals traditionally to treat intestinal worm infections, showed significant anthelmintic activity. The experimental evidence obtained in the laboratory model could provide a rationale for the traditional use of this plant as anthelmintic. The plant may be further explored for its phytochemical profile to recognize the active constituent accountable for anthelmintic activity.

Crowno	Concentration (mg/ml)	Eisenia foeteda					
Groups		Paralyzing time (minutes)	Death time (minutes)				
Distilled Water	-	8	8				
	10	95 ± 2.041	265 ± 8.660				
Aqueous Extract	20	28 ± 1.780	183.75 ± 9.869				
	40	9.5 ± 0.866	37.75 ± 1.652				
	10	35.50 ± 2.466	282.50 ± 9.869				
Ethanolic Extract	20	30.25 ± 1.031	193.75 ± 10.282				
	40	11 ± 1.225	51.75 ± 10.282				
Standard (Albendazole)	20	6 ± 0.408	36 ± 2.160				

Table 3: Results of Anthelmintic activity

5. CONCLUSION

Trichosanthes tricaspidata Lour. is native to India, occurring in wild area of the Sub-Himalayan region, upto the height of 1500 m. It is a vein and rich source of glycosides, tanins and flavonoids, which may be responsible for its pharmacological action. Traditionally it is used as anti-inflammatory, digestive, anthelmintic, expectorant, arthritis, inflammations etc. The present work embodies phytochemical investigation and biological evaluation on the aerial part of *Trichosanthes tricaspidata* Lour.. Phytochemical screening of ethanolic and aqueous extracts revealed the presence of carbohydrates, glycosides, phenolic compounds, flavonoids, proteins/amino acids, saponins and sterols etc. The result of *in vitro* anthelmintic study suggests that the aerial part of *Trichosanthes tricaspidata* Lour. show dose dependent anthelmintic action which is significant with respect to standard drug Albendazole and this is first ever reported.

REFERENCES

- Shiva MP. Inventory of forestry resources for sustainable management and biodiversity conservation. New Delhi: Indus Publishing Company. 1996.
- Ahmad RU, Medicinal plants used in ISM—their procurement, cultivation, regeneration and import/export aspects—a report. In: JN Govil, VK Singh and H Shamina (eds.), Medicinal Plants: New Vistas of Research (Part I). In: Glimpses in Plant Research, v. 10. New Delhi: Today and Tomorrow's Printers and Publishers. 1993, 221-258.
- Singh HP. National perspective on development of medicinal and aromatic plants. Technical report. Agri 2001.
- Dagar HS and SA Chaghtai *Trichosanthes bracteata* (Lam.) Voigt (Cucurbitaceae) - a promising ethnomedicinal taxon in Andaman and Nicobar Islands. *Indian Journal of Applied and Pure Biology*, 1989; 4(2): 131-132.
- Bhandari S., Trichosanthes Tricuspidata: A Medicinally Important Plant, Trees for Life Journal 2008. Bhandari S, Dobhal U, Sajwan M, Bisht NS. Trichosanthes tricuspidata: A medicinally important plant. Trees for Life J. 2008; 3(5): 1.
- Duvey BK, Goyel R., Parashar B. et al. "*Trichosanthes tricuspidata*: Exploration of Its Medicinal Value" Asian J. Pharm. Tech. 2012; 2(1), 26-28
- 7. Evans WC. *Pharmacognosy*, W B Saunders and company Ltd, Oval road, London, 24, 1997.p125-126.

- Gupta P, Vasudeva N, Sharma SK. Preliminary phytochemical and pharmacognostical studies of *Tagetes erecta* Linn., *Hamdard Medicus*, 52, 2009. p153-160.
- 9. Harbone JB. Phytochemical Methods; *A guide to modern techniques of plant analysis*, Chapman and Hall, London, 1988. p1-226.
- 10. Kokate CK. "*Practical Pharmacognosy*", Vallabh prakashan, New Delhi, 1991. p107-113.
- Mukurjee PK. Quality control of herbal drugs; an approach to evaluation of botanicals, 1st edition, Business Horizons, New Delhi. 2002.
- 12. Vogel AIC. A Text Book of Macro and Semi-macro Qualitative Inorganic Analysis, Longman, Green and Co. Ltd., London, 1953. p 489-563.
- Yadav N, Vasudeva N, Singh S, Sharma SK. Pharmacognostical and phytochemical study on *Chinopodium* album root, *Hamdard Medicus* 2007; 50(1), 95-102.
- Ajaiyeoba EO, Onocha PA, Olarenwaju OT. In vitro anthelmintic properties of *Buchholzia coriaceae* and *Gynandropsis gynandra* extract. Pharm Biol. 2001; 39: 217-20.
- 15. Vigar Z. Atlas of Medical Parasitology. 2 nd ed. P. G. Publishing House Singapore 1984. p. 242.
- Dash GK, Suresh P, Kar DM, Ganpaty S, Panda SB. Evaluation of *Evolvulus alsinoides* Linn. For anthelmintic and antimicrobial activities, *J Nat Rem.* 2002; 2: 182-185.
- 17. Shivkumar YM, Kumar VL. Anthelmintic activity of latex of *Calotropis procera. Pharma Biol.* 2003; 41: 263-265.
- Kaushik RK, Katiyar JC, Sen AB. Studies on the mode of action of anthelmintics with *Ascardia galli* as a test parasite. *Indian J Med Res.* 1974; 64: 1367-75.
- Yadav AK, Temjenmongla. Anthelmintic activity of *Gynura* angulosa against *Trichinella spiralis* infections in mice. *Pharmacologyonline* 2006; 2: 299-306
- Mali RG, Shailaja Mahajan, Patil KS. Anthelmintic activity of root bark of *Capparis spinosa*. Indian J Nat Prod. 2005; 21: 50-51.
- 21. Gbolade AA, Adeyemi AA. Investigation of in vitro anthelmintic activities of *Pycnanthus ngeolensis* and *Sphenocentrum jollyanum. Fitoterapia* 2008; 79: 200-222.
- Lal J, Chandra S, Raviprakash V, Sabir M. In vitro anthelmintic action of some indigenous medicinal plants on Ascardia galli worms. Indian J Physiol Pharmacology 1976; 20: 64-68.