

Prevalence of Obesity in Primary School Students and its Relation to Watching TV in Arar City

Afaf Amer Flag Alenazi^{1*}, Sabry Mohamed Hammad², Amel Elwan³

¹ Family Medicine Resident, Arar Primary Health Care Centers, KSA.

² Professor of Public Health & Community Medicine, Faculty of Medicine, Mansoura University, Egypt & Consultant Public Health & Community Medicine, Northern Borders General Health Affairs, KSA.

³ Lecturer of Public Health & Community Medicine, Consultant Public Health & Community Medicine, Northern

Borders General Health Affairs, KSA.

ABSTRACT

Background: The prevalence of obesity and overweight in children has risen in both males and females. **Objectives**: To investigate the prevalence of overweight and obesity among female primary school students (7-12years old) in Arar city, 2018, and explore its association with watching television. **Methods**: A cross-sectional study was conducted, including 330 girls, using the multistage cluster sampling technique. Parents completed a structured questionnaire about demographic characteristics and lifestyle factors (diet and physical activity). Every child's body weight and height were measured and plotted on Growth charts for female Saudi children. Data were collected during the first semester of 2018 **Results**: The prevalence of overweight and obesity was 15.9%, 9.6% had a family history of obesity, about 91.7% of the students consume \leq 3 meals but 8.3% consume >3 meals daily. 15% of them eat fast food in two and more of their daily meals. The daily time of video games was >2 hours in 37.6%. 32.8% were watching TV alone away from parents. Only 44.6% of girls practice physical exercise. There was significant relation (P<0.05) between obesity, fast food consumption, frequency of daily meals, watching TV alone away from parents, physical inactivity, daily time of video games, and family history of obesity. **Conclusion**: The prevalence of overweight and obesity among primary school girls was considerable. They have bad dietary habits and lack physical exercise. Family physicians counseling of parent on national education is recommended to educate girls on healthy lifestyle.

Key Words: obesity, prevalence, children, primary school, KSA.

eIJPPR 2021; 11(1):159-166

HOW TO CITE THIS ARTICLE: Afaf Amer Flag Alenazi, Sabry Mohamed Hammad, Amel Elwan (2021). "Prevalence of Obesity in Primary School Students and its Relation to Watching TV in Arar City", International Journal of Pharmaceutical and Phytopharmacological Research, 11(1), pp.159-166.

INTRODUCTION

Childhood obesity is a serious health problem internationally. World Health Organization (WHO) defines obesity as a condition of excessive body fat that exceeds healthy limits that risk health [1]. Overweight and obese children are expected to continue obesity into maturity and more probable to get non-communicable diseases like diabetes and cardiovascular problems at an earlier age [2, 3].

Identifying ecological effects that influence body weight in children is important for developing suitable protective strategies. One possible ecological issue is the extent of TV watched [4]. Childhood obesity is rising in Saudi Arabia with a distressing rate. In the prevalence of overweight and obesity, risk factors included higher parental education, and lifestyle factors. lifestyle factors included increased consumption of calorie-dense food and a sedentary lifestyle [5].

The prevalence rate of childhood overweight and obesity continued to increase reaching 48.3% in males in 2018 and to 27.6% in females in 2017. Management of adolescents with obesity in Saudi Arabia is considered a public health priority [6].

In a study in Alhsa, it was reported that the general prevalence of overweight and obesity among school children was 29.6% (10.8% overweight, 3.8% obese, and 15% extremely obese). The occurrence of overweight and

Corresponding author: Afaf Amer Flag Alenazi

E-mail: 🖂 afooof-2011 @ hotmail.com

Address: Family Medicine Resident, Arar Primary Health Care Centers, KSA.

Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 10 November 2020; Revised: 09 February2021; Accepted: 11 February 2021

obesity was considerably affected by maternal obesity, employment of the mother, income of the family, eating snacks and takeaway foods, bodily laziness, and period of time spent watching television [7].

In Taif city, Saudi Arabia, childhood overweight and obesity is a public health problem affecting more than onethird of girls and almost one-fifth of boys. It confirms the significant association between some unhealthy dietary habits and increases the prevalence of childhood obesity [8].

Several studies reported that higher TV numbers of watching hours are related to increasing body mass index (BMI), decreased body fitness, and increased blood cholesterol levels [9].

Three possible mechanisms have been projected to clarify the connection between TV watching and obesity; First, watching TV may lead to reduced physical activity, decreased energy expenditure comparative to energy consumption, and leads to obesity. Second, the mental immaturity of children rises their vulnerability to influential intent of advertisements for foods of deprived nutritious value. Food advertisements are broadcast frequently during children's television programs and prime to corrupt food favorites, and dietary inequities related to obesity. Third, action shift joint with vulnerability to convincing intent of televised food advertisements influences young children to early obesity [10].

There is no information on the prevalence of obesity among female primary school children in the Northern Borders region, Saudi Arabia. This study aimed to determine the prevalence of overweight and obesity among female primary school students (7-12 years old)

SUBJECTS AND METHODS

A cross-sectional study was conducted on Saudi female Primary schools in Arar city

Sample size was calculated using StatCalc of Epi Info software version 7.222 using the following criteria; population size 3700 students, female expected frequency (prevalence of overweight and obesity was 12.4% So, the final sample size was 330 students.

Total number of primary schools in Arar city is 25 schools. Sampling was selected using the multi-stage cluster sample technique. In the first stage, 2 schools were selected randomly. In the second stage, one class from each grade was randomly selected. All children of each chosen class were included in the study.

Official permission was obtained from the educational authorities in Arar. A structured questionnaire was used to collect demographic characteristics and lifestyle factors (diet and physical activity) from parents. Pilot study was carried out on 30 students to test the questioner. No modification was required.. The Questionnaire was given to the parents of the children to complete the data and to sign the consent. The questioner assessed the following variables; age, gender, grade, number of siblings, height, weight, any chronic disease, physical activity, number of TVs in the home, number of hours watching TV per day, eating junk foods, eating breakfast before going to school, eating vegetables, and usually drink fresh juice.

The researcher visited the school 3 times per week during the period from 1st January to 31st May 2019. Anthropometric assessment was done for the children. The children were asked to stand on weigh scale with minimal clothing and without shoes. The weight was measured in kg to the nearest decimal fraction. The height was measured in meter using a suitable tap meter scale fixed on the wall, height of the subjects was measured without shoes taking into that heels, buttocks, shoulders, and head were closed to the vertical wall surface. Diagnosis of overweight was done BMI-for-age percentile >85th≥95th. Obesity was diagnosed by BMI-for-age percentile >95th centile after plotting on age-specific growth charts for Saudi females [11]. For assessing the physical activity of children and adolescents, a physically active child should do a total of at least 60 min/day moderate-to-vigorous physical activity (MVPA) or vigorous physical activity in bouts of at least 10 min [12].

Statistical analysis was performed using the statistical package for social sciences (SPSS version 20.0). Quantitative data were expressed as mean \pm SD & (range), and qualitative data were expressed as absolute frequencies (number) & relative frequencies (percentage). Percentage of categorical variables were compared using the Chi-square test or Fisher exact test when appropriate. P-value <0.05 was considered statistically significant (S), and p-value \geq 0.05 was considered statistically insignificant (NS).

RESULTS:

According to (able 1, BMI distribution, 84.1% had normal BMI and 15.9% were overweight and/or obese (**Figure** 1). Family history of obesity was positive in 9.6% of them, 91.7% consumed \leq 3 meals but 8.3% consumed >3 meals daily and 15.3% ate fast food in 2 and more of their daily meals. A daily time of video games was >2 hours in 37.6% (**Figure** 2), 32.8% were watching TV alone away from parents (as shown in **Figure** 3) and physical exercise was performed by 44.6% of participants. There was a significant relationship (P<0.05) between the BMI of study children, family history of obesity, fast food consumption. However, no significant relationship was observed between the BMI of study children and their daily meals, watching TV alone away from parents, physical exercise, and daily time of video games.

Table (2) shows the relationship between BMI and the study group's socio-demographic characteristics. There

was a significant relationship between the BMI and fathers' education (P<0.05), while there was no significant relationship between the BMI and other socio-demographic characteristics.

Table (3) shows the relationship between the BMI of the study group and their dietary history and family history of obesity. There was a significant relationship between the

BMI and family history of obesity, fast food consumption, and the number of daily meals (P<0.05).

Table (4) shows the relationship between BMI of the study group and watching TV and physical exercise. Our study found that there was a significant relationship between the BMI and watching TV alone away from parents, physical exercise, and daily time of video games (P<0.05).

Table 1: Socio-demographic characteristics, BMI, dilatory history, family history of obesity, watching TV, a	nd
physical exercise in the study group	

Variables	Responses	No. (%)	
Ago	Mean(±SD)	10.2(±1.36)	
Age	Range	7-13	
Nationality	Saudi	241(76.0)	
Trationanty	Non-Saudi	73(23.2)	
Fathers' education	Less than secondary school education	27(8.6)	
Famers education	Secondary school education and above	287(91.4)	
Mathema advection	Less than secondary school education	39(12.4)	
Womers education	Secondary school education and above	275(87.6)	
	Unemployed	25(8.5)	
Fathers work	Government's employee	196(66.4)	
	Private sector employee	74(25.1)	
Working mother	Housewife	115(37.0)	
	Employed	196(63.0)	
	1-2	168(53.5)	
Child order	3-4	94(29.9)	
	≥5	52(16.6)	
	1 st	30(9.01)	
	2 nd	36(10.9)	
Child grade in the school	3 rd	80(24.2)	
	4 th	107(32.4)	
	5 th	77(23.3)	
	Insufficient	30(9.6)	
Family income	Satisfactory	138(43.9)	
	More than satisfactory	146(46.5)	
Family history	Yes	29(9.2)	
Failing instory	No	285(90.8)	
BMI distribution	Normal	264(84.1)	
Divit distribution	Overweight and obese	50(15.9)	
Fast food	Less than 2 meals	26684.7)	
Fast food	2 and more	48(15.3)	
Daily meals	≤3 meals	288(91.7)	
Daily incais	> 3 meals	26(8.3)	
TV number in the home	≤1	85(27.1)	
	>1	229(72.9)	
Daily time of video games	\leq 2 hours	196 (62.4)	
Dury time of video games	> 2 hours	118(37.6)	
Physical exercise	Yes	140(44.6)	
	No	174(55.4)	
Watching TV alone away	Yes	103(32.8)	
from parents	No	211(67.2)	

International Journal of Pharmaceutical and Phytopharmacological Research(eIJPPR) | February 2021 | Volume 11 | Issue 1 | Page 159-166 Afaf Amer Flag Alenazi, Prevalence of Obesity in Primary School Students and its Relation to Watching TV in Arar City



Figure 1: Percentage distribution of the study Children by BMI in Arar, 2019



Figure 2: Daily time of video games spent by children, Arar, 2019



Figure 3: Watching T.V alone away from parents, Arar, 2019

Table 2: Relationship between BMI and socio-demographic characteristics of the study group

		BM	Total			
Variables	Responses	No overweight and	Overweight and	(N=214)	P value	
		obesity (n=264)	obesity (n=50)	(N=314)		
Nationality	Saudi	206(78.0)	35(70.0)	241(76.0)	0.2	
	Non-Saudi	58(22.0)	15(30.0)	73(23.2)		
Fathers'	Less than secondary school education	27(10.2)	0(0.0)	27(8.6)	0.012	
education	secondary school education and above	237(89.8)	50(100)	287(91.4)	0.012	

International Journal of Pharmaceutical and Phytopharmacological Research(eIJPPR)| February 2021 | Volume 11 | Issue 1 | Page 159-166 Afaf Amer Flag Alenazi, Prevalence of Obesity in Primary School Students and its Relation to Watching TV in Arar City

Mothers	Less than secondary school education	33(12.5)	6(12.0)	39(12.4)	0.0
education	secondary school education and above	231(87.5)	44(88.0)	275(87.6)	0.9
Fathers work	Unemployed	24(9.6)	1(2.2)	25(8.5)	
	Government's employee	160(64.3)	36(78.3)	196(66.4)	0.1
	Private sector employee	65(26.1)	9(19.6)	74(25.1)	
Working	Housewife	93(35.6)	22(44.0)	115(37.0)	0.26
mother	Employed	168(64.4)	28(56.0)	196(63.0)	0.20
Child order	1-2	136(51.5	32(64.0)	168(53.5)	
	3-4	80(30.3)	14(28.0)	94(29.9)	0.14
	≥5	48(18.2)	4(8.0)	52(16.6)	
Family income	Not sufficient	25(9.5)	5(10.0)	30(9.6)	
	Satisfactory	117(44.3)	21(42.0)	138(43.9)	0.9
	More than satisfactory	122(46.2)	24(48.0)	146(46.5)	

Table 3: Relationship between BMI of the study group and their dietary history and family history of obesity

		BM		D	
Variables	Responses	No overweight and	Overweight and	Total (N=314)	r -
		obesity (n=264)	obesity (n=50)		value
Fast food	Less than 2 meals	229(86.7)	37(74.0)	26684.7)	0.02
	2 and more	35(13.3)	13(26.0)	48(15.3)	0.02
Daily meals	\leq 3 meals	246(93.2)	42(84.0)	288(91.7)	0.046
	> 3 meals	18(6.8)	8(16.0)	26(8.3)	0.040
Family history of	Yes	17(6.4)	12(24.0)	29(9.2)	0.000
obesity	No	247(93.6)	38(76.0)	285(90.8)	0.000

Table 4: Relationship between BMI of the study group and watching TV and physical exercise

		BMI			р
Variables	Responses	No overweight and obesity (n=264)	Overweight and obesity (n=50)	Total (N=314)	P- value
TV number in the home	≤ 1	76(28.8)	9(18.0)	85(27.1)	0.1
	> 1	188(71.2)	41(82.0)	229(72.9)	0.1
Daily time of video games	\leq 2 hours	171(64.8)	25(50.0)	196 (62.4)	0.048
	> 2 hours	93 (35.2)	25(50.0)	118(37.6)	0.048
Watching TV alone away	Yes	79(29.9)	24(48.0)	103(32.8)	0.01
from parents	No	185(70.1)	26(52.0)	211(67.2)	0.01
Physical exercise	Yes	125(47.3)	15(30.0)	140(44.6)	0.02
	No	139(52.7)	3570.0)	174(55.4)	0.02

DISCUSSION

The present cross-sectional study was conducted among 330 female students in the primary school in Arar, KSA. The study aimed to determine the prevalence of obesity among female primary school students (6-12 years old) in Arar city, Northern Saudi Arabia, during the academic year of 2018-2019 and explore its association with watching television.

As regards the percentage distribution of the study children by BMI, the present study found that 15.9% of female students were overweight and obese. In Taif, KSA, in schoolboys, the prevalence of overweight and obesity were 10.7% and 7.6%, respectively; while in girls, they were 16.8% and 18.2%, respectively [8]. In Jazan, overweight and obesity were 10.1% and 12.4%, respectively, and both obesity and overweight were more frequent in children consuming more fast food [13].

Aljaadi and Alharbi stated that the frequency of obesity in Saudi children is increasing and is associated with psychosomatic and societal consequences. It differs according to gender, area of residence, and description of cases [14].

As regards the association of the BMI of the study group with watching TV and physical exercise, our study reported a significant association between the BMI and watching TV alone away from parents, physical exercise, and daily time of video games (P<0.05).

Similar to our results another study reported that there were significant correlations between the prevalence of obesity and physical exercise and watching TV (P<0.050) and the findings showed that performing any type of sport as

biking, swimming, or football frequently is considerably associated with less prevalence of overweight and obesity (P<0.001) [15]. Also, another study found that there was a significant relationship between BMI, and physical exercise and computer games but no significant relationship was observed with watching TV [16]. In contrast to our results, another study found that there was no significant association between obesity and time spent watching TV and using the computer or playing video games [17].

Also, a systematic review that included a total of 66 analyses of data on the relationship between TV watching and obesity found that 47 studies stated a positive association, while 19 studies stated no relationship, and no studies reported an inverse association between watching TV and obesity. It was stated that watching TV reduced resting metabolic rate while it was not supported by other studies. The results of the studies that observed the link between TV watching and physical activity were unreliable. Of the 34 studies recognized, two stated a confident association, 16 found no association, 15 concluded there was an inverse association, and one had mixed results. The national Children's Nutrition Survey is among the studies that reported an association between watching TV and a less nutritious dietary intake [18].

On the other hand, Katzmarzyk et al. (1998) reported a positive association (P< 0.05) between the time of watching TV and obesity in boys aged 9-12 years, but not in any other gender-age subgroups [19]. Also, TV viewing was positively associated with BMI and skinfold thickness in both genders, although it was significant only for BMI in boys (P <0.05) as reported by Guillaume et al. (1997) [20]. This is in line with Ozdirenc et al. (2005) who reported a significant positive correlation between BMI and the amount of watching TV (P<0.05) [21]. Again, time per day spent watching TV was positively associated with having a BMI above the 50th percentile (P= 0.001) as reported in a cohort by Maffeis et al. (1998) [22]. In addition, Grund et al. (2001) reported that children who watched TV >1 hr/day had a higher BMI (22.0 vs 19.8) and fat mass (30.7% vs 24.5%) than those who watched less while children watching TV >3 hours/day had about 40% increased prevalence of obesity compared with children watching <1 hour/day as reported by Guan-Sheng et al. (2002) [23, 24]. Findings are consistent with Scragg et al. (2004) who reported children watching TV >2 hours per day had higher BMI than those watching <1 hour per day $(19.3 \text{ vs } 18.7 \text{ kg/m}^2; P= 0.005)$ [25].

Regarding overweight and obesity risk factors, the current study concluded that an unhealthy diet and sedentary life were significantly associated with increased BMI. These findings were similar to previous studies. A study in Riyadh on expatriate girls found that 34.2% were insufficiently active [26]. Burke et al. (2005) in a cohort study reported that the mean BMI increased by 0.185 units for each extra hour/day of watching TV [27].

As regards the relationship between BMI of the study group with their dietary history and family history of obesity, we found that there was a significant relationship between the BMI and family history of obesity, fast food consumption, and number of daily meals (P<0.05).

In a study on dietary habits of Saudi children, one-quarter of children recorded eating sweets daily, and >50% recorded eating sweets irregularly, 56.3% of children reported drinking carbonated beverages occasionally and 17.1% drank it daily. Energy drinks weekly were found in 21.9% and daily drinking was found in 4.3% [28]. Another study stated a strongly positive correlation between child BMI and parents' BMI (P<0.001). It is possibly caused by hereditary factors or lifestyle issues [15].

Numerous studies agreed with these findings. Mo-suwan et al. (2000) reported that the hazard of child obesity was augmented 2-3 times by a positive family history of obesity. However, Al-Enazy et al. (2014) found no substantial association between the occurrence of obesity and the history of paternal obesity.

CONCLUSION AND RECOMMENDATIONS:

The prevalence of overweight and obesity among primary school girls was considerable (15.9%). Family history, fast food consumption, watching TV alone away from parents, physical exercise, and daily time of video games were significant risk factors. So we recommend nutrition education for school girls and their parents.

Study limitations:

This was a cross-sectional study, which reflects the current situation only but cannot allow studying the causal relationship, which needs a follow-up study. Also, there were no detailed investigations of other causes as the genetic, dietary pattern from birth to now, and history of chronic diseases causing weight gain.

ACKNOWLEDGMENT:

We would like to thank our participants who were all dedicated to the study. We would also like to thank our professors who spared no effort in mentoring us and providing us with proper feedback.

Authors Contributions:

Afaf Amer Flag Alenazi shared in setting the study design, research objectives, preparing study instrument, pilot administration, data collection, and reviewing the results. Prof. Sabry Mohamed Hammad held study approvals and supervised study phases. Dr. Amel Elwan shared in International Journal of Pharmaceutical and Phytopharmacological Research(eIJPPR) | February 2021 | Volume 11 | Issue 1 | Page 159-166 Afaf Amer Flag Alenazi, Prevalence of Obesity in Primary School Students and its Relation to Watching TV in Arar City

preparing the study proposal, a logistics plan, data collection plan, supervised data entry, and conducted the statistical analysis, data display, discussion guidelines, and final write-up. All authors read and approved the final manuscript.

Funding

This study has received no external funding.

Conflict of interest

The authors would like to declare that there is no conflict of interest.

Informed consent

Written & Oral informed consent was obtained from all individual participants included in the study. Additional informed consent was obtained from all individual participants for whom identifying information is included in this manuscript.

Ethical approval for study protocol /study design /Methodology:

The study was approved by research ethical local committees of General Health Affairs of Northern Border region with researcher No. (0559748194), committee No. (H-09-A-51), and approval No. (1440/7).

REFERENCES

- [1] Ryabaya IN. Clinical laboratory and instrumental peculiarities of atrial fibrillation in obese patients. Arch. Pharma. Pract. 2020;11(2):1-6.
- [2] Bogoriani NW, Suaniti NM, Putra AA, Pradnya KD. The Activity of Cordyline Terminalis's Leaf Extract as Antidiabetic in Obese Wistar Rats. Int. j. pharm. res. Allied sci. 2019;8(2):206-13.
- [3] Nittari G, Scuri S, Petrelli F, Pirillo I, Di Luca N, Grappasonni I. Fighting obesity in children from European World Health Organization member states. Epidemiological data, medical-social aspects, and prevention programs. Clin Ter. 2019;170(3):e223e30.
- [4] Dennison BA, Erb TA, Jenkins PL. TV viewing and TV in bedrooms associated with overweight among low-income preschool children. Pediatrics 2002;109(6):1028-35.
- [5] Hammad SS, Berry DC. The child obesity epidemic in Saudi Arabia: A review of the literature. J Transcult Nurs. 2017;28(5):505-15.
- [6] Habbab RM, Bhutta ZA. Prevalence and social determinants of overweight and obesity in adolescents in Saudi Arabia: A systematic review. Clin Obes. 2020;10(6):e12400.

- [7] Saleh AAA, Alhaiz AS, Khan AR, Al-Quwaidhi AJ, Aljasim M, Almubarak A, *et al.* Prevalence of obesity in school children and its relation to lifestyle behaviors in Al-Ahsa district of Saudi Arabia. Glob J Health Sci. 2017;9(12):1-80.
- [8] AlGhamdi RA. Prevalence of Childhood Obesity among Primary School Students at Taif Governorate, Saudi Arabia. J Adv Med Med Res. 2019;31(5):1-21.
- [9] Park JH, Moon JH, Kim HJ, Kong MH, Oh YH. Sedentary Lifestyle: Overview of Updated Evidence of Potential Health Risks. Korean J Fam Med. 2020;41(6):365.
- [10] Garcia JD, Medina IG. Children, media and food. Food advertising aimed at children audience in Spanish TV. 2020.
- [11] Aljaaly EA. A Screening Program for Overweight Status among Schoolgirls in Saudi Arabia: A Proposed Co-design Project to Tackle the Problem. Journalism. 2017;7(5):225-43.
- [12] WHO. Global recommendations on physical activity for health. Geneva; 2010. ISBN: 9789241599979 http://www who int/dietphysicalactivity/publications/9789241599979/ en/ Accessed. 2017;1:60.
- [13] Fakeeh MI, Shanawaz M, Azeez FK, Arar IA. Overweight and obesity among the boys of primary public schools of Baish City in Jazan Province, Saudi Arabia: A cross-sectional study. Indian J Public Health. 2019;63(4):330.
- [14] Aljaadi AM, Alharbi M. Overweight and Obesity Among Saudi Children: Prevalence, Lifestyle Factors, and Health Impacts. Handbook of Healthcare in the Arab World Springer, Cham https://doi org/101007/978-3-319-74365-3_187-1. 2020.
- [15] Badawi NE, Barakat AA, El Sherbini SA, Fawzy HM. Prevalence of overweight and obesity in primary school children in Port Said city. Egypt Paediatr Assoc Gaz. 2013 Jan 1;61(1):31-6.
- [16] Ajayi EO, Elechi HA, Alhaji MA. Prevalence of overweight/obesity among primary school pupils in Urban Centre, Nigeria. Saudi J Obesity. 2015;3(2):59-65.
- [17] Al-Enazy WH, Al Dahi SK, Al Hariri IM. Prevalence of overweight and obesity among Saudi primary school students in Tabuk, Saudi Arabia. Saudi J Obesity 2014;2(1):13-8.
- [18] Scragg R, Quigley R, Taylor R. Does watching TV contribute to increased body weight and obesity in children. Scientific Committee of the Agencies for Nutrition Action: Wellington, New Zealand. 2006.
- [19] Katzmarzyk PT, Malina RM, Song TM, Bouchard C. TV viewing, physical activity and health-related

fitness of youth in the Quebec Family Study. J Adolesc Health 1998;23(5):318-25.

- [20] Guillaume M, Lapidus L, Björntorp P, Lambert A. Physical activity, obesity, and cardiovascular risk factors in children. The Belgian Luxemborg Child Study II. Obes Res 1997;5(6):549-56.
- [21] Özdirenç M, Özcan A, Akin F, Gelecek N. Physical fitness in rural children compared with urban children in Turkey. Pediatr Int 2005;47(1):26-31.
- [22] Maffeis C, Talamini G, Tato L, Influence of diet, physical activity and parent's obesity on children's adiposity: a four-year longitudinal study. Int J Obes 1998;22(8):758-64.
- [23] Grund A, Krause H, Siewers M, Rieckert H, Müller MJ. Is TV viewing and index of physical activity and fitness in overweight children and normal weight children? Publ Health Nutr 2001;4(6):1245-51.
- [24] Guan-Sheng M, Yan-Ping L, Xiao-Qi H, Wen-Jun M, Jin W, Effect of TV viewing on pediatric obesity. Biomed Environ Sci 2002;15(4):291-7.

- [25] Scragg R, Wilson N, Schaaf D, Fitzgerald E, Utter J. Risk factors for obesity in New Zealand children aged 5-14 years: results from the 2002 National Children's Nutrition Survey. Aust Epidemiol 2004;11:23-4.
- [26] Alhusaini AA, Melam GR, Buragadda S. Cross-Cultural Variation in BMI, Sedentary Behavior, and Physical Activity in International School Girls Residing in Saudi Arabia. Int J Environ Res Public Health. 2020;17(6):2057.
- [27] Burke V, Beilin LJ, Simmer K, Oddy WH, Blake KV, Doherty D, *et al.* Predictors of body mass index and associations with cardiovascular risk factors in Australian children: a prospective cohort study. Int J Obes 2005;29(1):15-23.
- [28] Alsubaie ASR. Consumption and correlates of sweet foods, carbonated beverages, and energy drinks among primary school children in Saudi Arabia. Saudi Med J. 2017;38(10):1045.