



Antimicrobial Effects of Methanolic Extract of Salvia Officinalis L, Including MRSA and Multidrug Resistant Acinetobacter Baumannii

Mohammad Akram Randhawa^{1*}, Abdulhakim Bawadekji², Mouhanad Al Ali³,
Mohamed Habib Oueslati⁴, Jamith Basha W⁵

¹Department of Pharmacology, College of Medicine, Northern Border University, P.O. Box 1321, 91431 Arar, Saudi Arabia.

²Deanship of Scientific Research, Northern Border University, P.O. Box 1321, 91431 Arar, Saudi Arabia,

³University of Angers, Superior Institute of Health and Bioproducts of Angers, 16 Boulevard Daviers, 49045 Angers Cedex, France.

⁴Department of Chemistry, College of Science, Northern Border University, P.O. Box 1321, 91431 Arar, Saudi Arabia,

⁵Department of Microbiology, College of Applied Medicine, Northern Border University, P.O.Box 1321, 91431 Arar, Saudi Arabia.

ABSTRACT

Microbial infections due to opportunistic bacteria and fungi in hospitals are becoming serious health problems, particularly in immune-compromised individuals. Moreover, microbes are becoming resistant to presently available drugs. Therefore, there is a need to develop new drugs to treat microbial infections. Salvia officinalis (S. officinalis) has been widely used in Middle East countries and Europe as a food preservative and for treatment of infections. Present study was aimed to investigate activity of S. officinalis against resistant and opportunistic microorganisms. ATCC strains of some bacteria (Staphylococcus aureus, Enterococcus faecalis, Escherichia coli, Klebsiella pneumoniae, Pseudomonas aeruginosa) were purchased from Micromaster, Saudi Arabia. Methicillin Resistant Staphylococcus aureus (MRSA), Multi-drug resistant Acinetobacter baumannii and some fungi (Candida albicans and Aspergillus flavus), commonly causing serious infections in hospitals and opportunistic infections, were obtained from the Microbiology Department, Prince Abdulaziz Bin Musaad Hospital, Arar, Saudi Arabia. Agar dilution method was used to determine minimum inhibitory concentrations of methanolic extract of S. officinalis against test organisms as recommended by Clinical and Laboratory Standards Institute. The extract showed great activity against Staphylococcus aureus, MRSA and the multidrug resistant Acinetobacter baumannii (MICs 5-10 mg/ml), moderate activity against Candida albicans (MIC 10-20 mg/ml) and weak activity against Gram negative bacilli and Aspergillus flavus (MICs 20-40 mg/ml). To conclude, methanolic extract of S. officinalis possesses good activity against resistant and opportunistic infections: Staphylococcus aureus, MRSA, multidrug resistant Acinetobacter baumannii and Candida albicans. Further investigations are needed to isolate active components from S. officinalis and develop new drugs.

Key Words: Salvia Officinalis, Methanolic Extract, Opportunistic Infections, MRSA, Multi Drug Resistant Acinetobacter Baumannii

eIJPPR 2018; 8(4):1-5

HOW TO CITE THIS ARTICLE: Mohammad Akram Randhawa, Abdulhakim Bawadekji, Mouhanad Al Ali, Mohamed Habib Oueslati, Jamith Basha W. (2018). "Antimicrobial Effects of Methanolic Extract of Salvia Officinalis L, Including MRSA and Multidrug Resistant Acinetobacter Baumannii", International Journal of Pharmaceutical and Phytopharmacological Research, 8(4), pp.1-5.

INTRODUCTION

Microbial infections due to opportunistic bacteria and fungi in hospitals, particularly in immune-compromised

individuals, are becoming a serious health problem. Microbes are also becoming resistant to available antimicrobial drugs, thus requiring alternative medications [1]. Moreover, antimicrobial drugs killing the bacteria and

Corresponding author: Mohammad Akram Randhawa

Address: Department of Pharmacology, College of Medicine, Northern Border University, P.O.Box 1321, 91431 Arar, Saudi Arabia.

E-mail: ✉ mrakramsa@yahoo.co.uk

Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 06 May 2018; **Revised:** 12 July 2018; **Accepted:** 28 August 2018



fungi are likely to damage the human cells. Antimicrobial resistance, toxicity and poor immune-system of the host are the major problems encountered in the management of infectious diseases, and require the development of new, more effective and less toxic drugs.

Salvia officinalis L, (*S. officinalis*) commonly named as 'Sage' in English, and 'Maramia' in Arabic, has been used since ancient times as herbal tea, flavoring agent in food, a food preservative, and for the treatment of skin and respiratory infections, as well as for the elevation of mood, feeling of wellbeing and improvement of memory in Saudi Arabia, other Middle East countries and Europe [1, 2].

Recently, extracts of *S. officinalis* and some other species of *Salvia* were shown to possess antibacterial, antifungal and insecticidal activities, against gram-positive bacteria (such as *Staphylococcus aureus*, *Micrococcus luteus*, *Bacillus cereus* and *Bacillus subtilis*), gram negative bacteria (such as *Escherichia coli*, *Salmonella enteritidis* and *Agrobacterium tumefaciens*), food-borne and opportunistic fungi (*Aspergillus* and *Fusarium* species) as well as insecticidal effects against *Spodoptera littoralis* larvae and *Tribolium castaneum* adults [3].

The present study was aimed to further explore the antimicrobial activities of the methanolic extract of *S. officinalis*. Besides the determination of its activity on representative Gram positive and gram negative bacteria (*Staphylococcus aureus*, *Enterococcus faecalis*, *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*), which commonly cause opportunistic and serious infections in hospitals, its activity was also investigated against Methicillin Resistant *Staphylococcus aureus* (MRSA), Multi-drug resistant *Acinetobacter baumannii* and some opportunistic fungi (*Candida albicans* and *Aspergillus flavus*).

METHODS

Plant

S. officinalis L plant was purchased from the local market, as it is commonly available in Saudi Arabia. Its identification was certified from a botanist at the College of Science, Northern Border University, Arar, Saudi Arabia. Aerial parts of these plants were washed with distilled water, and freeze-dried.

Preparation of Methanolic Extract

100 grams of dried and powdered plant was macerated in 1000 ml of methanol (100 g/ 1000 ml) for one week at room temperature (about 25°C). The extract was then filtered using a filter funnel and the excess methanol evaporated under reduced pressure in a rotary evaporator. Once evaporated, the extract remained as a dry residue, which was stored in the refrigerator at 4°C. Immediately

before use for the antimicrobial assay, the extract was suspended in DMSO to a final concentration of 400mg/ml. Fresh suspension of the extract in DMSO was used for the estimation of its activity against each microorganism.

Microorganisms

The ATCC strains (Numbers given in Table-1) of representative gram positive and gram negative bacteria (*Staphylococcus aureus*, *Enterococcus faecalis*, *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*) were purchased from Micromaster Saudi Arabia. MRSA, multi-drug resistant *Acinetobacter baumannii* and opportunistic fungi (*Candida albicans* and *Aspergillus flavus*) were obtained from the Microbiology Department of Prince Abdulaziz Bin Musaad Hospital, Arar, Saudi Arabia.

Table 1. Antibacterial activity of Methanolic extract of *S. officinalis* (Maramia)

S.No	Name of Bacterial ATCC Strains	Methanol Extract of <i>S. officinalis</i> (mg/ml)				
		40	20	10	5	2.5
1	<i>Staphylococcus aureus</i> ATCC 254996	S	S	S	S	R
2	<i>Staphylococcus aureus</i> (MRSA)	S	S	S	S	R
3	<i>Acinetobacter baumannii</i> (MDRO)	S	S	S	S	I
5	<i>Klebsiella pneumoniae</i> ATCC 700603	S	S	R	R	R
6	<i>Pseudomonas aeruginosa</i> ATCC 254992	S	S	R	R	R
7	<i>Escherichia coli</i> ATCC 25922	S	R	R	R	R

Growth media

Muller Hinton Agar (MHA) was used for bacteria and Sabouraud Dextrose agar (SDA) for fungi, and purchased from Micromaster, Saudi Arabia.

Minimum inhibitory concentration (MIC) value determination assay

The agar dilution method was used for the determination of the minimum inhibitory concentration (MIC) of the methanolic extract of *S. officinalis* against the standard (ATCC) strains of *Staphylococcus aureus*, *Enterococcus faecalis*, *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, as well as MRSA, Multi-drug resistant *Acinetobacter baumannii*, *Candida albicans*, and

Aspergillus flavus) as recommended by the Clinical and Laboratory Standards Institute [4].

2 ml of stock solution of the extract (800 mg/2ml DMSO) was added to 18 ml of MHA (For bacteria) and 18 ml of SDA (For fungi), then serially diluted to obtain final concentrations of 40, 20, 10, 5, and 2.5 mg/ml of extract in sterile petri plates. 2ml of DMSO was added to 18 ml of MHA (For bacteria) and 18 ml of SDA (For fungi), without *S. officinalis* extract, and then serially diluted to obtain concentrations of 10, 5, 2, and 1.25% DMSO in sterile petri plates as positive controls. The standard inoculum of 1µl for bacteria and of 0.5 MacFarland (10^5 CFU) for fungi, were spot inoculated in the test and control petri plates. All the plates were incubated aerobically for 12-24 hours at 37°C for bacteria, and at 25°C for fungi. The antimicrobial activity of the methanolic extract of *S. officinalis* was determined as Minimal Inhibitory Concentration (MIC) value, and was defined as the lowest concentration of the extract completely inhibiting the growth of the microorganism.

RESULTS AND DISCUSSION

Methanolic extract of *S. officinalis* had great activity against some bacteria, like *Staphylococcus aureus*, MRSA, and multi drug resistant *Acinetobacter baumannii*, (MIC of 5-10mg/ml of the extract); and there was a significant growth of these bacteria in corresponding DMSO controls. However, in the case of *Enterococcus faecalis*, there was also no growth in DMSO controls, i.e. were highly sensitive to DMSO. Because of that, the effect of *S. officinalis* against *Enterococcus faecalis* could not be determined. The extract had moderate activity for *Klebsiella pneumoniae* & *Pseudomonas aeruginosa* (MIC of 20mg/ml); and there was a significant growth in corresponding DMSO controls, whereas, *Escherichia coli* had low sensitivity to the extract as well as DMSO (MIC of 40mg/ml for the extract and no growth in the corresponding concentrations of DMSO, 10%), (Table-1 and Fig. 1-3).




Fig. 1. Growth of *Staphylococcus aureus* in different concentrations of methanolic extract of *S. officinalis* in Muller Hinton Agar, showing MIC of 5mg/ml




Fig. 2. Growth of MRSA in different concentrations of methanolic extract of *S. officinalis* in Muller Hinton Agar, showing MIC of 5mg/ml




Fig. 3. Growth of *Acinetobacter baumannii* in different concentrations of methanolic extract of *S. officinalis* in Muller Hinton Agar, showing MIC of 5mg/ml

Regarding fungi, *Candida albicans* were relatively more sensitive to the methanolic extract of *S. officinalis* (MIC of 20mg/ml) than *Aspergillus flavus* (Almost complete to around 50% inhibition of the growth of the extract (Table-2 and Fig. 4 & 5).

Table 2. Antifungal Activity of Methanolic Extract of *S. officinalis* (Maramia)

S.No	Name of Fungal ATCC Strains	Methanol Extract of Salvia officinalis (mg/ml)				
		40	20	10	5	2.5
1	<i>Candida albicans</i> ATCC	S	S	I	R	R
2	<i>Aspergillus flavus</i>	S	I	I	I	R



Fig. 4. Growth of *Candida albicans* in different concentrations of methanolic extract of *S. officinalis* in Sabouraud Dextrose Agar, showing MIC of 20 mg/ml




Fig. 5. Growth of *Aspergillus flavus* in different concentrations of methanolic extract of *S. officinalis* in Sabouraud Dextrose, showing almost complete to 50% inhibition of the growth with 40 to 5mg/ml of the extract

DMSO which is commonly used as a solvent in phytochemical studies, has been shown to possess antimicrobial activity and may interfere with the determination of MICs of plant extracts. Variable effects of DMSO against different microorganisms are reported in the literature. In general, 10-15% DMSO is reported to inhibit the growth of most organisms [5, 6], 1% showed no effect [7], whereas, between 2-8 gave variable results for different microorganisms [8].

In the present study, DMSO was used as a solvent for the methanolic extract of *S. officinalis* and as active control in the antimicrobial assays. DMSO 5% and below did not affect the growth of the organisms tested, except that DMSO 10 to 1.25% completely inhibited the growth of *Enterococcus faecalis* and partially inhibited *Aspergillus flavus*. Therefore, MIC values of the extract of *S. officinalis* between 2.5 to 20% observed for other microorganisms tested in the present study are least likely to be affected by the presence of DMSO. In a recent study also, *Staphylococcus aureus* was reported to be relatively more resistant and *Enterococcus faecalis* more sensitive to DMSO [9].

Staphylococcus aureus, gram positive cocci, reside normally in the upper respiratory tract, skin and intestinal mucosa. They can become pathogenic and may cause serious infections, like bacteremia, pneumonia, endocarditis and brain abscess, etc. Within days, they become resistant to commonly used drugs [10-12]. The results of the present study revealed that the methanolic extract of *S. officinalis* had promising activity against MRSA, which are becoming resistant to even Vancomycin, termed as vancomycin-resistant *Staphylococcus aureus* [13].

In an earlier study also *Staphylococcus aureus* was shown to be more sensitive to essential oil of *S. officinalis* (MIC 0.6 mg/ml) than gram negative bacilli, like *Salmonella* and *E. coli* (MICs 5 and 10 mg/ml, respectively). Their results also show that the essential oil was more effective than methanolic extract used in the present study, MICs for *Staphylococcus aureus* and MRSA were 5mg/ml and *Escherichia coli* 40 mg/ml³.

Acinetobacter baumannii is a Gram-negative bacillus, an emergent opportunistic pathogen and prone to cause

infections of respiratory tract, urinary tract, skin and eyes. Recently, it has been designated as a “red alert” human pathogen, because of its extensive antibiotic resistance [14]. The methanolic extract of *S. officinalis* was also very effective against multi-drug resistant *A. baumannii*. We could not find any study in the literature to compare the results of the extract against *Acinetobacter baumannii*.

Candida infections are the most frequently encountered nosocomial fungal infections [15]. In the present study, the methanolic extract of *S. officinalis* demonstrated a significant activity against *Candida albicans*. Essential oils of *S. officinalis* have also been demonstrated to inhibit *Candida albicans* [16].

Aspergillus flavus is commonly found in the air and is the most common cause of superficial infection. Aflatoxins released by *Aspergillus flavus* are the most potent hepatocarcinogenic natural compounds [17]. In the present study the methanolic extract of *S. officinalis* also demonstrated a moderate activity against *Aspergillus flavus*. In a previous study, food isolate of *Aspergillus flavus* was reported to be more resistant to essential oil of *S. officinalis* as compared to standard *Aspergillus niger* [3].

The present study has disclosed the potential of *S. officinalis* as a natural source for the production of new antibacterial and antifungal drugs. Other plants' extracts and their active principles have also been shown to possess antimicrobial activity. For example, various extracts of *Nigella sativa* and its active component, e.g. thymoquinone, were shown to be active against resistant *Streptococci*, *Staphylococci* and *Pseudomonas* species, as well as some opportunistic fungi and dermatophytes [18-20].

CONCLUSIONS

The methanolic extract of *S. officinalis* showed good activity against *Staphylococcus aureus*, MRSA, and the multidrug resistant *Acinetobacter baumannii*; moderate activity against *Candida albicans* and weak activity against some gram negative bacilli and *Aspergillus flavus*. Further researches are needed for the isolation of active principles from *S. officinalis* extracts and determination of their antimicrobial activities, which could lead to development of new drugs against resistant bacteria, like MRSA, and multidrug resistant *A. baumannii*.

ACKNOWLEDGMENTS

The authors gratefully acknowledge the financial support of the present study from the Deanship of the Scientific Research, Northern Border University, Arar, Saudi Arabia (Grant No. 5749-MED-2016-1-6-F).

REFERENCES

- [1] Bouajaj, S., Benyamna, A., Bouamama, H., Romane, A., Falconieri, D., Piras, A., Marongiu, B., 2013. Antibacterial, allelopathic and antioxidant activities of

- essential oil of *Salvia officinalis* L. growing wild in the Atlas Mountains of Morocco. *Nat Prod Res*, 27(18), pp. 1673-1676.
- [2] Cutillas, A. B., Carrasco, A., Martinez-Gutierrez, R., Tomas, V. and Tudela, J., 2017. *Salvia officinalis* L. Essential Oils from Spain: Determination of Composition, Antioxidant Capacity, Antienzymatic, and Antimicrobial Bioactivities. *Chem Biodivers*, 14(8). doi: 10.1002/cbdv.201700102. Epub 2017 Jul 1.
- [3] Ben-Khedher, M.R., Ben-Khedher, S., Ikbal Chaieb, I., Tounsi, S. and Hammami, M., 2017. Chemical composition and biological activities of *Salvia officinalis* essential oil from Tunisia. *EXCLI J*, 16, 160-173.
- [4] Weinstein, M.P., 2018. <https://clsi.org/standards/products/microbiology/documents/nm100/> ISBN:1-56238-838-X, 11 January.
- [5] Ansel, H.C., Norred, W.P. and Roth, I.L., 1969. Antimicrobial activity of dimethyl sulfoxide against *Escherichia coli*, *Pseudomonas aeruginosa*, and *Bacillus megaterium*. *Journal of Pharmaceutical Sciences*, 58(7), pp. fmi 777-910.
- [6] Bash, H., and Gadebusch, H.H., 1968. In Vitro antimicrobial Activity of Dimethylsulfoxide. *Applied Microbiology*, 16 (12), p. 1953.
- [7] Nostro, A., M.P. Germano, M.P., D'Angelo, A.V., Marino, A. and Cannatelli, M.A., 2000. Extraction methods and bioautography for evaluation of medicinal plant antimicrobial activity. *Letters in Applied Microbiology*, 30, pp. 379-384.
- [8] Hassan, A.S., 2014. The Antibacterial Activity of Dimethyl Sulfoxide (DMSO) with and without of Some Ligand Complexes of the Transitional Metal Ions of Ethyl Coumarin against Bacteria Isolate from Burn and Wound Infection. *Journal of Natural Sciences Research*, 4 (19), pp. 106-111. www.iiste.org. ISSN 2224-3186 (Paper) ISSN 2225-0921
- [9] Júnior, J.N.S.D. Bezerra, R.P. Porto, A.L.F, Herculano, P.N. and Marques, D.A.V., 2015. Effect of dimethyl sulfoxide (DMSO) in bacterial strains growth MIC determination. 23rd Congress of the International Union for Biochemistry and Molecular Biology, 44th Annual Meeting of the Brazilian Society for Biochemistry and Molecular Biology. Foz do Iguaçu, PRn, Brazil, 24-28 August.
- [10] Schenck, L.P., Surette, M.G. and Bowdish, D.M., 2016. Composition and immunological significance of the upper respiratory tract microbiota. *FEBS Letters*. 590 (21), pp. 3705-3720.
- [11] Wollina, U., 2017. Microbiome in atopic dermatitis. *Clinical, Cosmetic and Investigational Dermatology*, 10, pp. 51-56.
- [12] Liu, C.L., Bayer, A., Cosgrove, S.E., Daum, R.S., ridkin, S.K., Gorwitz, R. J., Kaplan, S.L., Karchmer, A.W., Levine, D.P., Murray, B.E., Rybak, M.J., Talan, D.A. and Chambers, H.F., 2011. Clinical Practice Guidelines by the Infectious Diseases Society of America for the Treatment of Methicillin-Resistant *Staphylococcus aureus* Infections in Adults and Children. *Clinical Infectious Diseases*, 52(3), pp. e18-e55.
- [13] Schito, G.C., 2006. The importance of the development of antibiotic resistance in *Staphylococcus aureus*. *Clin Microbiol Infect*, 12 (Suppl 1), pp. 3-8.
- [14] Howard, A., Donoghue, M.O., Feeney, A. and Sleator, R.D., 2012. *Acinetobacter baumannii*, An emerging opportunistic pathogen. *Virulence*. 3(3), pp. 243-250.
- [15] Karacaer, Z., Oncul, O., Turhan, V., Gorenk, L., and Ozyurt, M., 2014. A surveillance of nosocomial candida infections: epidemiology and influences on mortality in intensive care unit. *Afr Med J*, 19: 398 doi: 10.11604/pamj.2014.19.398.4960.
- [16] Cutillas, A.B., Carrasco, A., Martinez-Gutierrez, R., Tomas, V., Tudela, J., 2017. *Salvia officinalis* L. Essential Oils from Spain: Determination of Composition, Antioxidant Capacity, Antienzymatic, and Antimicrobial Bioactivities. *Chem Biodivers*, 14(8). doi: 10.1002/cbdv.201700102.
- [17] Hedayati MT, Pasqualotto AC, Warn PA, Bowyer P, Denning DW (2007). *Aspergillus flavus*: human pathogen, allergen and mycotoxin producer. *Microbiology*. 153(Pt 6): 1677-92.
- [18] Hanafi, M.S, Hatem, M.E., 1991. Studies on the antimicrobial activity of the *Nigella sativa* seed. (Black Cumin). *J Ethnopharmacol*, 31(2-3), pp. 275-278.
- [19] Khan, M.A., Ashfaq, M.K., Zuberi, H.S. and Zuberi A.H., 200). The in vivo antifungal activity of the aqueous extract from *Nigella sativa* seed, *Phytotherapy Research*, 17, pp. 183-186.
- [20] Aljabre, S.H.M., Randhawa, M.A., Akhtar, N., Alakloby, O.M., Alqurashi, A.M. and Aldossary, A., 2005. Antidermatophyte activity of ether extract of *Nigella sativa* and its active principle, thymoquinone. *J Ethnopharmacology*, 101, pp. 116-119