



Efficacy of *Equisetum Arvense* Extract Against Carbon Tetrachloride Induced Liver and Kidney Injury in Rats

Eman M. Ragheb^{1*}, Zaenah Z. Alamri²

¹ Agriculture Research Center, Regional Center for Food and Feed, Giza, Egypt.

² Biological Sciences Department, Faculty of Science, University of Jeddah, Jeddah, Saudi Arabia.

ABSTRACT

Medicinal plants are considered among the most important sources of antioxidants, which are proven to be highly effective against hepatic and nephrotoxicity of many chemical compounds. *Equisetum arvense* (*E. arvense*) plant family Equisetaceae has many uses in traditional medicine and possesses several pharmacological effects, most notably antioxidant effects. This study aimed first to assess the active constituents and antioxidants activities of *E. arvense* extract. Second to evaluate the protective action of *E. arvense* ethanolic extract against carbon tetrachloride (CCl₄) induced hepatic and renal toxicity in rats. This study was carried on 50 rats. Ten rats were served as a control group. Hepato and nephrotoxicity were induced in 10 rats by injection of CCl₄ (3ml/kg 2 times weekly for 2 weeks) and served as CCl₄ group. Thirty rats were sorted into 3 groups (n =10) and orally administered with *E. arvense* ethanolic extract (25, 50, and 75 mg/kg) for 2 weeks and then injected with CCl₄ for another 2 weeks. Results showed that *E. arvense* contains 3 main active constituents bergenin, nilotinib, and glafenin. It also has high total antioxidants and polyphenol contents. Administration of *E. arvense* at all dosage regimen significantly improved rats body weight gain percentage, liver functions (ALT, AST, ALP, total protein, and albumin), kidney functions (creatinine, urea, and uric acid), and lipid profiles (TC, TG, LDL-C, and HDL-C) matched to CCl₄ group. Oral feeding with *E. arvense* at all dosages regimen significantly ameliorates liver histopathology in favor of the highest *E. arvense* dose (75 mg/kg). Also, *E. arvense* at all dosages regimen significantly decreased lipid peroxidation products (MDA) matched to CCl₄ group. In conclusion, *E. arvense* exerted hepato and nephroprotective action as well as hypolipidemic effects against CCl₄-induced toxicity in rats. The mechanism may involve antioxidant effect and mitigation of lipid peroxidation.

Key Words: *Equisetum arvense*, hepatotoxicity, nephrotoxicity, dyslipidemia, antioxidant.

eIJPPR 2020; 10(5):43-52

HOW TO CITE THIS ARTICLE: Eman M. Ragheb, Zaenah Z. Alamri (2020). "Efficacy of *Equisetum arvense* Extract against Carbon Tetrachloride Induced Liver and Kidney Injury in Rats", International Journal of Pharmaceutical and Phytopharmacological Research, 10(2), pp.43-52.

INTRODUCTION

The liver is the largest and one of the most vital organs that functions to regulate detoxification and metabolism of exogenous and endogenous compounds [1, 2]. Hepatotoxicity is a prevalent health problem that represents 38% of all hepatic problems worldwide [3]. Toxic damage happened in the liver frequently compared to the other organs as all the absorbed substances first reach the liver to be metabolized and eliminated [4]. Carbon tetrachloride (CCl₄) has long been known as a toxicant in animal models for made of acute and chronic liver and renal injuries. CCl₄ model has been utilized in numerous *in vivo* and *in vitro* toxicological researches [5, 6]. It induces lipid peroxidation

and lowering antioxidant enzyme activities [7]. Although, when the balance between the oxidative stress and the antioxidant was impaired, the liver is the utmost organ to be in danger for tissue injury associated with the reactive oxygen species [8, 9].

Many *in vitro* and *in vivo* researches assess natural therapeutic medicine for curing and protecting several debilitating diseases [10, 11]. Nowadays, there is a growing interest in medicinal plant usage. *Equisetum arvense* (*E. arvense*) belongs to the Equisetaceae family famous as field horsetail. It is a plant with a wide prospectus. *E. arvense* contains numerous flavonoids, alkaloids, phenol, phenolic, petrosins, triterpenoids, sterols, saponins, phytosterols, tannin, volatile

Corresponding author: Eman M. Ragheb

Address: Agriculture Research Center, Regional Center for Food and Feed, Giza, Egypt.

E-mail: ✉ emanragheb07 @ gmail.com

Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 04 July 2020; **Revised:** 23 September 2020; **Accepted:** 25 September 2020



oils, minerals, ascorbic acid, silicic acid, and many other biologically active constituents [12-14].

In folk medicine, *E. arvense* is used to treat tuberculosis, pulmonary, gastric, hemorrhages, rheumatic diseases, gout, wound healing, ulcers, and fractures. It possessed numerous pharmacological properties including antimicrobial, antioxidant, anticancer, and anti-inflammatory actions [13, 15-20]. Numerous researches have documented the hypoglycemic action of *E. arvense* extract in diabetic models [21, 22]. The hepatoprotective action of the *E. arvense* extract versus the hepatitis model made by tetrachloromethane has been confirmed [23]. Moreover, a hepatoprotective effect of the phenolic petrosins and flavonoids separated from *E. arvense* has been documented [24]. Also, *E. arvense* extract showed a renoprotective and pressure-lowering impact in an experimental model of chronic kidney diseases [25].

As far as we know, no previous researches reported the liver and kidney protective efficacy of *E. arvense* ethanolic extract versus CCl₄. Subsequently, the current research was performed to estimate the preventive effect of ethanolic extract of *E. arvense* against CCl₄ produced hepatic and renal damage in rats.

MATERIAL AND METHODS

Plants, chemicals, and animals

E. arvense plant was obtained from Haraz for herbs and medicinal plants Company, Cairo, Egypt. Carbon tetrachloride (CCl₄) was bought from Sigma-Aldrich (St. Louis, USA). All chemicals were bought from EL-Gomhoria, and Biodiagnostic, Egypt. Fifty male Sprague Dawley rats (n=50) of 200 g ± 10 average body weights, were obtained from Helwan Experimental Animals Farm, Giza, Egypt.

Gas chromatography-mass spectroscopy (GC-MS) analysis of *E. arvense* active constituents

The assay was performed utilizing a GC-MS (Agilent Technologies 7890A) connected to a mass-specific detector (MSD, Agilent 7000). Helium was the carrier gas. The recognition of constituents was carried out by comparing their mass spectra and retention time with the library of authentic compounds (NIST and WILEY) [26].

Preparation of *E. arvense* ethanolic extract and estimation of its antioxidant content

The ethanolic extract of *E. arvense* was prepared according to Safiyeh et al. [21]. The phosphomolybdenum assay was adopted to estimate the antioxidant content [27], and Folin-Ciocalteu assay was adopted to estimate the total phenols content [28].

Induction of hepato-nephrotoxicity

Rats were subcutaneous (s.c.) injected 3ml/kg of 50% v/v CCl₄/olive oil (2 times weekly for 2 weeks) to induce liver and renal toxicity according to Hismiogullari et al. [6]; Jayasekhar et al. [29].

Experimental design

Rats were housed in suitable cages under optimal laboratory situation and fed a standard rats chow one week for adaptation [30]. They were sorted into five groups (10 rats). Group 1 (Control): rats s.c. injected with olive oil. Group 2 (CCl₄): rats s.c. injected with CCl₄. Groups 3-5 (*E. arvense* + CCl₄): rats received orally 25, 50, and 75 mg/kg *E. arvense* ethanolic extract, respectively, for two weeks before CCl₄ s.c. injection [21].

Feed intake was recorded over the four weeks, initial body weight (IBW), and final body weight (FBW) estimated to calculate body weight gain percent (BWG%). At the end of the experiment, rats in all groups were anesthetized, and blood was collected from the aorta and centrifuged 10 minutes at 3000 rpm. The serum was then collected and preserved at -80 °C for the measurement of the biochemical parameters. The liver and kidney were detached and washed with normal saline, then weighted.

Measurement of liver enzymes

In serum, the activities of aspartate aminotransaminase (AST), alanine aminotransaminase (ALT), and alkaline phosphatase (ALP) were measured using colorimetric assay kits as claimed by the manufacturer.

Measurement of total proteins and albumin

Serum total proteins and albumin were estimated using colorimetric assay kits as claimed by the manufacturer.

Measurement of renal functions

Creatinine, urea, and uric acid were measured in serum using colorimetric assay kits as claimed by the manufacturer.

Measurement of lipids

Triglycerides (TG), total cholesterol (TC), high-density lipoprotein cholesterol (HDL-C), and low-density lipoprotein-cholesterols (LDL-C) were estimated by colorimetric assay kits as claimed by the manufacturer.

Measurement of malondialdehyde

Malondialdehyde (MDA) was measured using ELISA kits, as claimed by the manufacturer.

Histopathological examination

The formalin-fixed liver tissues were dehydrated in graded alcohol, clarified in xylene, and paraffin-embedded. Then 3-5 µm thick sections were cut, stained with hematoxylin and eosin (H & E), and scanned *via* a light microscope.

Statistical calculations

Values are displayed as mean ± standard deviation (SD) and statistically tested for significance by ANOVA test preceded by LSD multiple comparison test utilizing SPSS software program, version 24 (p-value ≤ 0.05 was statistically significant).

RESULTS

Active constituents of *E. arvense*

The GC-MS analysis of *E. Arvense* is presented in Figure 1 and Table 1. The results showed that *E. arvense* contains several active constituents. There were 3 main compounds present in *E. arvense*, including bergenin (12.73%), nilotinib (11.55%), and glafenin (9.71%). Followed by gardenin (6.42%), dimethylfraxetin (5.86%), sepiapterin (5%), 3-(3,4-Dimethoxyphenyl)-4-methylcoumarin (4.97%), and (S)-(-)-Citronellic acid (4.66%).




Figure 1: Gas chromatography-mass spectroscopy spectra of *E. arvense* active contents

Table 1: Active constituents of *E. arvense* (gas chromatography-mass spectroscopy)

Constituents	RT (min)	Concentrations (%)
(S)-(-)-Citronellic acid	11.407	4.66
6,7-Dimethoxy-4-ethylcoumarin	12.515	1.77
Sepiapterin	12.997	5
Tetrahydro-L-biopterin	13.136	2.62
Retinyl propionate	13.271	2.45
3,5,3',5'-Tetra-tert-butylidiphenquinone	13.393	1.29
2'-Hydroxy-3,4,4',5-tetramethoxychalcone	13.798	1.09
Nilotinib	14.001	11.55
Dimethylfraxetin	14.622	5.86
3-(3,4-Dimethoxyphenyl)-4-methylcoumarin	15.095	4.97
7,3',4',5'-Tetramethoxyflavanone	15.244	1.93
Bergenin	15.428	12.73
3,2',4',5'-Tetramethoxyflavone	15.671	1.92
all trans-Retinal	15.816	1.51
Astilbin	16.014	4.79
Cholic acid	16.351	2.73
Glafenin	16.986	9.71
Gardenin	17.689	6.42
Linoleic acid	18.247	2.07

3-Hydroxy-7,8,2'-trimethoxyflavone	18.715	2.05
Isovitexin	19.04	0.96
Dodecanedioic acid	19.431	1.51
Phytanic acid	20.634	4.39
9-cis-Retinoic acid	21.872	1.58
Quercetin 3,5,7,3',4'-pentamethyl ether	22.926	3.51
Irbesartan	23.106	0.92
Non-identified compounds	> 23.2	0.01

Antioxidant contents of *E. arvense*

The total antioxidants content of *E. arvense* amounted to 2020.6 ± 20.0 mg/ 100 g ascorbic acid, where the total phenolic constituents amounted to 218.0 ± 18.0 mg/ 100 g gallic acid (Table 2).

Table 2: Total antioxidants and total phenols of *E. arvense*

Antioxidant constituents	Mean ± SD
Total antioxidants (mg/ 100 g ascorbic acid)	2020.6 ± 20.00
Total phenols (mg/ 100 g gallic acid)	218.0 ± 18.00

Values were presented as the mean of three replicates ± SD.

Impact of *E. arvense* on biological evaluation and organs (liver and renal) weight on CCl₄-induced toxicity in rats

Table 3: Impact of *E. arvense* on some biological parameters on CCl₄-induced toxicity in rats

Experimental groups	IBW (g)	FBW (g)	BWG %	FI (g/day/rat)
Control	202.4 ± 4.27	265.8 ± 10.43	31.32 ± 7.49	24.70 ± 2.96
CCl ₄	202.7 ± 4.90	187.2 ± 4.32 ^a	-7.65 ± 1.94 ^a	23.74 ± 4.14
<i>E. arvense</i> 25 mg kg+ CCl ₄	201.2 ± 5.27	220.3 ± 19.97 ^b	9.49 ± 3.62 ^b	24.52 ± 2.41
<i>E. arvense</i> 50 mg kg+ CCl ₄	200.1 ± 5.80	240.5 ± 5.99 ^{b,c}	20.19 ± 4.78 ^{b,c}	24.75 ± 3.24
<i>E. arvense</i> 75 mg kg+ CCl ₄	200.6 ± 5.76	254.3 ± 9.78 ^{b,c,d}	26.77 ± 5.09 ^{b,c,d}	25.36 ± 5.45

Values were offered as mean ± SD (n=10). Values were offered as mean ± SD (n=10). Results were significantly varied (p ≤ 0.05) from ^acontrol, ^bCCl₄ group, ^c*E. arvense* 25 mg/kg+CCl₄ group, ^d*E. arvense* 50 mg/kg+CCl₄ group.

Table 4: Impact of *E. arvense* on liver and renal weight on CCl₄-induced toxicity in rats

Experimental groups	Liver (g)	Renal (g)
Control	3.92 ± 0.43	0.61 ± 0.04
CCl ₄	5.07 ± 0.74 ^a	0.73 ± 0.08 ^a
<i>E. arvense</i> 25 mg kg+ CCl ₄	4.94 ± 0.64	0.70 ± 0.05
<i>E. arvense</i> 50 mg kg+ CCl ₄	4.84 ± 0.55	0.71 ± 0.06
<i>E. arvense</i> 75 mg kg+ CCl ₄	4.02 ± 0.45 ^{b,c,d}	0.70 ± 0.05

Values were offered as mean ± SD (n=10). Results were significantly varied (p ≤ 0.05) from ^a control, ^bCCl₄ group, ^c*E. arvense* 25 mg kg+CCl₄ group, ^d*E. arvense* 50 mg kg+CCl₄ group.

Impact of *E. arvense* on liver enzymes function on CCl₄-induced toxicity in rats

The rats injected with CCl₄ displayed a significant rise (p ≤ 0.001) in hepatic enzymes (ALT, AST, and ALP) matched to the control. *E. arvense* oral ingestion significantly decreased hepatic enzymes (ALT, AST, and ALP) in a

Administration of CCl₄ to rats significantly decreased FBW and BWG% (p ≤ 0.001), with significantly raised (p ≤ 0.001) liver and renal weight-matched to the control group. *E. arvense* oral ingestion significantly increased the FBW and BWG% in a dose-dependent way (p ≤ 0.001) matched to CCl₄ intoxicated rats. Besides, the ingestion of *E. arvense* 75 mg/kg significantly decreased the liver and renal weight matched to CCl₄ group, *E. arvense* 25 mg/kg+CCl₄, and *E. arvense* 50 mg/kg+CCl₄ (p ≤ 0.001). Concerning FI, there was a non-significant decline between CCl₄ intoxicated rats and control rats. Moreover, the ingestion of *E. arvense* in all doses non-significantly increased FI matched to CCl₄ group Tables 3 and 4.

dose-dependent way (p ≤ 0.001) matched to the CCl₄ intoxicated rats. Besides, there was a significant variation between the *E. arvense* 25 mg/kg and the other two doses. Also, there was a significant variation between the *E. arvense* 50 mg/kg+CCl₄ group and the *E. arvense* 75mg/kg+CCl₄ group Figure 2.




Figure 2: Impact of *E. arvense* on liver enzymes function on CCl₄-induced toxicity in rats. Values were offered as mean±SD (n=10). Results were significantly varied ($p \leq 0.05$) from ^acontrol, ^bCCl₄ group, ^c*E. arvense* 25 mg/kg+CCl₄ group, ^d*E. arvense* 50 mg/kg+CCl₄ group.

Impact of *E. arvense* on total protein and albumin on CCl₄-induced toxicity in rats

The rats injected with CCl₄ displayed significant decline ($p \leq 0.001$) in total protein and albumin values matched to the control group. *E. arvense* oral ingestion significantly increased total protein and albumin values in a dose-

dependent ($p \leq 0.001$) matched to the CCl₄ intoxicated rats. Besides, there was a significant variation between the *E. arvense* 25 mg/kg and the other two doses. Also, there was a significant variation between the *E. arvense* 50 mg/kg+CCl₄ group and the *E. arvense* 75 mg/kg+CCl₄ group Figure 3.




Figure 3: Impact of *E. arvense* on total protein and albumin on CCl₄-induced toxicity in rats. Values were offered as mean ± SD (n=10). Results were significantly varied ($p \leq 0.05$) from ^acontrol, ^bCCl₄ group, ^c*E. arvense* 25 mg/kg+CCl₄ group, ^d*E. arvense* 50 mg/kg+CCl₄ group.

Impact of *E. arvense* on kidney function on CCl₄-induced toxicity in rats

The rats injected with CCl₄ displayed a significant rise ($p \leq 0.001$) in serum creatinine, uric acid, and urea levels matched to the control group. *E. arvense* oral ingestion significantly decreased serum creatinine, uric acid, and urea values in a dose-dependent way ($p \leq 0.001$) matched

to the CCl₄ intoxicated rats. Concerning uric acid, there was a significant variation between the *E. arvense* 25 mg/kg and the other two doses. Also, there was a significant variation between the *E. arvense* 50 mg/kg+CCl₄ group and the *E. arvense* 75 mg/kg+CCl₄ group Figure 4.




Figure 4: Impact of *E. arvense* on kidney function on CCl₄-induced toxicity in rats. Values were offered as mean ± SD (n=10). Results were significantly varied (p ≤ 0.05) from ^a control, ^bCCl₄ group, ^c*E. arvense* 25 mg/kg+CCl₄ group, ^d*E. arvense* 50 mg/kg+CCl₄ group.

Impact of *E. arvense* on lipid profile parameters on CCl₄-induced toxicity in rats

The rats injected with CCl₄ displayed a significant elevation (p ≤ 0.001) in serum values of TC, TG, and LDL-C, concurrent with a significant elevation (p ≤ 0.001) in serum HDL-C concentration matched to control. *E.*

arvense oral ingestion significantly improves all lipid profile parameters in a dose-dependent way (p ≤ 0.001) matched to CCl₄ intoxicated rats. Besides, there was significant variation between the *E. arvense* 25 mg/kg+CCl₄ group and the *E. arvense* 75 mg/kg+CCl₄ group in all tested lipid parameters Table 5.

Table 5: Impact of *E. arvense* on lipid profile parameters on CCl₄-induced toxicity in rats

Experimental groups	TC(mg/dl)	TG (mg/dl)	LDL-C (mg/dl)	HDL-C (mg/dl)
Control	81.52 ± 3.89	73.84 ± 5.58	30.06 ± 5.87	54.11 ± 8.26
CCl ₄	115.31 ± 7.22 ^a	147.04 ± 19.06 ^a	49.39 ± 10.79 ^a	33.85 ± 9.04 ^a
<i>E. arvense</i> 25 mg kg+ CCl ₄	89.30 ± 5.15 ^b	90.97 ± 9.88 ^b	35.06 ± 4.02 ^b	40.16 ± 9.15 ^b
<i>E. arvense</i> 50 mg kg+ CCl ₄	86.68 ± 5.88 ^b	82.72 ± 5.15 ^b	30.07 ± 9.41 ^b	42.13 ± 6.14 ^b
<i>E. arvense</i> 75 mg kg+ CCl ₄	82.09 ± 6.24 ^{b,c}	78.01 ± 5.45 ^{b,c}	23.28 ± 7.38 ^{b,c}	50.02 ± 12.62 ^{b,c}

Values were offered as mean ± SD (n=10). Results were significantly varied (p ≤ 0.05) from ^a control, ^bCCl₄ group, ^c*E. arvense* 25 mg kg+CCl₄ group.

Impact of *E. arvense* on serum malondialdehyde on CCl₄-induced toxicity in rats

The rats injected with CCl₄ displayed a significant rise (p ≤ 0.001) in MDA matched to control. *E. arvense* oral

ingestion significantly decreases MDA in a dose-dependent way (p ≤ 0.001) matched to the CCl₄ intoxicated rats. Besides, there was a significant decrease between the *E. arvense* 75 mg/kg+CCl₄ group and the *E. arvense* 25 mg/kg+CCl₄ group Figure 5.




Figure 5: Impact of *E. arvense* on serum malondialdehyde (MDA) on CCl₄-induced toxicity in rats. Values were offered as mean ± SD (n=10). Results were significantly varied ($p \leq 0.05$) from ^a control, ^bCCl₄ group, ^c*E. arvense*25 mg/kg + CCl₄ group.

Impact of *E. arvense* on liver tissue histopathology

The histological results of hepatic sections of the control rats displayed a normal histological structure of hepatic lobules (Figure 6A). Hepatic sections of rats from the CCl₄ group displayed hepatic sinusoids congestion, Kupffer cell activation, presence of karyomegaly, binucleated cells, sporadic hepatocytes necrosis, and inflammatory leukocytes infiltration (Figure 6B&C). Liver sections of CCl₄ pretreated with *E. arvense*25 mg/kg displayed regeneration

nodules separated by fibrous septa with slight loss of normal architecture (Figure 6D). Liver sections of CCl₄ pretreated with *E. arvense*50 mg/kg displayed proliferated hepatocytes replacing the degenerated ones (Figure 6E). Liver sections of CCl₄ pretreated with *E. arvense*75 mg/kg displayed apparent normal hepatic lobule structure where the portal tract was surrounded by proliferated hepatocytes replacing the degenerated ones (Figure 6F).




Figure 6. Effect of *E. arvense* on hepatic tissue histopathology displayed in hepatotoxic rats. The histological examination of liver sections of control displayed a healthy histological appearance of hepatic lobules (A). The liver of rats from the CCl₄ group displayed hepatic sinusoid congestion, kupffer cell activation, presence of karyomegaly, binucleated cells, sporadic cell necrosis (B), along with sporadic hepatocytes necrosis and inflammatory leukocytes infiltration (C). Liver sections of *E. arvense*25 mg/kg +CCl₄group displayed regeneration nodules separated by fibrous septa with loss of normal architecture of the liver (D). Liver sections of *E. arvense*50 mg/kg+CCl₄group displayedproliferated hepatocytes replacing the degenerated ones (E). Liver sections of *E. arvense*75 mg/kg +CCl₄group displayed a portal tract surrounded by proliferated hepatocytes replacing the degenerated ones (F)(H&E X 200).

DISCUSSION

The liver regulates numerous functions of the body [31]. CCl₄ is well documented to induce hepatic and nephrotoxicity via oxidative stress mechanisms [1]. The

extract *E. arvense* has many bioactive components that quench free-radicals and possess antioxidant properties [15, 17, 18]. Since the damaging effects of CCl₄ involve oxidative stress, this study postulated that *E. arvense* extract

might protect the liver and renal versus CCl₄ induced toxicity.

In this study, the CCl₄ induced significant decreases in FBW, BWG%, with significant increases in liver and kidney weight. The obtained results agree with Ezejindu et al. [32], who revealed that injection with CCl₄ caused a significant loss of body weight of rats concomitant with a relative increase in liver weight. This increase in liver weight was not growth, but inflammation occurs *via* CCl₄. Besides, recently Ullah et al. [33] attributed the reduction in body weight to the decreased feed intake, which was observed in the present study.

Liver enzymes are utilized to estimate the liver cell damage, while total protein is usually applied to estimate the hepatic activity [1, 10]. This study showed that CCl₄-induced severe hepatic and renal destruction as evidence by significant increases in hepatic enzyme levels (ALT, AST, and ALP) coupled with significant increases in renal function levels (creatinine, urea, and uric acid), as well as disturbing in serum total protein and albumin values. The biochemical findings confirmed with the histopathological results of CCl₄ intoxicated rat's liver sections, which showed the occurrence of focal hepatocytes necrosis together with the infiltration of inflammatory cells. These findings pointed to kidney and liver impairment, cellular infiltration, injury, and disturbed cell membrane integrity in the kidney and liver.

The obtained results agree with several studies [3, 33-36] who showed that CCl₄ treated mice had extended necrosis around the central vein and vacuole formation, thus indicated increased hepatic injury. The hepatotoxicity following CCl₄ application explained *via* membrane damage and significant disturbance in renal and liver tissues induced by CCl₄, which is metabolized by cytochrome p450-2E1 to trichloromethyl radicals that begin free radical-induced lipid peroxidation, thus cause hepatic and renal damage [33, 37].

The nephrotoxic effect of CCl₄ was confirmed in several previous studies [38, 39]. Adewole et al. [40] revealed that CCl₄ caused severe kidney damage as estimated by elevated serum creatinine, blood urea nitrogen, and urea concentration, which was explained through CCl₄-induced oxidative stress that boosts the production of diverse vasoactive substances which directly disturbed the kidney function by prompting renal vasoconstriction and impaired the glomerular filtration [41].

Furthermore, CCl₄- induced overt oxidative stress was marked by significant elevation of serum MDA level, thus confirmed in several types of researches [10, 34, 40]. Aziz et al. [7] revealed that many xenobiotics, including CCl₄ toxicity induced *via* the free radicals' production, which are toxic and implicated in the pathophysiology of ailments. Recently, Ullah et al. [33] reported that the CCl₄ brought a high level of MDA and noticeable exhaustion of endogenous antioxidant molecules. The CCl₄ intoxication

formed free radicals that induced a cascade of actions inducing in its toxicity [38, 39].

Besides, in this study CCl₄ injection caused a significant increase in lipid profile parameters. Recently Ullah et al. [33] and Elsayy et al. [35] revealed that CCl₄ injection induced a statically significant increase in liver and serum lipids (free fatty acids, TC, total lipids, and TG), while decrease serum HDL-C. CCl₄-induced oxygen-free radicals generation, which catalyzes the oxidation of LDL that caused cell injury [42]. Besides, alteration of lipid profile considered a causal factor for oxidative stress and excessive MDA as found in this study.

The pretreatment effect of *E. arvense* extract (25, 50, and 75 mg/kg) is remarkably protected against both liver and renal injury caused by CCl₄. There were significant decreases in serum liver and kidney markers, as well as improve serum protein and albumin levels. The biochemical results confirmed by the histopathological investigation, which showed that pretreatment of intoxicated rats with *E. arvense* extract significantly protects the liver as minimal changes were seen. The high dose of *E. arvense* (75 mg/kg) was the most effective. *E. arvense* extract administration maintains liver and renal function homeostasis *via* acting as a membrane stabilizing agent through its active antioxidant constituents effects, which was confirmed by the analysis of the active constituents of *E. arvense* by GC-MS done in the current study. *E. arvense* comprises antioxidant activity through its numerous biological active constituents including flavonoids, phenolic, and phytosterols [13, 14, 43].

The hepatoprotection activity of *E. Arvense* herbs extract was investigated in a model of acute hepatitis produced by tetrachloromethane. The results offered that the extract protected the membrane through antioxidant action. This was displayed through lowered liver enzymes, total bilirubin, and lipid peroxidation products, besides the absence of reduced endogenous alpha-tocopherol and glutathione-based enzymes [23]. Oh, et al. [24] showed that the methanolic extract of *E. arvense* produced a marked protective action against tacrine-prompted cytotoxicity in the Hep G2 cell line. *E. arvense* extract decline serum level of MDA induced by CCl₄ injection. This effect was further explained by *E. arvense* phytochemical antioxidant constituents, which possesses a potent radical scavenging ability [44-47]. To the best of our knowledge, no previous study was reported concerning the protective role of *E. arvense* extract against the toxic effects induced by CCl₄ in rats, and our study is the first in this line.

CONCLUSION

The biochemical and the histopathological findings of this study concluded that *E. arvense extract* dose-dependently protects against hepatotoxicity, nephrotoxicity, and hyperlipidemia induced by CCl₄ in rats. The mechanism

behind *E. arvense* action could be explained by its antioxidant and free radicals scavenging efficacy.

REFERENCES

- [1] Al-Seeni MN, El Rabey HA, Zamzami MA, Anefayee AM. The hepatoprotective activity of olive oil and *Nigella sativa* oil against CCl₄-induced hepatotoxicity in male rats. *BMC Complement Altern Med.* 2016; 16(438): 1–14.
- [2] Ozougwn JC. Physiology of the liver. *IntJ Res in Pharm and Biosciences.* 2017; 4(8):13–24.
- [3] Khan RA, Khan MR, Sahreen S. CCl₄-induced hepatotoxicity: protective effect of rutin on p53, CYP2E1, and the antioxidative status in rat. *BMC Complement Altern Med.* 2012; 12(178):1–6.
- [4] Tavga AA. Effect of benfotiamine on hepatic tissue levels of free calcium, copper, iron, and zinc during CCl₄-induced hepatotoxicity in rats. *Zanco J. Med. Sci.* 2011; 15(2): 10–14.
- [5] Hartmut J. Toxic responses of the liver. In: Casarett and Doull's Toxicology: The basic science of poisons. 7th edition. 2008:557–82.
- [6] Hismiogullari AA, Hismiogullari SE, Karaca O, Sunay FB, Paksoy S, Can M, Kus I, SeyrekK, Yavuz O. The protective effect of curcumin administration on carbon tetrachloride (CCl₄)-induced nephrotoxicity in rats. *Pharmacol Rep.* 2015; 67(3):410–6.
- [7] Aziz TA, Ahmed ZA, Juma'a KM, Abdulrazzaq MH, Hussain SA. Study of the protective effects of benfotiamine against CCl₄-induced hepatotoxicity in rats. *Iraqi J Pham Sci.* 2009; 18:47–53.
- [8] McGill MR, Jaeschke H. Metabolism and disposition of acetaminophen: recent advances in relation to hepatotoxicity and diagnosis. *Pharm Res.* 2013; 30(9):2174–87.
- [9] Ritesh K, Suganya A, Dileepkumar H, Rajashekar Y, Shivanandappa T. A single acute hepatotoxic dose of CCl₄ causes oxidative stress in the rat brain. *Toxicol Rep.* 2015; 2:891–5.
- [10] Rusu M, Tamas M, Puica C, Roman I, Sabadas M. The hepatoprotective action of ten herbal extracts in CCl₄ intoxicated liver. *Phytother Res.* 2005; 19(9):744–9.
- [11] Zhao L, Pickering G. Paracetamol metabolism and related genetic differences. *Drug Metab Rev.* 2011; 43(1):41–52.
- [12] Radulović N, Stojanović G, Palić R. Composition and antimicrobial activity of *Equisetum arvense* L. essential oil. *Phytotherapy Res.* 2006; 20(1):85–8.
- [13] Mimica-Dukic N, Simin N, Cvefic J, Jovin E, Orcic D, Bozin B. Phenolic compounds in field horsetail (*Equisetum arvense* L) as natural antioxidants. *Molecule.* 2008; 13: 1455–64.
- [14] Navdeep SS, Sarabjit K, Divneet C. *Equisetum arvense*: pharmacology and phytochemistry- A review. *Asian J Pharmaceut and Clin Res.* 2010; 3(3):146–9.
- [15] Trouillasa P, Callistea CA, Allaisc DP, Simonb A, Marfaka A, Delageb C, Durouxa JL. Antioxidant, anti-inflammatory, and antiproliferative properties of sixteen water plant extracts used in the Limousin countryside as herbal teas. *Food Chemist.* 2003; 80: 399–407.
- [16] AlexandruV, Petrusca DN, Gille E. Investigation of pro-apoptotic activity of *Equisetum arvense* L. water extract on human leukemia U 937 cells. *Romanian Biotechnol Letters.* 2007; 12(2):3139–47.
- [17] Stajner D, Popović BM, Canadanović-Brunet J, Anackov G. Exploring *Equisetum arvense* L., *Equisetum ramosissimum* L. and *Equisetum telmateia* L. as sources of natural antioxidants. *Phytother Res.* 2009; 23(4):546–50.
- [18] Cetojevic-Simin DD, Canadanovic-Brunet JM, Bogdanovic GM, Djilas SM, Cetkovic GS, Tumbas VT, Stojiljkovic BT. Antioxidative and antiproliferative activities of different horsetail (*Equisetum arvense* L.) extracts. *J Med Food.* 2010; 13(2): 452–9.
- [19] Geetha RV, Lakshmi T, Roy A. In vitro evaluation of antibacterial activity of *Equisetum arvense* Linn on urinary tract pathogens. *Int J Pharm and Pharmaceutical Sci.* 2011; 3(4): 323–5.
- [20] Sinha NS. In vitro antibacterial activity of ethanolic extract of *Equisetum arvense* L. *IntJPharmaceut and Biol Res.* 2012; 3(1): 19–21.
- [21] Safiyeh S, Fathallah FA, Vahid N. The effect of *Equisetum arvense* L. (Equisetaceae) in histological changes of pancreatic beta-cells in streptozotocin-induced diabetic in rats. *Pak J BiolSci.* 2007; 10(23):4236–40.
- [22] Al-Snafi AE. The pharmacology of *Equisetum arvense*- A review. *IOSR J Pharm.* 2017; 7(2): 31–42.
- [23] Katikova OI, Kostin I, Tishkin VS. Hepatoprotective effect of plant preparations. *Eksp Klin Farmakol.* 2002; 65: 41–3.
- [24] Oh H, Kim DH, Cho JH, Kim YC. Hepatoprotective and free radical scavenging activities of phenolic petrosins and flavonoids isolated from *Equisetum arvense*. *J Ethnopharmacol.* 2004; 95:421–4.
- [25] Ülle P, Ingrid K, Mai O. Renoprotective and blood pressure lowering impact of *Equisetum arvense* and Viscumalbumtherapy in experimental model of chronic kidney disease. *World J Cardiovascular Dis.* 2018; 8: 545–56.
- [26] Santana PM, Miranda M, Payrol JA, SilvaM, Hernández V, Peralta E. Gas chromatography-mass spectrometry study from the leaves fractions obtained

- of *Vernonanthura patens* (Kunth) H. *Rob Int J Org Chem.* 2013;03(02):105–9.
- [27] Prieto P, Pineda M, Aguilar M. Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: Specific application to the determination of vitamin E. *Anal Biochem.* 1999; 269(2):337–41.
- [28] Singleton VL, Orthofer R, Lamuela-Raventós RM. Analysis of total phenols and other oxidation substrates and antioxidants by means of folin-ciocalteu reagent. *Methods Enzymol.* 1999; 299:152–78.
- [29] Jayasekhar P, Mohanan PV, Rathinam K. Hepatoprotective activity of ethyl acetate extract of *Acacia catechu*. *Indian J of Pharmacol.* 1997; 29(6):426–8.
- [30] Reeves PG, Nielsen FH, Fahey GC. AIN-93 purified diets for laboratory rodents: final report of the American Institute of Nutrition ad hoc writing committee on the reformulation of the AIN-76A rodent diet. *J Nutr.* 1993;123 (11):1939–51.
- [31] Bertoni G, Trevisi E, Han X, Bionaz M. Effects of inflammatory conditions on liver activity in puerperium period and consequences for performance in dairy cows. *J Dairy Sci.* 2008;91(9):3300–10.
- [32] Ezejindu DN, Ulolene GC, Ihentuge CJ. Evaluation of toxicological effects of carbon tetrachloride on liver enzymes of Wistar rats. *IntScientifTechnol Res.* 2013; 2 (12):372-4.
- [33] Ullah H, Khan A, Baig MW, Ullah N, Ahmed N, Tipu MK, Ali H, Khan S. Poncirin attenuates CCL4-induced liver injury through inhibition of oxidative stress and inflammatory cytokines in mice. *BMC Complement Altern Med and Therapy.* 2020; 20 (115): 1–14.
- [34] Jayakumar T, Ramesh E, Geraldine P. Antioxidant activity of the oyster mushroom, *Pleurotus ostreatus*, on CCl4-induced liver injury in rats. *Food Chem Toxicol.* 2006;44(12):1989–96.
- [35] Elsayy H, Badr GM, Sedky A, Abdallah BM, Alzahrani AM, Abdel-Moneim AM. Rutin ameliorates carbon tetrachloride (CCl4)-induced hepatorenal toxicity and hypogonadism in male rats. *PeerJ.* 2019; 7:1–20.
- [36] Chao-Lin Y, Yu-Shih L, Keng-Fan L, Wen-Huang P, Chih-Ming H. Hepatoprotective mechanisms of taxifolin on carbon tetrachloride-induced acute liver injury in mice. *Nutrients.* 2019; 11(2655): 1–7.
- [37] Aleynik SI, Leo MA, MaX, Aleynik MK, Lieber CS. Polyenylphosphatidylcholine prevents carbon tetrachloride-induced lipid peroxidation while it attenuates liver fibrosis. *J Hepatol.* 1997; 27(3): 554–61.
- [38] Khan MR, Zehra H. Amelioration of CCl4-induced nephrotoxicity by *Oxalis corniculata* in rat. *Experiment Toxicol Pathol.* 2013; 65(3):327–34.
- [39] Jan S, Khan MR. Protective effects of *Monotheba buxifolia* fruit on renal toxicity induced by CCl4 in rats. *BMC Complement Altern Med.* 2016; 16289: 1–15.
- [40] Adewole SO, Salako AA, Doherty OW, Naicker T. Effect of melatonin on carbon tetrachloride induced kidney injury in Wistar rats. *African J of Biomedical Res.* 2007; 10:153–64.
- [41] Garcia-Cohen EC, Marin J, Diez-Picazo LD, Baena AB, Salaices M, Rodriguez-Martinez MA. Oxidative stress induced by tertbutylhydroperoxide causes vasoconstriction in the aorta from hypertensive and aged rats: role of cyclooxygenase 2 isoform. *J Pharmacol Exp Ther.* 2000; 293(1): 75–81.
- [42] Kosola J, Vaara JP, Ahotupa M, Kyröläinen H, Santtila M, Oksala N, Atalay M, Vasankari T. Elevated concentration of oxidized LDL together with poor cardiorespiratory and abdominal muscle fitness predicts metabolic syndrome in young men. *Metabol: Clin Experiment.* 2013; 62(7):992–9.
- [43] Ozay Y, Cayci MK, Guzel-Ozay S, Cimbiz A, Gurlek-Olgum E, Ozyurt MS. Effects of *Equisetum arvense* ointment on diabetic wound healing in rats. *Wounds.* 2013;25(9):234–41.
- [44] Park EY, Jeon H. Antioxidant and anti-inflammatory activities of *Equisetum hyemale*. *Nat Prod Sci.* 2008; 14: 239–43.
- [45] Fu LN, Chen LY, Liu RH, Chen DF. Chemical composition of *Equisetum debile*. *J Chin Med Mat.* 2010, 33(9): 1871–6.
- [46] Khan MI, Ahmad M, Khan RA, Ullah A, Rehman S, Ullah B. Phytotoxic, antioxidant and antifungal activity of crude methanolic extract of *Equisetum debile*. *Int J Biosci.* 2013; 3: 130–5.
- [47] Rehman T, Shad MA, Nawaz H, Andaleeb H, Aslam M. Biochemical, phytochemical and antioxidant composition of *Equisetum debile* Roxb. *Biochemist & Analytical Biochemist.* 2018; 7(4):1–6.