



International Journal of Pharmaceutical and Phytopharmacological Research (eIJPPR)

[Impact Factor – 0.852]

Journal Homepage: www.eijppr.com

Research Article

Article ID: 404

Evaluation of *in vitro* anti-diabetic activity of *Elaeocarpus serratus* fruit

D. H. Geetha*, Indhiramuthu Jayashree and M. Rajeswari

PG and Research Department of Botany, Vellalar College for Women, (Autonomous), Erode- 638012, Tamil Nadu, India

*Corresponding Author: Email - geethadhandapani2010@gmail.com

Article info

Article History:
Received 28 August 2015
Accepted 23 September 2015

Keywords:

Elaeocarpus serratus, Anti-diabetic activity, α -amylase, α -glucosidase, acarbose

Abstract

Diabetes is a clinical syndrome characterized by hyperglycemia due to absolute or relative deficiency of insulin. Recent decades have experienced a sharp increase in the incidence and prevalence of Diabetes mellitus. Pancreatic α -amylase and α -glucosidase inhibitors offer an effective strategy to lower levels of post prandial hyperglycemia via control of starch breakdown. The objective of the present work was to evaluate the ethanolic extract of fruit of *Elaeocarpus serratus* by using the *in vitro* anti-diabetic activity. The results suggested that the maximum α -amylase inhibitory activity and moderate α -glucosidase inhibitory activity. Acarbose was used as a standard drug.

1. INTRODUCTION

Diabetes mellitus is a chronic disease caused by inherited or acquired deficiency in insulin secretion and by decreased responsiveness of the organs to secreted insulin. Such a deficiency results in increased blood glucose level, which in turn can damage many of the body's systems, including blood vessels and nerves¹. For its therapy, along with the synthetic drugs, many agents of the plant origin are also in use particularly for the treatment of non-insulin dependent diabetes mellitus. According to world ethno botanical information reports, almost 800 plants may possess anti-diabetic potential². Medicinal plants with anti-diabetic potentials decrease the absorption of glucose by inhibiting the carbohydrate hydrolyzing enzymes, such as pancreatic amylase^{3,4}. In the past decade, research has been focused on scientific evaluation of traditional drugs of plant origin and screening of more effective and safe hypoglycemic agents has continued to be an important area.

Natural products, such as plants extract, either as pure compounds or as standardized extracts, provide unlimited opportunities for new drug discoveries because of the unmatched availability of chemical diversity⁵. *Elaeocarpus* is a genus of tropical and subtropical evergreen trees and shrubs belonging to family Elaeocarpaceae. Studies indicate that various species contain chemical constituent such as triterpenes, tannins, indolizidine alkaloids, flavonoids, and ellagic acid derivatives⁶⁻⁷. Several species have been known to possess anti-inflammatory⁸, antimicrobial⁹, analgesic¹⁰, antioxidant activity¹¹ and antihypertensive¹² activities. *Elaeocarpus serratus* leaves are used in the treatment of rheumatism¹³, diuretic, cardiovascular stimulant and as antidote to poison, antimicrobial activity¹⁴ while the fruits are locally prescribed for the treatment of diarrhea and dysentery. The fruit juice is given for stimulating secretions from taste buds thus increasing appetite in patients¹⁵. Leaves contain Myricitrin, Myricetin, Mearnsitrin and Ellagic acid. Fruits contain tannin and large amount of plant acids¹⁶⁻¹⁷. In continuation of the above studies, the present work is aimed to evaluate the anti-diabetic activity of ethanolic fruit extract using *in vitro* assay system.

2. MATERIALS AND METHODS

2.1 Plant Collection and Extraction

The fruit of *Elaeocarpus serratus* L. were collected from Upper Palani Hills of Western Ghats (Kodaikanal Forest Division), India and were authenticated at Botanical Survey of India (BSI), Southern Circle, Coimbatore, India and the herbarium of Voucher specimen number BSI/SRC/5/23/2011-12/Tech. 454 has been deposited at the PG and Research Department of Botany, Vellalar College for Women, Erode

(T.N), India. Coarse powder from the shade dried fruit of *E. serratus* (500g) were exhaustively extracted using soxhlet apparatus with absolute ethanol (78.5°C). The extract were dried (free of solvent) using a vacuum evaporator. The extract thus obtained was stored in refrigerator and used for the experiment.

2.2 Chemicals

The chemicals used in the present work were acquired from E. Merck (INDIA) Limited, Hi-media, Pvt. Ltd, India and Sigma-Aldrich, India.

2.2.1 In Vitro α -amylase Inhibition Assay

The α -amylase inhibitory activity was determined according to the method described by¹⁸. 20-100 μ g/ml of plant extract and 1ml enzyme solution were mixed in a tube and incubated at 25°C for 30 minutes. To 1ml of this mixture was added 1ml of 0.5% starch solution and the tube was incubated at 25°C for 3 minutes. Then, 1ml of the color reagent Dinitro Salicylic acid (DNSA) was added and the closed tube was placed in a water bath at 85°C. After 15minutes, the reaction mixture was removed from the water bath and cooled. Thereafter, it was diluted with 9ml distilled water and the absorbance value determined at 540nm. Acarbose was used as positive control. The percentage of α -amylase inhibition was assessed using the formula:-

$$\text{Percentage of } \alpha\text{-amylase inhibition} = \frac{A_{540} \text{ control} - A_{540} \text{ test}}{A_{540} \text{ control}} \times 100$$

2.2.2 In Vitro α -glucosidase Inhibition Assay

The assay was carried out following the standard protocol¹⁹. α -glucosidase inhibitory activity was determined by premixing α -glucosidase (0.07units) with 20-100 μ g /ml of plant extract. Then 3mM p-nitrophenyl glucopyranoside was added as a substrate. This reaction mixture was incubated at 37°C for 30minutes and the reaction was terminated by addition of 2ml of sodium carbonate. The α -glucosidase activity was determined by measuring the p-nitrophenyl release from p-nitrophenyl glucopyranoside at 400nm. Acarbose was used as positive control. The percentage of α -amylase inhibition was assessed using the formula:

$$\text{Percentage of } \alpha\text{-glucosidase inhibition} = 1 - B/A \times 100$$

A - Absorbance of control; B - Absorbance of sample containing extracts.

3. RESULTS

The results showed that ethanolic extract of fruit of *Elaeocarpus serratus* exhibited different degree of α -amylase and α -glucosidase inhibitory activities in *in vitro* assays using starch and p-nitrophenyl glucopyranoside as substrate, respectively. The percentage inhibition at 20, 40, 60, 80 and 100 μ g/ml concentration of plant extract showed a dose-dependent reduction in percentage inhibition. Maximum inhibition of 89.54 \pm 2.99% was shown by 100 μ g/ml concentration of the fruit extract. The standard drug acarbose showed 94.65 \pm 0.82% at 100 μ g/ml (Figure 1). The plant extract displayed moderate α -glucosidase inhibitory activity (6.68 \pm 0.59% to 25.93 \pm 0.34%). The standard acarbose (at concentrations 20-100 μ g/ml) showed α -glucosidase inhibitory activity from 43.12 \pm 0.59% to 75.80 \pm 1.01%. According to the experimental results, it was certainly confirmed that the plant extract exhibited weak α -glucosidase enzyme inhibition when compared to α -amylase enzyme inhibition (Figure 2).




Figure 1: Effect of ethanolic extract of fruit of *E. serratus* on α -amylase inhibition

Values are mean of three independent analyses of the extract \pm standard deviation (n=3)
Standard - Acarbose (20-100 μ g/ml), EESF - Ethanolic extract of *E. serratus* fruit (20-100 μ g/ml)




Figure 2: Effect of ethanolic extract of fruit of *E. serratus* on α -glucosidase inhibition

Values are mean of three independent analyses of the extract \pm standard deviation ($n=3$)
Standard - Acarbose (20-100µg/ml), EESF - Ethanol extract of *E. serratus* fruit (20-100 µg/ml)

4. DISCUSSION

Diabetes is a debilitating disease affecting millions of people worldwide. Since the disease has no known modern allopathic cure, it requires lifelong health. One therapeutic approach is the prevention of carbohydrate absorption after food intake, which is facilitated by inhibition of enteric enzymes including α -amylase and α -glucosidase present in the brush borders of intestine. A major goal in the treatment of diabetes mellitus is to maintain near normal blood glucose levels in both the fasting and postprandial state²⁰. Furthermore, some inhibitors of α -amylase and α -glucosidase such as phaseolamin, acarbose and voglibose are currently used to suppress postprandial glucose levels in diabetic patients²¹. Hence, plants have been suggested as a rich, as yet unexplored source of potentially useful anti-diabetic drugs. However, only a few have been subjected to detailed scientific investigation due to a lack of mechanism-based available *in vitro* assays^{22,23}.

The present investigation indicated that the plant extract exhibited weak α -glucosidase enzyme inhibition when compared to α -amylase. In earlier studies, the hypoglycemic potential of *Carpesium abrotanoides* was evaluated by the α -amylase and α -glucosidase inhibition assay. The optimal concentration of *C. abrotanoides* required for the 50% inhibition (IC_{50}) against α -glucosidase was 44.22µg/ml. Acarbose was used as positive control with IC_{50} value of 2.5µg/ml²⁴. According to²⁵ the acetone extract of both fruits (68.9%) and leaves (89.6%) of *Terminalia bellirica* showed strong inhibitory activity against α -amylase whereas the aqueous extract of fruit (6.89%) and leaf (7.5%) was found to exhibit highest α -glucosidase activity. Similar trend was noted by^{26,27} who showed that the methanolic extracts of *Amaranthus cruentus* and *Moringa oleifera* showed promising levels of α -amylase (10-45%) and α -glucosidase (13-80%) inhibition activities. According to²⁸ the methanolic extract of leaves of *Psidium guava* efficiently inhibited both α -amylase (89.4%) and α -glucosidase (96.3%) enzymes *in vitro* in a dose-dependent manner. Our experiment was in accordance with the results reported by²⁹⁻³¹.

5. CONCLUSION

We, for the first time, have demonstrated the ability of ethanolic fruit extract of *Elaeocarpus serratus* to inhibit glucose diffusion using an *in vitro* model of glucose absorption. Further studies are required to elucidate whether *in vitro* effects represented therapeutic potential by limiting postprandial glucose absorption and for improving glycemic control in type 2 diabetic subjects.

REFERENCES

- Matsui T, Tanaka T, Tamura S, Toshima A, Miyata Y, Tanaka K. Alpha-glucosidase inhibitory profile of Catechins and Theaflavins. J.Agric. Food Chem, 2007, 55:99-105.
- Alarcon-Aguilara FJ, Roman-Ramos R, Perez-Gutierrez S, Aguilar-Contreras A, Contreras-Weber CC, Flores-Saenz JL. Study of the anti-hyperglycemic effect of plants used as anti-diabetics. J. Ethnopharmacol, 1998, 61:101-110.
- Tripathi KD. Essential of medical pharmacology, 5th Edition. 2003, 235.
- Gulati V, Ian HH, Palombo EA. Enzyme inhibitory and antioxidant activities of traditional medicinal plants: Potential application in the management of hyperglycemia. Complementary and Alternative Medicine, 2012, 12 (77):1-9.
- Bhaduri SK, Chanda S, Majumdar P. Bioresource Technology, 1998, 63: 279-281.
- Fang X, Phoebe Jr CH, Pezzuto JM, Fong HH, Farnsworth NR, Yellin B, Hecht SM. Plant anticancer agents, XXXIV. Cucurbitacins from *Elaeocarpus dolichostylus*. J.Nat.Prod, 1984, 47 (6): 988-993.

7. Ito A, Chai HB, Lee D, Kardono LBS, Riswan S, Farnsworth NR, Cordell GA, Pezzuto JM, Kinghorn AD. Ellagic acid derivatives and cytotoxic cucurbitacins from *Elaeocarpus mastersii*. *Phytochemistry*, 2002, 61 (2):171-174.
8. Singh RK, Bhattacharya SK, Acharya SB. Studies on extracts of *Elaeocarpus sphaericus* fruits on *in vitro* rat mast cells. *Phytomedicine*, 2000, 7(3): 205-207.
9. Singh R and Nath G. Antimicrobial Activity of *Elaeocarpus sphaericus*. *Phytother. Res*, 1999, 13: 448-450.
10. Bhattacharya SK, Debnath PK, Pandey VB. Pharmacological investigations on *Elaeocarpus ganitrus*. *Planta Medica*, 1975, 28: 174-177.
11. Sathish Kumar T, Shanmugam S, Palvannan T, Bharathi Kumar VM. Evaluation of antioxidant properties of *Elaeocarpus ganitrus* Roxb. Leaves. *Iranian Journal of Pharmaceutical Research*, 2008, 7(3): 211-215.
12. Pandey VB and Bhattacharya SK. Scientific appraisal of rudraksha (*Elaeocarpus ganitrus*): chemical and pharmacological studies. *J. Res. Edu. Ind. Med*, 1985, 4: 47-50.
13. Geetha DH, Indhiramuthu Jayashree, Rajeswari M. *In vitro* anti-arthritic activity of *Elaeocarpus serratus* Linn. (Elaeocarpaceae). *IJPSR*, 2015, 6(6): 2649-2651.
14. Jayashree I, Geetha DH, Rajeswari M. Evaluation of antimicrobial potential of *Elaeocarpus serratus* L. *Int J Pharm Sci Res*, 2014, 5(8): 3467-3472.
15. Ghani A. Monographs: *Elaeocarpus serratus* Linn. In: Medicinal plants of Bangladesh: Chemical constituents and uses. Asiatic. Soc. Bangla. Publ, 1998, 167-168.
16. Chopra RN, Chopra IC, Hunda KI, Kumar LD. Chopra's indigenous drug of India. India: Academic Publishers. 1982.
17. Most NP, Sarwar S, Chowdhury SA, Zakaria HM. and Huda NH. *In vitro* cytotoxicity and antioxidant studies of *Elaeocarpus serratus*. *Stamford J.Pharm.Sci*, 2009, 2(2):86-90.
18. Giancarlo S, Rosa LM, Nadjafi F, Francesco M. Hypoglycaemic activity of two species extracts: *Rhus coriaria* L. and *Bunium persicum* Boiss. *Nat. Prod. Res*, 2006, 20:882-886.
19. Miller GL. Use of dinitro salicylic acid reagent for determination of reducing sugar. *Anal. Chem*, 1959, 31:426-428.
20. Bailey CJ. Potential new treatments for type 2 diabetes. *Trends in pharmacol sci*, 2000, 21: 259-265.
21. Sanap SP, Ghosh S, Jabgunde AM, Pinjari RV, Gejji SP, Singh S, Chopade BA, Dhavale DD. Synthesis, computational study and glycosidase inhibitory activity of polyhydroxylated conidine alkaloids- a bicyclic iminosugar. *Org Biomol Chem*, 2010, 8: 3307-3315.
22. Saxena A. and Vikram NK. Role of selected Indian plants in management of type 2 diabetes. *J. Altern. Comp. Med*, 2004, 10: 369-378.
23. Ahamad J, Naquvi JK, Showkat R, Ali MM, Shuaib M. "Review on role of natural alpha-glucosidase inhibitors for management of diabetes mellitus", *IJBR*, 2011, 2(6):374-380.
24. Mayur B, Sandesh S, Shruti S, Yum SS. "Antioxidant and α -glucosidase inhibitory properties of *Carpesium abrotanoides* L". *Journal of Medicinal Plants Research*, 2010, 4(15):1547-1553.
25. Gayathri Devi S, Manila TN, Mary Shoba Das C. *In vitro* anti-diabetic properties of the fruits and leaves of *Terminalia bellirica*. *Journal of Natural Sciences Research*, 2012, 2(9):54-62.
26. Catherine KH, Pandey P, Arya CM, Ahmed Z. "Evaluation of anti-diabetic activity of two important memory enhancing medicinal plants *Baccopa monnieri* and *Centella asiatica*", *Indian. J. Pharmacol*, 2010, 44(1): 114-117.
27. Rani BA, Pugin A, Wendehenne D. "New insights into nitric oxide signaling in plants", *Annual Review of Plant Biology*, 2011, 59: 21-39.
28. Manikandan R, Vijaya Anand A, Durai Muthamani G. Phytochemical and *in vitro* anti-diabetic activity of methanolic extract of *Psidium guajava* leaves. *Int. J. Curr. Microbiol. App. Sci*, 2013, 2(2): 15-19.
29. Deguchi Y, Osada K, Uchida K, Kimura H, Yoshikawa M, Kudo T, Yasui H, Watanuki M. Effects of extract of guava leaves on the development of diabetes in the db/db mouse and on the postprandial blood glucose of human. *Nippon geikagaku Kaishi (in Japanese)*, 1998, 72: 923-931.
30. Kavitha S, Kamaraj M, Rajeshwari S. *In vitro* alpha amylase and alpha glucosidase inhibition activity of crude ethanol extract of *Cissus arnottiana*. *Asian Journal of Plant Science and Research*, 2012, 2 (4):550-553.
31. Kavimani S, Saminathan K, Senthil Kumar R. *In vitro* anti-diabetic activity of *Dolichandrone atrovirens* - An Indian medicinal plant. *Int. J. of Pharmacotherapy*, 2014, 4(3): 107-113.