



Internal Mammary Lymph Node Volume: What Radiation Dose Is Received with Modified Wide Tangents?

Reem Ujaimi, MD, FRCPC^{1*}, Nisreen Awad², MD, Zakaria Hasan³, Noura Abbas⁴,
Wejdan Baageel⁴, Salma Khayyat⁴, Mohammed Attar¹ MD, Shyma M Al-Khateeb PhD,
MP⁴

¹ Faculty of Medicine - King Abdulaziz University, Radiotherapy section- Department of Radiology, Jeddah, Saudi Arabia.

² King Abdulaziz University Hospital, Radiotherapy section- Department of Radiology, Jeddah, Saudi Arabia.

³ King Faisal Specialist Hospital and Research Center, Biomedical Physics, Jeddah, Saudi Arabia.

⁴ Faculty of applied medical Sciences - King Abdulaziz University, Diagnostic radiography technology, Jeddah, Saudi Arabia.

ABSTRACT

Introduction: There is a growing interest in radiating the internal mammary (IM) nodes in the context of adjuvant radiotherapy in breast cancer. There have been variations in defining its CTV and determining acceptable coverage. Clear definitions are necessary; as such, this paper sought to estimate the dose delivered to an institutionally defined IM CTV using the standard modified wide tangential (MWT) fields. **Materials and Methods:** IM vessels in the first three intercostal spaces of 10 randomly selected patients were contoured. Then, 5 mm was added and trimmed from anatomical boundaries to create the CTV. Three planning target volumes (PTVs) were created (PTV1, PTV2, and PTV3). Plans were created using a 3D technique with the aim of covering the target volumes by 90%–95% of the dose while keeping the V20 of the ipsilateral lung <35% and the V25 of the heart <10%, with a minimum mean heart dose. Two MWT covered the IM lymph nodes and breast or chest wall; one anterior field was used for the supraclavicular lymph nodes. A dose of 50 Gy/25 fractions was prescribed at the isocenter. DVH curves for normal tissues were generated. **Results:** Six left- and four right-sided patients were included. Nine and one patients underwent mastectomy and lumpectomy, respectively. The mean percentage of the volume that received 95% (V95) \pm standard deviation (SD) was: IM vessel (V95) = 85.49 \pm 14.06; CTV (V95) = 78.73 \pm 14.30; PTV1 (V95) = 65.88 \pm 11.99; PTV2 (V95) = 76.05 \pm 12.31; PTV3 (V95) = 70 \pm 12.84. The mean heart V25 was 5.5 \pm 3.4 for left-sided patients. The mean heart dose was 4.5 \pm 1.8 for left-sided patients. The mean ipsilateral lung V20 was 35.15 \pm 0.39 and 34.31 \pm 1.17 for right- and left-sided patients, respectively. **Conclusions:** In conclusion, we found that using our CTV volume, covering 80-85% of the CTV by 95-90% of the dose, respectively, is considered acceptable. We should aim for better coverage of our target volumes but proceed with caution to maintain and improve tumor control without exceeding normal tissue toxicity.

Key Words: Cancer, internal mammary, MVT, breast.

eIJPPR 2020; 10(5):76-83

HOW TO CITE THIS ARTICLE: Reem Ujaimi, Nisreen Awad, Zakaria Hasan, Noura Abbas, Wejdan Baageel, Salma Khayyat and *et al.* (2020). "Internal mammary lymph node volume: What radiation dose is received with modified wide tangents?", International Journal of Pharmaceutical and Phytopharmacological Research, 10(5), pp.76-83.

INTRODUCTION

Breast cancer is the most common cancer affecting women in Saudi Arabia and around the world. [1-4] A number of studies have shown that adjuvant breast or chest wall and regional nodal radiation improves local control and overall survival, especially in high-risk breast cancer patients. [5-

11] As a result, adjuvant radiotherapy has been an integral part in the management of breast cancer.

While these studies included the internal mammary (IM) chain of lymphatics in the target, many radiation oncologists do not routinely include the IM when treating the regional nodes, [12, 13] probably due to a fear of lung

Corresponding author: Reem Ujaimi

Address: Assistant Professor Radiation Oncology, King Abdul-Aziz University, Jeddah, Saudi Arabia.

E-mail: ✉ rkujaimi@kau.edu.sa

Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 20 July 2020; **Revised:** 07 October 2020; **Accepted:** 10 October 2020



and cardiac toxicity and the lack of strong evidence to support this practice. Irradiating the IM has recently become a matter of interest, especially after studies related survival advantage to IM radiation, [9, 14 15] and given improvements in treatment planning techniques that allow radiation to reach the IM without exceeding doses to the lung and heart. [16-19] Including the regional lymph nodes in treatment planning should be carried out with the least possible dose to the organs at risk. The aim of this research is to evaluate the coverage of the IM clinical target volume (CTV) and different planning target volumes (PTV), assuming that modified wide tangential fields are the optimal standard.

METHODS AND MATERIALS

The unit of biomedical ethics at our institution approved the study. In order to reduce selection bias, 10 unilateral breast cancer patients were randomly selected with the aim of recruiting a comparable number of right- and left-sided patients. All patients were planned using the modified wide tangent technique, as described in the NCIC CTG MA.20 protocol. [7] The simulation process was done for all patients using computed tomography (CT) scan with slices 2.5–3 mm in thickness. All patients were in the supine position on a breast board with both arms raised above their head. IM vessels in the first three intercostal spaces were contoured following the Radiation Therapy Oncology Group (RTOG) guidelines by three students and reviewed by two radiation oncologists. Subsequently, a 5-mm margin was added for the CTV IM and trimmed from the muscles, bones, lungs, and heart. Another 5-mm margin was added all-around to create PTV1. Afterward, PTV1 was trimmed from the lung and heart to create PTV2 (PTV-eval). PTV3 was a 5-mm margin added to the IM vessels (5 mm around the RTOG contours) (Figure 1). The heart and lungs were contoured using the RTOG consensus guidelines. A plan optimization target volume was created by adding 1 cm around the vessels following the MA 20 protocol for CT-based planning. [7] Apart from the latter volume, the planner was blinded to all other volumes. The prescribed dose for all patients was 50 Gy/25 fractions to the isocenter. Treatment planning was forward using 3–4 fields with the aim of covering the target volume by 90% or 95% of the prescribed dose while minimizing the dose to the heart and lungs to maintain the V20 of the lung below 35% and the V25 of the heart below 10%. Two tangential fields were used to cover the IM lymph nodes, breast or chest wall, and one anterior field, with or without a posterior field for the supraclavicular lymph nodes. Monaco® 5.11 from Elekta, using Monte Carlo treatment planning algorithm was used for planning. Dose–volume histogram (DVH) curves for normal tissues were generated. The information from each treatment plan was used for data analysis.

RESULTS

The prescribed dose was 50 Gy/25 fractions in all cases. Four patients had right-sided breast cancer and six were left-sided. Nine patients underwent mastectomy and one patient underwent lumpectomy. The mean percentage for volumes that received 95% of the prescribed dose (V95) ± standard deviation (SD) were as follows: IM vessel (V95) = 85.49±14.06; CTV (V95) = 78.73±14.29; PTV1 (V95) = 65.88±11.99; PTV2 (V95) = 76.05±12.31; and PTV3 (V95) = 70.00±12.84. Table 1 shows the V95 of all 10 patients. Figure 2 shows an example of the isodose lines.

Table 1: V95

No.	Laterality	V95 IM PTV1 (%)	V95 IM PTV2 (%)	V95 IM PTV3 (%)	V95 IM CTV (%)	V95 IM Vessels (%)
1	LT	60.93	70.72	63.76	68.67	77.94
2	LT	54.24	57.07	62.71	62.02	86.25
3	LT	78.54	87.61	82.87	90.66	92.69
4	LT	68.42	76.63	71.28	83.19	78.57
5	LT	76.77	88.08	78.88	95.6	97.42
6	LT	77.9	82	81.7	85.8	80.6
7	RT	59.83	65	70.52	76.94	98.92
8	RT	43.57	59.38	41.72	50.72	52.44
9	RT	60.76	83.67	63.63	82.30	94.54
10	RT	77.83	90.31	82.92	91.37	95.53
	Mean	65.88	76.05	70.00	78.73	85.49
	SD	11.988	12.31	12.84	14.29	14.06

The mean percentage for volumes that received 90% of the dose (V90) were as follows: IM vessel (V 90) = 91.87±10.87; CTV (V90) = 86.04±12.79; PTV1 (V90) = 75.34±11.55; PTV2 (V90) = 81.25±11.61; and PTV3 (V90) = 81.31±11.7. The mean V90% for all patients is presented in Table 2.

Table 2: V90

No.	Laterality	V90 IM PTV1 (%)	V90 IM PTV2 (%)	V90 IM PTV3 (%)	V90 IM CTV (%)	V90 IM Vessels (%)
1	LT	69.66	74.65	74.9	72.45	81.61
2	LT	62.8	63.36	76.01	72.43	94.26
3	LT	88.54	93.70	93.38	98.19	100
4	LT	78.1	83.35	83.37	88.93	86.18
5	LT	84.85	92.14	89.38	98.79	99.73
6	LT	85.3	86.9	88.6	90.3	89.3
7	RT	69.99	70.48	83.38	87.29	100

8	RT	54.14	66.25	53.89	61.68	67.63
9	RT	72.33	88.05	77.72	93.77	100
10	RT	87.71	93.58	92.49	96.55	99.95
	Mean	75.34	81.24	81.31	86.04	91.87
	SD	11.55	11.61	11.70	12.79	10.87

The mean heart (V25) dose was 0.037 ± 0.035 SD for right-sided patients and 5.51 ± 3.4 SD for left-sided patients. The mean heart dose was 1.2 ± 0.5 for right-sided patients and 4.5 ± 1.8 for left-sided patients. The mean lung (V20) dose was 35.15 ± 0.39 SD for right-sided patients and 34.31 ± 1.17 SD for left-sided patients. Table 3 shows the heart and lung doses for left-sided patients.

Table 3: Left-sided lung and heart constraints

No.	Lung V20 (%)	Heart V25 (%)
1	34.78	3.14
2	34.53	10.01
3	34.49	7.72
4	32.09	2.10
5	35.6	7.75
6	34.4	2.3
Mean	34.31	5.50
SD	1.17	3.40

DISCUSSION

Adjuvant, locoregional radiotherapy is an integral part of breast cancer management [20]. The extent of surgical resection, particularly in the axilla, and the role of radiotherapy is continuously evolving [21]. The inclusion of the IM chain of lymph nodes has represented an ongoing controversy. On the one hand, most major trials on which we base our recommendations for adjuvant radiation include radiation to the IM. [6, 7, 11] On the other hand, an isolated recurrence rate in the IM is low, [22] and the potential toxicities associated with increasing doses to the heart and lung, which is inevitable with IM radiation, is concerning. [23, 24] Recently, however, there has been a renewed interest in IM radiation, partly due to the emerging evidence that linked it to improvements in disease-free survival. [9, 14] Moreover, advances in radiotherapy techniques, including the use of deep inspiration breath hold (DIBH), have made it possible to treat the IM area with relatively small doses to the heart. [25]

Locoregional breast radiotherapy has largely continued to be field-based. Nonetheless, volumetric planning is essential, particularly with the use of intensity-modulated radiotherapy. As such, developing a standard contouring guideline and coverage criteria is important. The consensus reached by the RTOG group is to contour the IM/thoracic vessels in the first three intercostal spaces, starting at the cranial border of the medial first rib. [26] Using these

contours, a standard monoisocentric 2 tangential field, an anterior electron field, and a supraclavicular field, 80% of the target would be covered by 90% of the dose. With contour-based planning, this will go up to 85%. [26] In other words, a conventional, anterior electron-based field does not cover the RTOG-defined contours. Moreover, it is questionable as to whether these contours represent the area at risk of recurrence. Jethwa et al. have found that among 130 IM node metastases, 78% were in the first three intercostal spaces, and of those, 53% were covered by RTOG-defined contours. By adding 4 mm, 5 mm, and 6 mm medial and lateral to the vessels, 90%, 93%, and 95% of the metastatic nodes would have been encompassed, respectively. [27]

The authors also found that 8% and 14% of the recurrences were located cranial to the first rib, but caudal to the confluence of the IM vessels and the brachiocephalic vessels, and caudal to the third intercostal space, respectively. The ESTRO group began contouring the IM CTV 5 mm from the subclavian vein and down to the cranial aspect of the fourth rib. They added 5 mm medial and lateral to the vessels. [28] Further, the Danish group also contoured 5 mm around the vessels but included the four first intercostal spaces. [29] Dijkima et al., from the Netherlands, increased the contours to reach the jugulosubclavian junction. [30] To our knowledge, it is not clear what percentage of those contours are covered with a modified wide tangent technique.

The CTV contours described in the methods section of this study adopted the approach agreed upon by the involved radiation oncologist after reviewing all the above evidence. The addition of a PTV margin was thought to be crucial in the era of conformal radiotherapy. We chose to use different PTVs to account for variations in the PTV margins among different institutions and radiation oncologists.

Our results confirmed that the standard modified wide tangential technique does not adequately cover our target volumes. The RTOG target V95 and V90 was covered by 85.5% and 91.8%, respectively, which was comparable to the contour-based V90 of 85% achieved by Fontanilla et al. [26] Our CTV was covered by 78.7%. It is reasonable to assume that if contour-based planning was used, the results would have been better. In fact, in real life, we do follow contour-based planning to allow for better target coverage and maintain organ-at-risk (OAR) constraints. To cover our PTV volumes, advanced treatment techniques with IMRT and DIBH would have to be used, particularly in the setting of positive IM lymph nodes, where one would consider extending the coverage to include the space cranial and caudal to the first and fourth rib, respectively. Assuming the modified wide tangential technique is the standard approach for elective IM radiation, one would accept 85% coverage to the vessels and 80% to our CTV; however, one

should aim for better coverage with advanced techniques, particularly in the context of positive IM nodes.

CONCLUSIONS

In conclusion, we found that using our CTV volume, covering 80-85% of the CTV by 95-90% of the dose, respectively, is considered acceptable. We should aim for better coverage of our target volumes but proceed with caution to maintain and improve tumor control without exceeding normal tissue toxicity.

Conflict of Interest

No conflict of interest to be disclosed by any of the authors.


Funding:

No funding sources associated with this research.


REFERENCES

- [1] McGuire S. World Cancer Report 2014. Geneva, Switzerland: World Health Organization, International Agency for Research on Cancer, WHO Press, 2015, 2016.
- [2] Saggi S, Rehman H, Abbas ZK, Ansari AA. Recent incidence and descriptive epidemiological survey of breast cancer in Saudi Arabia. *Saudi Med J* 2015;36(10):1176-1180.
- [3] Saher A, Heba A, Hanan A. Knowledge, Attitude and Practices of Working Women in Tabuk University Regarding Breast Cancer Int. *J. Pharm. Res. Allied Sci.* 2018;7(3):198-208.
- [4] Samir D, Naouel A, Safa G. Assessment of Hematological Parameters, Enzymes Activities, and Oxidative Stress Markers in Salivary and Blood of Algerian Breast Cancer Patients Receiving Chemotherapy. *J. Biochem. Technol.* 2019;10(4):50-8.
- [5] McGale P, Taylor C, Correa C, Cutter D, Duane F, Ewertz M, Gray R, Mannu G, Peto R, Whelan T, Wang Y. Effect of radiotherapy after mastectomy and axillary surgery on 10-year recurrence and 20-year breast cancer mortality: meta-analysis of individual patient data for 8135 women in 22 randomised trials. *Lancet* 2014;383(9935):2127-2135.
- [6] Ragaz J, Olivotto IA, Spinelli JJ, Phillips N, Jackson SM, Wilson KS, Knowling MA, Coppin CM, Weir L, Gelmon K, Le N. Locoregional radiation therapy in patients with high-risk breast cancer receiving adjuvant chemotherapy: 20-year results of the British Columbia randomized trial. *J Natl Cancer Inst* 2005;97(2):116-126.
- [7] Whelan TJ, Olivotto IA, Parulekar WR, Ackerman I, Chua BH, Nabid A, Vallis KA, White JR, Rousseau P, Fortin A, Pierce LJ. Regional nodal irradiation in early-stage breast cancer. *N Engl J Med* 2015;373(4):307-316.
- [8] Whelan TJ, Olivotto IA, Levine MN. Regional nodal irradiation in early-stage breast cancer. *N Engl J Med* 2015;373(19):1878-1879.
- [9] Poortmans PM, Collette S, Kirkove C, Van Limbergen E, Budach V, Struikmans H, Collette L, Fourquet A, Maingon P, Valli M, De Winter K. Internal mammary and medial supraclavicular irradiation in breast cancer. *N Engl J Med* 2015;373(4):317-327.
- [10] Ragaz J, Jackson SM, Le N, Plenderleith IH, Spinelli JJ, Basco VE, Wilson KS, Knowling MA, Coppin CM, Paradis M, Coldman AJ. Adjuvant radiotherapy and chemotherapy in node-positive premenopausal women with breast cancer. *N Engl J Med* 1997;337(14):956-962.
- [11] Overgaard M, Hansen PS, Overgaard J, Rose C, Andersson M, Bach F, Kjaer M, Gadeberg CC, Mouridsen HT, Jensen MB, Zedeler K. Postoperative radiotherapy in high-risk premenopausal women with breast cancer who receive adjuvant chemotherapy. Danish Breast Cancer Cooperative Group 82b Trial. *N Engl J Med* 1997;337(14):949-955.
- [12] Belkacemi Y, Kaidar-Person O, Poortmans P, Ozsahin M, Valli MC, Russell N, Kunkler I, Hermans J, Kuten A, van Tienhoven G, Westenberg H. Patterns of practice of regional nodal irradiation in breast cancer: results of the European Organization for Research and Treatment of Cancer (EORTC) Nodal Radiotherapy (NORA) survey. *Ann Oncol* 2015;26(3):529-535.
- [13] Park HJ, Oh DH, Shin KH, Kim JH, Choi DH, Park W, Suh CO, Kim YB, Ahn SD, Kim SS. Patterns of practice in radiotherapy for breast cancer in Korea. *J Breast Cancer* 2018;21(3):244-250.
- [14] Thorsen LB, Offersen BV, Danø H, Berg M, Jensen I, Pedersen AN, Zimmermann SJ, Brodersen HJ, Overgaard M, Overgaard J. DBCG-IMN: A population-based cohort study on the effect of internal mammary node irradiation in early node-positive breast cancer. *J Clin Oncol* 2016;34(4):314-320.
- [15] Thorsen LB, Thomsen MS, Berg M, Jensen I, Josipovic M, Overgaard M, Overgaard J, Skogholt P, Offersen BV, Danish Breast Cancer Cooperative Group Radiotherapy committee. CT-planned internal mammary node radiotherapy in the DBCG-IMN study: Benefit versus potentially harmful effects. *Acta Oncol* 2014;53(8):1027-1034.
- [16] Bergom C, Currey A, Desai N, Tai A, Strauss JB. Deep inspiration breath hold: Techniques and advantages for cardiac sparing during breast cancer irradiation. *Front Oncol* 2018;8:87.
- [17] Lemanski C, Thariat J, Ampil FL, Bose S, Vock J, Davis R, Chi A, Dutta S, Woods W, Desai A, Godinez J. Image-guided radiotherapy for cardiac sparing in


- patients with left-sided breast cancer. *Front Oncol* 2014;4:257.
- [18] Mast ME, van Kempen-Harteveld L, Heijenbrok MW, Kalidien Y, Rozema H, Jansen WP, Petoukhova AL, Struikmans H. Left-sided breast cancer radiotherapy with and without breath-hold: does IMRT reduce the cardiac dose even further? *Radiother Oncol* 2013;108(2):248-253.
- [19] Coon AB, Dickler A, Kirk MC, Liao Y, Shah AP, Strauss JB, Chen S, Turian J, Griem KL. Tomotherapy and multifield intensity-modulated radiotherapy planning reduce cardiac doses in left-sided breast cancer patients with unfavorable cardiac anatomy. *Int J Radiat Oncol, Biol, Phys* 2010;78(1):104-110.
- [20] Hosseini SH, Rafiei A, Janbabai G, Tirgari A, Zakavi A, Yazdani J, Bolhari J, Golzari M, Vaezzadeh N, Douki ZE. Comparison of Religious Cognitive Behavioral Therapy (RCBT), Cognitive Behavioral Therapy (CBT) And Citalopram On Depression and Anxiety Among Women with Breast Cancer: A Randomized Controlled Trial. *Pharmacophores*. 2018 May 1;9(3):37-47.
- [21] Mousavizadeh SM, Mahmoudvand H, Rezaadeh H, Otaghvar HR, Mousavizadeh SM, Rouietan A. Using CAD/CAM Technology to Prepare Skin Incision Template and Predict Size Reduction of Each Breast Separately and Final Shape of Breast After Reduction Mammoplasty Surgery. *Entomol. appl. sci. lett.* 2018 Jun 23;5(1):51-8.
- [22] Chen L, Gu Y, Leaw S, Wang Z, Wang P, Hu X, Chen J, Lu J, Shao Z. Internal mammary lymph node recurrence: Rare but characteristic metastasis site in breast cancer. *BMC Cancer* 2010;10(1):479.
- [23] Darby SC, Ewertz M, McGale P, Bennet AM, Blom-Goldman U, Brønnum D, Correa C, Cutter D, Gagliardi G, Gigante B, Jensen MB. Risk of ischemic heart disease in women after radiotherapy for breast cancer. *N Engl J Med* 2013;368(11):987-998.
- [24] Choi JY, Cho EY, Choi YJ, Lee JH, Jung SP, Cho KR, Kim CY, Kim YH, Park KH. Incidence and risk factors for congestive heart failure in patients with early breast cancer who received anthracycline and/or trastuzumab: A big data analysis of the Korean Health Insurance Review and Assessment service database. *Breast Cancer Research and Treatment* 2018;171(1):181-188.
- [25] Yeung R, Conroy L, Long K, Walrath D, Li H, Smith W, Hudson A, Phan T. Cardiac dose reduction with deep inspiration breath hold for left-sided breast cancer radiotherapy patients with and without regional nodal irradiation. *Radiat Oncol* 2015;10(1):200.
- [26] Fontanilla HP, Woodward WA, Lindberg ME, Kanke JE, Arora G, Durbin RR, Yu TK, Zhang L, Sharp HJ, Strom EA, Salehpour M. Current clinical coverage of Radiation Therapy Oncology Group-defined target volumes for postmastectomy radiation therapy. *Pract Radiat Oncol* 2012;2(3):201-209.
- [27] Jethwa KR, Kahila MM, Hunt KN, Brown LC, Corbin KS, Park SS, Yan ES, Boughey JC, Mutter RW. Delineation of internal mammary nodal target volumes in breast cancer radiation therapy. *Int J Radiat Oncol Biol Phys* 2017;97(4):762-769.
- [28] Offersen BV, Boersma LJ, Kirkove C, Hol S, Aznar MC, Sola AB, Kirova YM, Pignol JP, Remouchamps V, Verhoeven K, Weltens C. ESTRO consensus guideline on target volume delineation for elective radiation therapy of early stage breast cancer, version 1.1. *Radiother Oncol* 2016;118(1):205-208.
- [29] Nielsen MH, Berg M, Pedersen AN, Andersen K, Glavicic V, Jakobsen EH, Jensen I, Josipovic M, Lorenzen EL, Nielsen HM, Stenbygaard L. Delineation of target volumes and organs at risk in adjuvant radiotherapy of early breast cancer: National guidelines and contouring atlas by the Danish Breast Cancer Cooperative Group. *Acta Oncol* 2013;52(4):703-710.
- [30] Dijkema IM, Hofman P, Raaijmakers CP, Lagendijk JJ, Battermann JJ, Hillen B. Loco-regional conformal radiotherapy of the breast: Delineation of the regional lymph node clinical target volumes in treatment position. *Radiother Oncol* 2004;71(3):287-295.



1A: Orange: vessels; red: CTV; blue: PTV1.



1B: Orange: vessels; red: CTV; purple: PTV2.



1C :Orange: vessels; yellow: PTV3.

Figure 1: Contours




Figure 2: Example of dose distribution

Red: 90% isodose line; green: 95% isodose line.