

Antidepressant-like activity of Abietic Acid in Unstressed Mice and Chronic Unpredictable Mild Stressed Mice

Sudha, Dinesh Dhingra

Department of Pharmaceutical Sciences, Guru Jambheshwar University of Science and Technology, Hisar -125 001 (Haryana), India.

ABSTRACT

In the present study, abjetic acid was evaluated for antidepressant-like activity in normal mice and chronic unpredictable mild stress (CUMS) --induced depressed mice. Swiss albino male mice were subjected to CUMS for 21 successive days. Abietic acid (7.5, 15, and 30 mg/kg, p.o.) and fluoxetine (20 mg/kg, p.o.) per se were given for 3 successive weeks to separate groups of normal mice and CUMS-induced depressed mice. Tail suspension test (TST) and sucrose preference test were employed to study the effect of the drugs on the depressive-like behavior of mice. CUMS produced depressionlike behavior in mice. In TST, the immobility period was significantly decreased by abietic acid (30 mg/kg) and fluoxetine in normal mice as well as in CUMS-induced depressed mice as compared to their respective control groups. A decrease in sucrose preference due to CUMS was significantly restored by abietic acid (15 and 30 mg/kg) and fluoxetine. Locomotor activities of mice were not significantly changed by abietic acid and fluoxetine. Plasma nitrite, brain monoamine oxidase -A (MAO-A) activity, and brain malondialdehyde were significantly decreased; and brain catalase activity and reduced glutathione levels were significantly increased by abietic acid (15 and 30 mg/kg) and fluoxetine in both normal mice and CUMS-induced depressed mice. CUMS-induced increase in plasma corticosterone levels was significantly lowered by abietic acid (15 and 30 mg/kg) and fluoxetine. Abietic acid exerted significant antidepressant-like activity in both normal mice and CUMS-induced depressed mice possibly through mitigation of oxidative stress and decrease of brain MAO-A activity. Additionally, lowering of plasma corticosterone concentration by abietic acid in CUMS-induced depressed mice might also contribute to its antidepressant-like effect.

Key Words: Abietic acid, Antidepressant, Chronic unpredictable mild stress, Depression, Sucrose preference test, Tail suspension test.

eIJPPR 2021; 11(1):1-9

HOW TO CITE THIS ARTICLE: Sudha, Dinesh Dhingra (2021). "Antidepressant-like activity of Abietic Acid in Unstressed Mice and Chronic Unpredictable Mild Stressed Mice", International Journal of Pharmaceutical and Phytopharmacological Research, 11(1), pp.1-9.

INTRODUCTION

Depression is a common mental disorder worldwide, with more than 264 million people affected. In a depressed person, there is a decrease in mood, loss of interest and pleasure, low self-esteem, feeling of guilt, changes in sleep, and appetite [1-3]. It may be due to a decrease in norepinephrine, dopamine, and serotonin levels in the brain [4-6]. Stress has a major role in the development of human depression [7]. In a stressful situation, there is activation of the hypothalamic-pituitary-adrenal (HPA) axis, leading to elevation of the concentration of glucocorticoids like cortisol in primates or corticosterone in rodents, resulting in depression [8]. At present, the antidepressants available for clinical use include selective serotonin reuptake inhibitors, serotonin- noradrenergic reuptake inhibitors, tricyclic antidepressants, and monoamine oxidase inhibitors [9]. But these produce lots of side effects like cognitive impairment, sedation, fatigue, hypertensive crisis, sexual dysfunction [10, 11]. So plants and their bioactive compounds can be explored as an alternative and safe option for the treatment of depression. Hypericum perforatum has been proven to be an effective antidepressant for the treatment of depression in clinical studies [12]. Abietic acid is a tricyclic diterpene which is found in resins of different species of Pinus such as P. mugo, P. palustris, P. strobus, [13] P. sylvestris, P. roxburghii, [14] and also in Abies alba (Family -Pinaceae) [15]. Abietic acid has antioxidant and anticholinesterase, [16] antiepileptic, [17] anti-obesity [18] and

Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 04 October 2020; Revised: 23 December 2020; Accepted: 29 December 2020

Corresponding author: Sudha, Dinesh Dhingra

Address: Department of Pharmaceutical Sciences, Guru Jambheshwar University of Science and Technology, Hisar -125 001 (Haryana), India. E-mail: din_dhingra @ yahoo.com; din_dhingra @ rediffmail.com

anti-inflammatory [19] activities. But the effect of abietic acid on depression has not been mentioned in the literature. Therefore, abietic acid was studied for its effect on normal mice and CUMS-induced depressed mice.

MATERIALS AND METHODS

Experimental animals

Animals used were Swiss male albino mice (two to three months old and having body weight in the range 21-29 g). The mice were purchased from Disease Free Small Animal House, Lala Lajpat Rai University of Veterinary and Animal Sciences, Hisar (Haryana, India). Only male mice were used in the present study because the female sex hormone (estrogen) has been found to possess antidepressant activity [20]. The animals were housed in an air-conditioned room with a 12:12 h light-dark cycle. Drinking water and food were provided ad libitum to the animals, but the food was withdrawn 2 h before and 2 h after administration of the drugs, to rule out the effect of food on the absorption of the drugs [21]. The mice were acclimatized for 5 days before being used for the experiments. The experimental study was conducted between 9 AM and 5 PM. The experimental protocol and the number of animals to be used for the experiments were approved by Institutional Animals Ethics Committee (IAEC) in its meeting held on 6th October 2017 (Letter No. of Minutes- IAEC/2017/44-52, dated 02-11-2017).

Drugs and chemicals

Drugs used were abietic acid and fluoxetine (Sigma-Aldrich, St Louis, USA). Chemicals used were p-nitroso-N,N-dimethylaniline, N-(1- Naphthyl) ethylenediamine dihydrochloride, 5-hydroxytryptamine, thiobarbituric acid (HiMedia Laboratories Private Limited, Mumbai, India), sulfanilamide, metaphosphoric acid, potassium ferricyanide, hydrogen peroxide, trichloroacetic acid (CDH Private Limited, New Delhi, India), 5,5, Dithiobis-2-(nitro benzoic acid) (SRL Private Limited, Mumbai, India), sulfosalicylic acid (Spectrochem Private Limited, Mumbai, India). The kit was used for total protein estimation (Transasia Bio-medicals Ltd., Baddi (Solan), India).

Vehicle

The vehicle used for abietic acid was a 10% v/v solution of Tween 80 in distilled water. Normal saline was used as a vehicle for dissolving fluoxetine. An oral route of administration was employed for both the drugs.

Procedure for CUMS

CUMS was applied to the mice as per the method reported by Willner et al. (1987) with slight modifications [22]. The animals were exposed to various stressors only once a day for 3 successive weeks between 11 AM and 5 PM. The stressors were given to the mice in the following order:

Weeks	Day-1	Day-2	Day-3	Day-4	Day-5	Day-6	Day-7
1 st week	Ι	Е	F	0	T2	Х	T1
2 nd week	Ι	0	Х	T2	Е	T1	F
3 rd week	0	F	T1	Х	T2	Ι	Е

I— Immobilization for 2 h.

E— Exposure to an empty water bottle for 1 h.

F- Exposure to a foreign object (for example, a piece of plastic) for 24 h.

O— Overnight illumination.

T2— A tail pinch for 60 sec.

X— Tilted cage at 45 degrees for 7 h.

T1— A tail pinch for 30 sec.

Tail suspension test (TST)

The procedure for TST was the same as reported by Steru *et al.* [23]. The animals were suspended one by one fifty centimeters above the floor by adhesive tape which was placed approximately 1 cm from the tip of the tail. The immobility period of each mouse was recorded for 6 min. Immobility was characterized by lack of movements of limb or body except for respiratory movements which were produced during the passive and motionless hanging of animals.

Sucrose preference test

Sucrose preference test is mainly used to evaluate anhedonia, which is the core symptom of depression [24]. Initially, mice were exposed to a drinking water bottle containing 1% w/v sucrose solution until they are trained to take this solution. Then, the mice were not provided food and water for forty-eight hours but were allowed to take sucrose solution (1% w/v). After 3 days, mice were not given food and water for twenty-three hours; followed by a one-hour baseline test, where each mouse was exposed to two weighed bottles, one bottle filled with sucrose solution (1% w/v), and the other bottle filled with tap water. The sucrose preference was calculated as per the following formula:

Sucrose Preference

 $= \frac{\text{Sucrose solution intake (g)}}{[\text{Sucrose solution intake (g)} + \text{water intake (g)}]} \times 100$

On the 21st day, the sucrose preference test was again performed to evaluate the effect of CUMS and drugs.

International Journal of Pharmaceutical and Phytopharmacological Research (eIJPPR) | February 2021 | Volume 11| Issue 1| Page 1-9 Sudha, Dinesh Dhingra, Antidepressant-like activity of Abietic Acid in Unstressed Mice and Chronic Unpredictable Mild Stressed Mice

Measurement of locomotor activity

Horizontal locomotor activities of the mice were noted for 5 min [25] by using a photoactometer [INCO, Ambala (Haryana), India].

Details of experimental design

The animals were divided into the following 20 groups, each group having 8 mice.

For TST and locomotor activity

Groups 1–5: Tween 80 (10% v/v), abietic acid (7.5, 15 and 30 mg/kg) and fluoxetine (20 mg/kg) respectively were orally administered to mice for 3 successive weeks. The mice were subjected to TST one hour after vehicle/drug administration on the 22^{nd} day, and this was followed by testing of their locomotor scores 30 min later.

Groups 6–10: Tween 80 (10% v/v), abietic acid (7.5, 15 and 30 mg/kg) and fluoxetine (20 mg/kg) respectively were orally administered 30 min before giving stressors to the mice for 3 successive weeks. The mice were tested by TST 60 min after administration of vehicle/ drug on the 22^{nd} day, which was followed by testing of their locomotor scores 30 min later.

For sucrose preference test

Groups 11–20: Separate animals were used for this test, but the details of vehicle /drug treatments are the same as mentioned under groups 1 - 10. The test was carried out before the start of drug administration (baseline test) on the 1^{st} day and 60 min after drug administration on the 21^{st} day.

Estimation of various biochemical parameters Collection of blood samples and separation of plasma

After performing TST and testing of locomotor activity of mice of groups 1-10 on the 22^{nd} day, these mice were sacrificed on the 23^{rd} day by cervical dislocation and a blood sample (1.0 - 1.2 ml) was withdrawn from the carotid artery. Separation of plasma from the blood samples was carried out using a cooling centrifuge (Remi, Mumbai, India) at the speed of 2500 rpm for 10 min. Plasma was used for the estimation of nitrite and corticosterone.

Analysis of plasma nitrite levels

Plasma nitrite was analyzed by following the method reported by Green et al, 1982 [26].

Analysis of plasma corticosterone levels

Plasma corticosterone levels were determined by the method of Bartos and Pesez, 1979 [27].

Estimation of biochemical parameters in brain homogenate

Following the withdrawal of blood samples on the 23rd day, the mouse brain was isolated. This was followed by washing of brain samples with cold buffer solution (pH

7.4) containing 0.25 M sucrose - 0.1 M Tris-0.02 M ethylenediamine tetra-acetic acid. Then, the brain samples were weighed. Each brain sample was homogenized in 9 volumes of the above-mentioned buffer, followed by its centrifugation two times at the speed of 2500 rpm for 10 min at 4°C using a refrigerated centrifuge (Remi Instruments, Mumbai, India). The supernatant collected was centrifuged at 12,000 rpm for 20 min at 4°C. The precipitates (mitochondrial fraction) collected were used for the determination of monoamine oxidase- A (MAO-A) activity. The supernatant obtained was used to analyze malondialdehyde and, glutathione (GSH) levels; and catalase activity.

Analysis of MAO-A activity

MAO-A activity was measured spectrophotometrically by the method as reported in the literature [28, 29].

Analysis of total protein

Total protein was determined in the brain homogenate by using a kit (Erba, Transasia, Baddi (Solan, H.P.), using semi-automatic auto-analyzer (Chem5 plus-V2 semi-AutoAnalyzer; Erba Mannheim, Germany) [30].

Estimation of malondialdehyde (lipid peroxidation)

Malondialdehyde levels were determined spectrophotometrically by quantifying thiobarbituric acid-reactive substances (TBARS) by the method reported by Wills, 1965 [31].

Analysis of reduced glutathione (GSH)

Reduced glutathione levels were analyzed by the method reported by Jollow *et al.*, 1974 [32].

Analysis of catalase activity

Catalase activity was determined by the method reported by Claiborne, 1985 [33].

Statistical analysis of data

The mean and standard error of the mean of the data were calculated using Graphpad Instant statistical software. Then, the results were analyzed by one-way analysis of variance (ANOVA) followed by Tukey-Kramer multiple comparison test using the same statistical software.

RESULTS

Effect on immobility periods of mice in TST

Mice exposed to CUMS markedly (p <0.01; q= 6.141) enhanced immobility period of mice when tested in TST in comparison to control unstressed mice. Fluoxetine (20 mg/kg), the standard antidepressant drug employed, markedly (p <0.001) reduced the immobility periods of both unstressed mice (q= 16.764) and stressed mice (q=13.694) in comparison to their respective control International Journal of Pharmaceutical and Phytopharmacological Research (eIJPPR) | February 2021 | Volume 11| Issue 1| Page 1-9 Sudha, Dinesh Dhingra, Antidepressant-like activity of Abietic Acid in Unstressed Mice and Chronic Unpredictable Mild Stressed Mice

groups. The lowest dose (7.5 mg/kg) of abietic acid did not markedly affect immobility periods of both unstressed mice as well as CUMS exposed mice. But abietic acid when used in higher doses (15 and 30 mg/kg) markedly (p <0.001; q= 7.922 and 10.654 respectively) reduced immobility periods of CUMS exposed mice in comparison to their control group. On the other hand, abietic acid when used in the dose of 30 mg/kg markedly (p<0.001; q=6.540)) reduced immobility periods of unstressed mice in comparison to their control group (Table 1).

Table	1: Effect of	abietic acid	and fluox	etine on
	immobility	periods of n	nice in TS	Т

S.	Treatment for 3	Dose	Immobility
No.	successive weeks	(kg ⁻¹)	Period (sec)
1.	Control	10 ml	151.87 ± 5.27
2.	Control + CUMS	10 ml	176.87 ± 3.67**
3.	Fluoxetine (U)	20 mg	$83.62 \pm 3.50^{***}$
4.	Abietic acid (U)	7.5 mg	144.38 ± 6.94
5.	Abietic acid (U)	15 mg	135.62 ± 3.03
6.	Abietic acid (U)	30 mg	$125.25 \pm 2.99^{***}$
7.	Fluoxetine + CUMS	20 mg	$121.12 \pm 3.50^{\dagger\dagger\dagger}$

8.	Abietic acid + CUMS	7.5 mg	171 ± 2.51
9.	Abietic acid + CUMS	15 mg	$144.62 \pm 3.77^{\dagger\dagger\dagger}$
10.	Abietic acid + CUMS	30 mg	$133.5 \pm 3.63^{\dagger\dagger\dagger}$

n=8 in each group. U = unstressed mice; CUMS = chronic unpredictable mild stress.

Values are expressed as Mean \pm S.E.M. Data were analyzed by one-way ANOVA followed by Tukey–Kramer multiple comparison test. F (9,70) = 42.24: p<0.05

, * = p < 0.01, p < 0.001 respectively as compared to vehicle treated unstressed mice.

 $\dagger\dagger\dagger$ = p<0.001 as compared to vehicle treated stressed mice.

Effect on sucrose preference

Mice subjected to CUMS markedly (p <0.001; q= 9.714) reduced sucrose preference in comparison to control unstressed mice. In the baseline test, sucrose preference did not significantly differ among all the groups. In unstressed mice, all three doses of abietic acid, as well as fluoxetine (20 mg/kg), were devoid of any significant effect on sucrose preference. CUMS-induced decrease in sucrose preference was markedly reversed by abietic acid (15 and 30 mg/kg) and fluoxetine (20 mg/kg) [p <0.001; q= 10.088, 13.556 and 15.202 respectively] in comparison to CUMS exposed control mice (Table 2).

	Table 2:	: Effect a	of abietic	acid and	fluoxetine	on sucrose	preference
--	----------	------------	------------	----------	------------	------------	------------

S. No.	Treatment for 3 successive weeks	Dose (kg ⁻¹)	Sucrose preference (%)- Baseline test	Sucrose preference (%) - After 21 days
1.	Control	10 ml	61.13 ± 0.31	37.89 ± 3.19
2.	Control + CUMS	10 ml	60.89 ± 0.46	7.19 ± 0.65***
3.	Fluoxetine (U)	20 mg	61.76 ± 0.62	47.34 ± 5.30
4.	Abietic acid (U)	7.5 mg	61.35 ± 0.48	36.71 ± 0.28
5.	Abietic acid (U)	15 mg	61.28 ± 0.50	42.14 ±0.47
6.	Abietic acid (U)	30 mg	60.83 ± 0.41	47.33 ± 2.20
7.	Fluoxetine + CUMS	20 mg	61.45 ± 0.38	$55.23\pm4.29\dagger\dagger\dagger$
8.	Abietic acid + CUMS	7.5 mg	61.42 ± 0.44	20.45 ± 3.34
9.	Abietic acid + CUMS	15 mg	60.64 ± 0.74	$39.07\pm2.55\dagger\dagger\dagger\dagger$
10.	Abietic acid + CUMS	30 mg	61.09 ± 0.59	$50.03\pm4.42\dagger\dagger\dagger\dagger$

n =8 each group, U = unstressed mice; CUMS = chronic unpredictable mild stress.

Values are expressed as Mean ± S.E.M. The data were analyzed by one-way ANOVA followed by Tukey–Kramer multiple comparison test.

For Sucrose preference (%)- baseline test; F(9, 70) = 0.4246; p > 0.05.

For Sucrose preference (%)- after 21 days; F(9, 70) = 21.04; p < 0.05.

*** = p < 0.001, as compared to vehicle treated unstressed mice.

 $\dagger \dagger \dagger = p < 0.001$, as compared to vehicle treated stressed mice.

Effect on locomotor activity

Locomotor activities of unstressed mice and CUMS exposed mice were not significantly affected by abietic acid and fluoxetine in comparison to their respective controls (Table 3).

 Table 3: Effect of abietic acid and fluoxetine on

 locomotor activity

S.	Treatment for 3	Dose	No. of locomotor
No.	successive weeks	(kg-1)	counts

1.	Control	10 ml	246.87 ± 4.93
2.	Control + CUMS	10 ml	252.00 ± 3.12
3.	Fluoxetine (U)	20 mg	245.75 ± 4.61
4.	Abietic acid (U)	7.5 mg	254.75 ± 4.56
5.	Abietic acid (U)	15 mg	256.75 ± 7.00
6.	Abietic acid (U)	30 mg	238.37 ± 6.30
7.	Fluoxetine + CUMS	20 mg	256.62 ± 7.42
8.	Abietic acid + CUMS	7.5 mg	253.62 ± 6.83

9.	Abietic acid + CUMS	15 mg	269.12 ± 4.71
10.	Abietic acid + CUMS	30 mg	257.50 ± 6.71

n= 8 in each group. U = unstressed mice; CUMS = chronic unpredictable mild stress.

Values are expressed as Mean \pm S.E.M. Data were analyzed by one-way ANOVA followed by Tukey–Kramer multiple comparison test. F (9, 70) = 2.049; p > 0.05

Effect on plasma nitrite levels

CUMS markedly (p <0.001; q= 10.993) elevated plasma nitrite levels in mice. There was a significant (p <0.001) decrease of plasma nitrite levels by fluoxetine (20 mg/kg) in both unstressed mice (q= 7.441) as well as CUMS exposed mice (q= 19.112) in comparison to their respective controls. Abietic acid used in the lowest dose (7.5 mg/kg) was devoid of any significant effect on plasma nitrite levels in unstressed mice as well as in CUMS exposed mice. Abietic acid used in the dose of 30 mg/kg markedly (p <0.001; q= 7.543) reduced plasma nitrite levels in unstressed mice in comparison to their control mice. Abietic acid used in higher doses (15 and 30 mg/kg) markedly (p<0.001; q= 10.298 and 16.769 significantly) reduced nitrite levels in mice exposed to CUMS in comparison to CUMS exposed control mice (Table 4).

Table 4: Effect of abietic acid and fluoxetine on plasma nitrite and corticosterone levels

S No	Treatment for 3 successive	Dese (kg·l)	Plasma nitrite level	Plasma Corticosterone
5. NO.	weeks	Dose (kg)	(µg/ml)	Levels (µg/ml)
1.	Control	10 ml	22.83 ± 0.93	22.17 ± 0.57
2.	Control + CUMS	10 ml	$31.68 \pm 0.97^{***}$	$28.18 \pm 0.52^{***}$
3.	Fluoxetine (U)	20 mg	$16.84 \pm 0.91^{***}$	19.80 ± 0.62
4.	Abietic acid (U)	7.5mg	20.94 ± 0.88	21.87 ± 0.71
5.	Abietic acid (U)	15 mg	20.82 ± 0.68	20.68 ± 0.53
6.	Abietic acid (U)	30 mg	$16.76 \pm 0.37^{***}$	20.45 ± 0.58
7.	Fluoxetine + CUMS	20 mg	$16.30 \pm 0.61^{\dagger\dagger\dagger}$	$22.13 \pm 0.64^{\dagger\dagger\dagger}$
8.	Abietic acid + CUMS	7.5 mg	28.68 ± 0.88	$25.16 \pm 0.56^{\dagger}$
9.	Abietic acid + CUMS	15 mg	$23.40\pm0.87\dagger\dagger\dagger$	$24.12 \pm 0.47^{\dagger\dagger\dagger}$
10.	Abietic acid + CUMS	30 mg	$18.19 \pm 0.73^{\dagger\dagger\dagger}$	$21.56 \pm 0.48^{\dagger\dagger\dagger}$

n=8 in each group. U = unstressed mice; CUMS = chronic unpredictable mild stress.

Values are expressed as Mean ± S.E.M. Data were analyzed by one-way ANOVA followed by Tukey-Kramer multiple comparison test.

For plasma nitrite levels: F (9,70) = 41.726; p < 0.05

For plasma corticosterone levels: F(9, 70) = 18.289; p<0.05

***= p<0.001 as compared to vehicle-treated unstressed mice.

 \dagger , \dagger , \dagger , \dagger , \dagger , \dagger = p<0.05, p<0.001 respectively as compared to vehicle-treated stressed mice.

Effect on plasma corticosterone levels

CUMS exposed mice showed marked (p<0.001; q= 10.500) elevation in corticosterone levels in mice in comparison to control unstressed mice. Abietic acid at all the doses (7.5, 15 and 30 mg/kg) as well as fluoxetine (20 mg/kg) markedly (p<0.05, p<0.001, p<0.001 and p<0.001 respectively; q= 5.287, 7.046, 11.574 and 10.570 respectively) decreased corticosterone levels in CUMS exposed mice in comparison to CUMS exposed control mice. There was no significant effect of abietic acid and fluoxetine on corticosterone levels in unstressed mice in comparison to control unstressed mice (Table 4).

Effect on brain MAO-A activity

CUMS markedly (p<0.001; q= 24.920) elevated MAO-A activity in comparison to control unstressed mice. There was a marked (p<0.001) decrease in MAO-A activity by fluoxetine (20 mg/kg) in unstressed mice (q= 8.293) as well as in mice subjected to CUMS (q= 43.979) in comparison to their respective controls. Abietic acid used in the lowest dose (7.5 mg/kg) was devoid of any marked effect on MAO-A activity in unstressed mice as well as in CUMS exposed mice. But abietic acid used in the dose of 30 mg/kg markedly (p<0.001; q= 8.644) reduced brain MAO-A activity in unstressed mice in comparison to their control. The higher doses (15 and 30 mg/kg) of abietic acid significantly (p<0.001; q= 9.533 and 41.154 respectively) reduced MAO-A activity in CUMS exposed mice in comparison to control mice exposed to CUMS (Table 5).

Table 5: Effect of abietic acid and fluoxetine on brain MAO-A activity and malondialdehyde levels

S. No.	Treatment for 3 successive	Dose	MAO-A activity	Malondialdehyde levels
	weeks	(kg ⁻¹)	(nmol/mg protein)	(nmols/mg protein)
1.	Control	10 ml	67.97 ± 0.54	1.13 ± 0.003

International Journal of Pharmaceutical and Phytopharmacological Research (eIJPPR) | February 2021 | Volume 11| Issue 1| Page 1-9 Sudha, Dinesh Dhingra, Antidepressant-like activity of Abietic Acid in Unstressed Mice and Chronic Unpredictable Mild Stressed Mice

2.	Control + CUMS	10 ml	$89.87 \pm 0.57^{***}$	$2.20 \pm 0.08^{***}$
3.	Fluoxetine (U)	20 mg	$60.69 \pm 1.23^{***}$	$0.84 \pm 0.04^*$
4.	Abietic acid (U)	7.5mg	66.92 ± 0.69	1.06 ± 0.02
5.	Abietic acid (U)	15 mg	64.48 ± 0.53	1.09 ± 0.06
6.	Abietic acid (U)	30 mg	60.38 ±0.57***	$0.76 \pm 0.03^{**}$
7.	Fluoxetine + CUMS	20 mg	$51.22 \pm 1.21^{\dagger\dagger\dagger}$	$1.41 \pm 0.05^{\dagger\dagger\dagger}$
8.	Abietic acid + CUMS	7.5 mg	87.80 ± 0.64	2.13 ± 0.12
9.	Abietic acid + CUMS	15 mg	$81.50 \pm 1.38^{\dagger\dagger\dagger}$	$1.89 \pm 0.05^{\dagger\dagger}$
10.	Abietic acid + CUMS	30 mg	$53.71 \pm 0.84^{\dagger\dagger\dagger}$	$1.58 \pm \overline{0.02^{\dagger\dagger\dagger}}$

n=8 in each group. U = unstressed mice; CUMS = chronic unpredictable mild stress.

Values are expressed as Mean ± S.E.M. Data were analyzed by one-way ANOVA followed by Tukey-Kramer multiple comparison test.

For MAO- A activity: F (9,70) = 257.89; p<0.05

For Malondialdehyde levels: F(9,70) = 86.26; p<0.05

*, ** and ***= p<0.05, p<0.01, p<0.001 respectively as compared to vehicle-treated unstressed mice.

 \dagger , \dagger , \dagger , \dagger , \dagger , = p<0.01, p<0.001 respectively as compared to vehicle treated stressed mice.

Effect on brain malondialdehyde levels:

There was a significant (p < 0.001; q = 18.553) increase in malondialdehyde levels in CUMS exposed mice in comparison to unstressed control mice. Fluoxetine (20 mg/kg) and abietic acid (15 and 30 mg/kg) significantly (p<0.001, p<0.01 and p<0.001 respectively; q= 13.705,5.366 and 10.731 respectively) decreased malondialdehyde levels in CUMS exposed mice in comparison to control mice subjected to CUMS. There was no significant effect of the lowest dose (7.5 mg/kg) of abietic acid on malondialdehyde levels in mice subjected to CUMS in comparison to control mice subjected to CUMS. Fluoxetine (20 mg/kg) and only the highest dose (30 mg/kg) of abietic acid markedly (p<0.05 and p<0.01 respectively; q= 4.978 and 6.228 respectively) reduced malondialdehyde levels in unstressed mice in comparison to their control (Table 5).

Effect on brain reduced glutathione levels:

There was a significant (p<0.001; q= 34.829) decrease in reduced glutathione levels in CUMS exposed mice in comparison to control unstressed mice. Fluoxetine (20 mg/kg) and only the highest dose (30 mg/kg) of abietic acid markedly (p<0.001; q= 61.698 and 8.274 respectively) increased brain reduced glutathione levels in unstressed mice in comparison to their control. Abietic acid (15 and 30 mg/kg) and fluoxetine (20 mg/kg) markedly (p<0.001; q= 8.558, 31.162 and 33.123 respectively) increased brain reduced glutathione levels in comparison to control mice subjected to CUMS (Table 6).

S. No.	Treatment for 3 successive	Dose	GSH Levels (µmol/mg	Catalase activity (µg/mg
	weeks	(kg ⁻¹)	protein)	protein)
1.	Control	10 ml	0.247 ± 0.007	45.47 ± 0.38
2.	Control + CUMS	10 ml	$0.094 \pm 0.001^{***}$	$23.90 \pm 1.00^{***}$
3.	Fluoxetine (U)	20 mg	$0.517 \pm 0.004^{***}$	$89.60 \pm 0.37^{***}$
4.	Abietic acid (U)	7.5 mg	0.247 ± 0.005	46.34 ± 0.44
5.	Abietic acid (U)	15 mg	0.264 ± 0.007	47.91 ± 1.55
6.	Abietic acid (U)	30 mg	$0.283 \pm 0.003^{***}$	$72.39 \pm 1.00^{***}$
7.	Fluoxetine + CUMS	20 mg	$0.239 \pm 0.003^{\dagger\dagger\dagger}$	$68.58 \pm 2.18^{\dagger\dagger\dagger}$
8.	Abietic acid + CUMS	7.5 mg	0.104 ± 0.003	28.58 ± 1.17
9.	Abietic acid + CUMS	15 mg	$0.132 \pm 0.002^{\dagger\dagger\dagger}$	$31.43 \pm 1.30^{\dagger\dagger\dagger}$
10.	Abietic acid + CUMS	30 mg	$0.231 \pm 0.003^{\dagger\dagger\dagger}$	$40.82 \pm 0.56^{\dagger\dagger\dagger}$

Table 6: Effect of abietic acid and fluoxetine on brain reduced glutathione levels and catalase activity

n=8 in each group. U = unstressed mice; CUMS = chronic unpredictable mild stress.

Values are expressed as Mean ± S.E.M. Data were analyzed by one-way ANOVA followed by Tukey–Kramer multiple comparison test.

For reduced glutathione: F (9,70) = 791.03; p < 0.05

For catalase activity: F (9,70) = 344.54; p < 0.05

*** = p<0.001 as compared to vehicle-treated unstressed mice.

 \dagger \dagger \dagger = p<0.001 as compared to vehicle-treated stressed mice.

Effect on brain catalase activity:

There was a significant (p<0.001; q=18.915) decrease in brain catalase activity in mice subjected to CUMS in comparison to control unstressed mice. Fluoxetine (20 mg/kg) and only the highest dose (30 mg/kg) of abietic acid markedly (p<0.001; q= 38.709 and 23.610 respectively) increased brain catalase activity in unstressed mice in comparison to their control. Abietic acid (15 and 30 mg/kg) and fluoxetine (20 mg/kg) markedly (p<0.001; q= 6.601, 14.837 and 39.162 respectively) increased brain catalase activity in CUMS exposed mice in comparison to control mice subjected to CUMS (Table 6).

DISCUSSION

The present study demonstrated the effects of abietic acid (7.5, 15, and 30 mg/kg, p.o) on depression-like behavior in unstressed mice and CUMS-induced depressed mice using the behavioral models (TST and sucrose preference test). CUMS-induced model of depression is broadly employed for screening of drugs for antidepressant activity in mice and rats. This model results in depression-like behavior similar to that observed in humans [34, 35]. In TST, CUMS markedly increased immobility periods of mice, which indicated the development of depression-like behavior in mice. Immobility time in TST indicates despair behavior, which is a notable symptom of depression. There was a marked decrease in immobility periods of CUMS exposed mice by abietic acid (15 and 30 mg/kg) and fluoxetine (20 mg/kg), indicating antidepressant-like effects of these drugs. But in unstressed mice, fluoxetine (20 mg/kg) and only the highest dose (30 mg/kg) of abietic acid produced significant antidepressant-like activity. Fluoxetine was used as a standard drug to validate the employed animal models of depression. Abietic acid and fluoxetine did not produce any marked effect on locomotor activities of unstressed mice as well as in CUMS exposed in comparison to their respective controls, indicating that antidepressant-like effects of these drugs were specific and not due to their stimulant or depressant effects on the central nervous system. The sucrose preference test is considered the most reliable behavioral test to study the effects of drugs on CUMS-induced depression in mice and rats. A decrease in sucrose preference indicates anhedonia, that is, loss of interest or pleasure [34], which is commonly seen in depressive patients. There is damage to nerve cells by CUMS in the neural reward system linked to serotonergic and dopaminergic systems; and this leads to the abolition of experiencing happiness or pleasure [36]. In the present study, sucrose preference markedly decreased in CUMS exposed mice in comparison to control unstressed mice. Fluoxetine (20 mg/kg) and abietic acid (15 and 30 mg/kg) significantly reversed CUMS-induced decrease in sucrose preference. This indicated the antidepressant-like effects of abietic acid and fluoxetine in mice subjected to CUMS. Restoration of reduced sucrose preference by fluoxetine is also supported by the literature [37]. HPA axis plays an important role in bringing out various physiological changes in response to stressful stimuli [38]. Stress-induced activation of the HPA axis leads to a rise in plasma corticosterone levels in rodents. Cortisol levels elevate in depressive patients. Antidepressants such as fluoxetine decrease cortisol levels [39]. In the current study, elevated plasma corticosterone levels in CUMS exposed mice were markedly decreased by abietic acid and fluoxetine. In unstressed mice, no significant effect on plasma corticosterone level was shown which indicates that hyperactivation of the HPA axis occurs only in stressful conditions. Oxidative stress leads to the production of reactive oxygen species and decreases the levels of endogenous antioxidants, resulting in depression [40]. There is damage to lipids and proteins; a decrease in antioxidant enzymes and reduced glutathione levels in rodents' brains by repeated and unpredictable stress [41]. Catalase and glutathione are antioxidant defense systems against reactive oxygen species [42]. CUMS leads to a significant increase in reactive oxygen species accumulation in the brain which leads to abnormality in the normal physiology of the central nervous system [43]. In the present study, CUMS markedly increased brain lipid peroxidation and plasma nitrite levels; and diminished brain reduced glutathione levels and catalase activity. These findings are supported by earlier studies [44]. Abietic acid (15 and 30 mg/kg, p.o) markedly reversed CUMS-induced changes in oxidative stress parameters. But in unstressed mice, abietic acid in the highest dose (30 mg/kg) employed showed antioxidant activity per se. Thus, the antidepressant-like effect of abietic acid might be due to the amelioration of oxidative stress.CUMS markedly enhanced brain MAO-A activity. This finding is also supported by the literature [45]. Abietic acid (15 and 30 mg/kg) and fluoxetine (20 mg/kg) markedly decreased brain MAO-A activity in CUMSinduced depressed mice. But in unstressed mice, abietic acid in the highest dose (30 mg/kg) employed significantly reduced brain MAO-A activity. Therefore, the antidepressant-like activity of abietic acid in CUMSinduced depressed mice and also in unstressed mice might also be due to brain MAO-A inhibition. Literature also supports MAO-A inhibition by fluoxetine [46]. Thus, it may be concluded that the administration of abietic acid for 3 consecutive weeks exerted marked antidepressant-like activity in CUMS-induced depressed mice as well as in unstressed mice. This antidepressant-like effect of abietic acid might be to decrease oxidative stress, restorative antioxidant enzymes, and brain MAO-A inhibition. Additionally, in CUMS-induced depressed mice, abietic acid showed antidepressant-like activity through a decrease of plasma corticosterone levels.

International Journal of Pharmaceutical and Phytopharmacological Research (eIJPPR) | February 2021 | Volume 11 | Issue 1| Page 1-9 Sudha, Dinesh Dhingra, Antidepressant-like activity of Abietic Acid in Unstressed Mice and Chronic Unpredictable Mild Stressed Mice

CONCLUSION

In the present research study, abietic acid exerted significant antidepressant-like activity in both normal mice and CUMS-induced depressed mice possibly through alleviation of oxidative stress and decrease of brain MAO-A activity. Additionally, lowering of plasma corticosterone concentration by abietic acid in CUMS-induced depressed mice might also contribute to its antidepressant-like effect.

REFERENCES

- WHO. Depression- Fact sheet. January 2020. https://www.who.int/news-room/factsheets/detail/depression.
- [2] Mosarrezaii A, Kargar K. Frequency of Depression in Patients with Seizure Referring to the Urmia Neuromedical Clinic and Some of the Factors Affecting it. J. Adv. Pharm. Educ. Res. 2018;8(2):16-20.
- [3] Fouladi A, Goli S. Comparing working memory, verbal memory and keeping attention in the manic phase and depression in bipolar disorder. J. Adv. Pharm. Educ. Res. 2018;8(2):82-4.
- [4] Raeisi M, Navidian A, Rezaee N. Comparison of the Effect of Nurses' Education on Stress, Anxiety, and Depression of Family Caregivers of Patients Hospitalized with Schizophrenia Disorder. Arch. Pharm. Pract. 2020;11(1):82-7.
- [5] Abdelbasset WK, Nambi G, Moawd SA, Alrawaili SM, Elnegamy TE, Elshehawy AA, Eltrawy AH. Evaluating the depression status following backward walking exercise in diabetic neuropathy patients. Archives of Pharmacy Practice. 2020;11(1):105-9.
- [6] Gold PW, Goodwin FK, Chrousus G. Clinical and biochemical manifestations of depression in relation to the neurobiology of stress: Part 1. N Engl J Med 1988;319:348–53.
- [7] Nestler EJ, Barrot M, DiLeone RJ, Eisch AJ, Gold SJ, Monteggia LM. Neurobiology of depression. Neuron 2002;34:13-25.
- [8] Johnson SA, Fournier NM, Kalynchuk LE. Effect of different doses of corticosterone on depression-like behavior and HPA axis responses to a novel stressor. Behav Brain Res 2006; 168:280–88.
- [9] Lee B, Sur B, Yeom M, Shim I, Lee H, Hahm DH. Effect of berberine on depression- and anxiety-like behaviors and activation of the noradrenergic system induced by development of morphine dependence in rats. Korean J. Physiol. Pharmacol 2012;16:379–86.
- [10] Sathyanarayana Rao TS, Yeragani VK. Hypertensive crisis and cheese. Indian J Psychiatry 2009;51:65–6.
- [11] Papakostas GI. Tolerability of modern antidepressants. J Clin Psychiatry 2008;69:8–13.

- [12] Rahimi R, Nikfar S, Abdollahi M. Efficacy and tolerability of Hypericum perforatum in major depressive disorder in comparison with selective serotonin reuptake inhibitors: A meta-analysis. Prog Neuropsychopharmacol Biol Psychiatry 2009;33:118-21.
- [13] Corrigan D. Abies, Picea and Pinus Species. In: Peter A. G. M. De Smet, Keller K, Hansel R, Frank Chandler R. Adverse Effects of Herbal Drugs, Vol. 2. Verlag Berlin Heidelberg: Springer; 1993. p. 91-104.
- [14] Kaushik P, Kaushik D, Khokra SL. Ethnobotany and phytopharmacology of Pinus roxburghii Sargent: a plant review. J Integr Med 2013;11:371-76.
- [15] Wajs-Bonikowska A, Smeds A, Willfor S. Chemical composition and content of lipophilic seed extractives of some Abies and Picea species. Chem Biodivers 2016;13:1194-1201.
- [16] Ramnath MG, Thirugnanasampandan R, Sadasivam M, Subramaniam Mohan P. Antioxidant, antibacterial and antiacetylcholinesterase activities of abietic acid from Isodon wightii (Bentham) H. Hara. Free Rad. Antiox 2015;5(1):1-5.
- [17] Talevi, A, Cravero MS, Castro EA, Bruno-Blanch LE. Discovery of anticonvulsant activity of abietic acid through application of linear discriminant analysis. Bioorg Med Chem Lett 2017;17:1684-1690.
- [18] Hwang KH, Ahn JY, Kim S, Park JH, Ha TY. Abietic acid has an anti-obesity effect in mice fed a high-fat diet. J Med Food 2011;14:1052-56.
- [19] Fernandez MA, Tornos MP, Gracia MD, Heras BDL, Villar AM, Saenz MT. Anti-inflammatory activity of abietic acid, a diterpene isolated from Pimenta racemosa var. grissea. J Pharm Pharmacol 2001;53:867-72.
- [20] Pu W, Changhong L, Lei L, Xingyi Z, Bingzhong R, Bingjin L. The Antidepressant-like effects of estrogen-mediated ghrelin. Curr Neuropharmacol 2015;13:524-35.
- [21] Koziolek M, Alcaro S, Augustijns P, Basit AW, Grimm M, Hens B, Hoad CL, Jedamzik P, Madla CM, Maliepaard M, Marciani L. The mechanisms of pharmacokinetic food-drug interactions – A perspective from the UNGAP group. Eur J Pharm Sci 2019;134:31-59.
- [22] Willner P, Towell A, Sampson D, Muscat R, Sopholeous S. Reduction of sucrose preference by chronic mild stress and its restoration by a tricyclic antidepressant. Psychopharmacology 1987;93:358-64.
- [23] Steru L, Chermat R, Thierry B, Simon P. The tail suspension test: A new method for screening antidepressants in mice. Psychopharmacology (Berl) 1985;85:367-70.

- [24] Willner P, Towell A, Sampson D, Sophokleous S, Muscat R. Reduction of sucrose preference by chronic unpredictable mild stress, and its restoration by a tricyclic antidepressant. Psychopharmacology (Berl) 1987;93:358-64.
- [25] Dhingra D, Bansal Y. Antidepressant-like activity of beta-carotene in unstressed and chronic unpredictable mild stressed mice. J Funct Foods 2014;7:425-34.
- [26] Green LC, Wagner DA, Glogowski J, Skipper PL, Wishnock JS, Tannenbaum SR. Analysis of nitrate, nitrite, and [N-15N]-labeled nitrate in biological fluids. Anal Biochem 1982;126:131-38.
- [27] Bartos J, Pesez M. Colorimetric and fluorimetric determination of steroids. Pure Appl Chem 1979;151:2157-69.
- [28] Charles M, McEwen J. MAO activity in rabbit serum. In: H. Tabor, C.W. Tabor, editor. Methods in Enzymology (XVIIB). New York and London: Academic Press; 1977. p. 692-98.
- [29] Schurr A, Liven A. Differential inhibition of mitochondrial monoamine oxidase from brain by hashish components. Biochem Pharmacol 1976;25:1201-03.
- [30] Gornall AG, Bardawill CJ, David MM. Determination of serum protein by means of the biuret reaction. J Biol Chem 1949;177:751-66.
- [31] Wills ED. Mechanisms of lipid peroxide formation in tissues. Role of metals and haematin proteins in the catalysis of the oxidation of unsaturated fatty acids. Biochim Biophys Acta 1965;98:238-51.
- [32] Jollow DJ, Mitchell JR, Zampaglione N, Gillette JR. Bromobenzene induced liver necrosis: Protective role of glutathione and evidence for 3,4bromobenzenoxide as the hepatotoxic metabolite. Pharmacol 1974;11:151-69.
- [33] Claiborne A. Catalase activity. In Greenwald RA, editor. CRC Handbook of Methods for Oxygen Free Radical Research. Florida: CRC Press, Boca Raton; 1985. p. 283-84
- [34] Willner P, Towell A, Sampson D, Sophokleous S, Muscat R. Reduction of sucrose preference by chronic unpredictable mild stress, and its restoration by a tricyclic antidepressant. Psychopharmacology (Berl) 1987;93:358-64.
- [35] Willner P. Validity, reliability, and utility of the chronic mild stress model of depression. A 10-year review and evaluation. Psychopharmacology (Berl) 1997;134:319-29.
- [36] Bekris S, Antoniou K, Daskas S, Papadopoulou-

Daifoti Z. Behavioural and neurochemical effects induced by chronic mild stress applied to two different rat strains. Behav Brain Res 2005;161:45-59.

- [37] Deng XY, Li HY, Chen JJ, Li RP, Qu R, Fu Q, Ma SP. Thymol produces an antidepressant-like effect in a chronic unpredictable mild stress model of depression in mice. Behav. Brain Res. 2015;291:12–19.
- [38] Pan Y, Zhang WY, Xia X, Kong LD. Effects of icariin on hypothalamic-pituitary-adrenal axis action and cytokine levels in stressed Sprague-Dawley rats. Biol Pharm Bull 2006;29: 2399–03.
- [39] Chau DT, Rada PV, Kim K, Kosloff RA, Hoebel BG. Fluoxetine alleviates behavioral depression while decreasing acetylcholine release in the nucleus accumbens shell. Neuropsychopharmacol 2011;36:1729–37.
- [40] Maes M, Galecki P, Chang YS, Berk M. A review on the oxidative and nitrosative stress (O & NS) pathways in major depression and their possible contribution to the (neuro)degenerative processes in that illness. Prog. Neuropsychopharmacol Biol Psychiatry 2011;35:676–92.
- [41] Sahin E, Gumuslu S. Alterations in brain antioxidant status, protein oxidation, and lipid peroxidation in response to different stress models. Behav Brain Res 2004;155:241–48.
- [42] Ng F, Berk M, Dean O, Bush AI. Oxidative stress in psychiatric disorders: Evidence base and therapeutic implications. Int. J. Neuropsychopharmacol 2008;11:851-76.
- [43] Madrigal JL, Olivenza R, Moro MA, Lizasoain I, Lorenzo P, Rodrigo J, Leza JC. Glutathione depletion, lipid peroxidation, and mitochondrial dysfunction are induced by chronic stress in rat brain. Neuropsychopharmacol 2001;24(4):420-429.
- [44] Bhatt S, Mahesh R, Devadoss A, Dhar AK. Neuropharmacological evaluation of a novel 5-HT3 receptor antagonist (6g) on chronic unpredictable mild stress-induced changes in behavioral and brain oxidative stress parameters in mice. Indian J Pharmacol 2014;46:191-96.
- [45] Liu J, Qiao W, Yang Y, Ren L, Sun Y, Wang S. Antidepressant-like effect of the ethanolic extract from Suanzaorenhehuan formula in mice models of depression. J Ethnopharmacol 2012;141:257–64.
- [46] Mukherjee J, Yang Z-Y. Monoamine oxidase- A inhibition by fluoxetine: an in vitro and in vivo study. Synapse 1999;31:285-89.