



International Journal of Pharmaceutical and Phytopharmacological Research (eIJPPR)

[Impact Factor – 0.852]

Journal Homepage: www.eijppr.com

Research Article

In vitro Evaluation of Antimicrobial Activity of Methanolic Extract of *Murraya Koenigii* Leaves (Curry Leaves)

Jayasree Tirumalasetti^{1,*}, Anuradha Basavaraju², Harini³, Praveena⁴¹ Professor and Head, Department of Pharmacology, Mamata Medical College, Khammam – 507 002, Andhra Pradesh, India.² Professor, Department of Microbiology, Mamata Medical College, Khammam – 507 002, Andhra Pradesh, India.³ MD Student, Department of Pharmacology, Mamata Medical College, Khammam – 507 002, Andhra Pradesh, India.⁴ PhD Scholar, Department of Microbiology, Mamata Medical College, Khammam – 507 002, Andhra Pradesh, India.

Article info

Article History:
Received 30 May 2014
Accepted 30 Oct 2014

Keywords:

Anti microbial activity, *Candida albicans*, disc diffusion method, gram negative bacteria, gram positive bacteria, Soxhlets apparatus.

Abstract

The microorganisms have developed resistance to many commercial antibiotics due to the indiscriminate use of antibacterial drugs. Therefore, investigation of new molecules has become necessary. As medicinal plants have chemical compounds with antimicrobial action in abundance, vast research is going on those plants. One of such plant is *Murraya Koenigii*. *Murraya Koenigii* also known as sweet neem is used in traditional medicine to treat various diseases ascardio protective, antimicrobial, antifungal, and antiulcer, antioxidant, antifungal, and antiulcer. The Methanolic Extract of *Murraya Koenigii* Leaves (curry leaves), were investigated against different species of Gram –ve bacteria i.e. *Pseudomonas aeruginosa*, *Escherichia Coli*, *Klebsiella* and Gram +ve *staphylococcus aureus* and fungus *Candida albicans* in our study. The screening was performed by standard disc diffusion method. It demonstrated moderate antibacterial activity against *staphylococcus aureus* having the diameter of zone of inhibition of 15 mm and good antibacterial activity against *E.coli* with zone of inhibition of 20 mm and *pseudomonas aeruginosa* with zone of inhibition of 10 mm and no zone of inhibition on *Klebsiella*, The leaf extract has shown anti fungal activity on *Candida albicans* with 16mm zone of inhibition.

1. INTRODUCTION

Various antimicrobial peptides, proteins, and small molecular weight organic substances are present in Plants acting as host defense mechanisms^{1,2}. Medicinal herbs and plant products were used in treating a wide spectrum of infections and other diseases in traditional medicine like Ayurveda and Unani^{3,4}. As many microorganisms developed resistance due to the indiscriminate use of antibacterial drugs, there is a need to develop new molecules with minimum side effects. Therefore investigation of the chemical compounds within medicinal plants has become desirable⁵. *Murraya Koenigii* is a medically important herb of Indian origin (Family-Rutaceae). Meethineem (Hindi), karivepaaku (Telugu) its extract is known to show antidiabetic⁶, cardio protective, antimicrobial, antifungal, and antiulcer, antioxidant⁷, anti-inflammatory⁹, hypolipidemic activities¹⁰, anticancer¹¹, analgesic properties and indigenous medicine as tonic, stomach stimulant and carminative. This plant has been used in poly herbal formations like Siddha medicine. The extracts were also proved to be effective in gastric ulceration and was suggested as protective as ranitidine.¹² Biologically active carbazole alkaloids are reported to have antimicrobial properties.¹³ Curry leaves have been reported to contain tocopherol, b-carotene, lutein and alkaloids.¹⁴ The aim and Purpose of this study was to evaluate antimicrobial activity of *Murraya Koenigii* (curry leaves) against different bacteria and fungi- *staphylococcus aureus*, *E.coli*, *Klebsiella* species and *Candida albicans*.

*Corresponding Author:

Dr. Jayasree Tirumalasetti,
Professor and Head, Department of pharmacology,
Mamata Medical College, Khammam-507002,
Andhra Pradesh, India.
Contact No. +91-9848127116
Email: drtjayasreemd@gmail.com

2. MATERIALS AND METHODS

2.1 Plant material

Murray Koenigii curry leaves were procured in bulk from a farm and the leaves were identified for the strain and the species by a botanist^{5,7}. The leaves were washed to remove dirt. After washing leaves were shade and dried, followed by hot air oven drying at 40^o C for 7 to 8 hours. Powder was made from dried leaves and stored in air tight containers for further analysis. Antimicrobial analysis (0-6 months) were done in the leaf powder *Murry Koenigii* 20 %, were taken for the study.

2.2 Extract Preparation

The extraction of *Murray Koenigii* curry leaves was done with 95% ethanol by Soxhlets apparatus in department of Pharmacology. The extract were dried under vacuum, stored at room temperature and protected from direct sunlight and stored in air tight containers for further analysis.

2.3 Phytochemical Analysis

Preliminary photochemical analysis was made for presence of alkaloid, flavonoids, carbohydrates, glycosides, proteins and amino acids, steroids, vitamin C, fat and fixed oil. Briefly, following tests has been performed for identifying the class of compounds.

a) Test for alkaloids

Two ml of extract was acidified with a few drops of dilute hydrochloric acid and then 1 ml of Dragendorffs reagent was added. The appearance of orange to red precipitate indicates the presence of alkaloids.

b) Test for tannins

To 2 ml of extract, a few drops of 10% lead acetate were added. The appearance of white precipitate indicates the presence of tannins.

c) Test for saponins

To 1 ml of extract taken in a measuring jar, 9 ml of distilled water was added and shaken vigorously for 15 s and the extract was

allowed to stand for 10 min. Formation of stable foam (1 cm) indicates the presence of saponins.

d) Test for steroids

Chloroform 10 ml was added to 2 ml of extract. To this extract, 1 ml of acetic anhydride was added: then, 2 ml of concentrated sulphuric acid was added along the sides of the test tube. Colour formation at the junction is noted. The appearance of blue-green colour indicates the presence of steroids.

e) Test for triterpenoids

Presence of triterpenoids is tested same as that of steroids. The appearance of red, pink or violet colour at the junction indicates the presence of triterpenoids.

f) Test for cardiac glycosides

To 1 ml of extract, a few drops of glacial acetic acid and ferric chloride, and 3-4 drops of concentrated sulphuric acid were added. The appearance of blue-green colour indicates the presence of glycosides.

2.4 Organisms Used

The test organisms included four clinical isolates of *Staphylococcus aureus* which is gram positive and *Pseudomonas aeruginosa*, *E. coli* and *Klebsiella* which are gram negative organisms and *Candida albicans* a fungus. An attempt has been made to test the in vitro antibacterial activity of above leaf extract against above said pathogens. Clinical isolates of these organisms isolated during the study period from pus for *Staphylococcus aureus* and from urine for *Escherichia coli* and *Klebsiella* and *Pseudomonas* sp. were utilized for this study.

2.5 Micro Dilution Assay

The minimum inhibitory concentration was defined as the lowest concentration of the compound to inhibit the growth of microorganisms. The minimum inhibitory concentration values were determined by broth dilution assay of micro dilution assay. Varying Concentrations of the extracts (500mg/ml, 400mg/ml, 300mg/ml, 200mg/ml, 100mg/ml and 50mg/ml) were prepared. 0.1ml of standardized test organism of controls was equally set up by using solvents and test organisms without extract. The inoculum was prepared from fresh overnight broth culture in nutrient broth. Plates were incubated for 24 h at 37°C. MIC was recorded as the lowest extract concentration demonstrating no visible growth in the broth

2.6 Test Micro-organisms

Antimicrobial activity of plant extract in different concentrations — 10000,30000,50000 µg/ml is tested against micro organisms

Staphylococcus aureus, *E. coli*, *Klebsiella pneumonia*, *Pseudomonas aeruginosa* and *Candida albicans* by Kirby-Bauer's disc diffusion method with Amikacin / Ketoconazole as positive control and normal saline as negative control.

2.7 Antimicrobial Screening

Antimicrobial Assay / Antimicrobial activity was evaluated by the Disc diffusion method on nutrient agar medium (NAM) for bacteria and Sabouraud's dextrose agar medium (SDA) for fungi. The Nutrient Agar Medium was prepared by dissolving Beef extract-0.3%, Yeast extract- 0.3%, Peptone-0.5%, NaCl-0.5%, Agar medium-1.0gram, Distilled water-1000ml and maintained PH 7.0. The Sabouraud's Dextrose Agar Medium was prepared by dissolving Peptone 10 grams, Dextrose- 20.0grams, Agar-15.0 grams, cycloheximide and Chloramphenicol in 1000ml water and pH maintained at 5.4 and autoclaved at 115°C for 15 mins. The sterile medium (20ml) was uniformly inoculated by using sterile cotton swabs with test pure cultures of bacteria *Escherichia coli*, *Pseudomonas aeruginosa*, *Klebsiella*, *Staphylococcus aureus* and fungus *Candida albicans*. The discs (5mm in diameter) were impregnated with 10000µg/ml, 30000µg/ml and 50000µg/ml extract of *Murray Koenigii* curry leaves (MEKML) followed by air drying and were placed on seeded agar plates. A disc with normal saline kept as negative control. Another disc with Amikacin kept as Positive Control and Ketoconazole was kept as positive control for *Candida albicans*. For each treatment two replicates were prepared. The plates were incubated for bacteria at 37°C for 24 hours and for fungi 28°C for 48 hours. After incubation the resulting zone of inhibition were measured.

3. RESULTS AND DISCUSSION

The antimicrobial and antifungal action of extract of *Murray Koenigii* curry leaves (MEMKL) was confirmed in our study. The extract showed inhibition of bacterial growth. The zones of inhibition for bacteria i.e., *Escherichia coli* is 20 mm for 50000µg/ml; showing 83.33% inhibition zone. For *Pseudomonas* the inhibition zone was 10 mm with 50000µg/ml exhibiting 45.5 % inhibition. There was no inhibition zone with *Klebsiella pneumonia* with 50000µg/ml. *Staphylococcus aureus* exhibited 15 mm inhibitory zone with 65.21% of inhibition when compared with positive controls Amikacin for gram negative and Augmentin (Amoxicillin / Cloxacillin combination) for gram positive. Whereas *Candida albicans* has shown 15mm inhibitory zone with 50000µg/ml extract, the percentage of inhibition being 65.21 when compared to Positive Control ketoconazole as shown in Table 1.

Table 1: Minimum Inhibitory Concentration of Methanolic Extract of *Murraya Koenigii* Leaves (MEMKL) and Amikacin / Ketoconazole in Serial Dilution Method

Organism	Negative Control (Normal Saline)	Positive Control	MEMKL 10000 µg/ml	MEMKL 30000µg/ml	MEMKL 50000 µg/ml	% of Inhibition
<i>Staph. aureus</i>	Nil	Augmentin 23 mm	Nil	Nil	15 mm	65.21%
<i>E. coli</i>	Nil	(Amikacin) 24mm	Nil	Nil	20mm	83.33%
<i>Pseudomonas aeruginosa</i>	Nil	(Amikacin) 28mm	Nil	Nil	10mm	45.5%
<i>Klebsiella pneumonia</i>	Nil	(Amikacin) 22mm	Nil	Nil	Nil	Nil
<i>Candida albicans</i>	Nil	ketoconazole 22 mm	Nil	Nil	16mm	72.72%

MEMKL- Methanolic Extract of *Murraya Koenigii* Leaves, Positive—Control Discs = Amikacin for *E. coli*, *Klebsiella*, *Pseudomonas aeruginosa*), Augmentin- (Amoxicillin/Cloxacillin combination) for *Staphylococcus aureus*, Ketoconazole for *Candida albicans*

5. CONCLUSION

In conclusion, this study confirms that *Murray Koenigii* curry leaves possesses in-vitro antimicrobial activity. This obviously justifies the use of *Murray Koenigii* curry leaves in traditional medicine. Further research has to be carried out to elucidate basic mechanism of antimicrobial action of above plant *Murray Koenigii*.

REFERENCES

- Gopalan C, Ramasastri BV and Balasubramaniam, SC: Nutritive value of Indian foods, National Institute of nutrition, Hyderabad. Indian council of medical research, New Delhi, 1984; 66 -117.
- Dutta S. Ind Soap J 1958; 23:201
- The Wealth of India, Vol – VI, CSIR, New Delhi. 1962; 448 – 452.
- Satyavathi GV, Riana MK and Sharma M. Indian Medicinal Plants, Vol – II, INMR, New Delhi. 1983; 289–299.
- Ahmad I, Mehamood Z and Mohammad F (1998), "Screening of Some Indian Medicinal Plants for their Antimicrobial Properties", *J. Ethnopharmacol.*, Vol. 62, pp. 183-193.

6. Arulselvan P, Senthil kumar G P, Sathish Kumar D and Subramanian S "Anti-Diabetic Effect of *Murraya Koenigii* Leaves on Streptozotocin Induced Diabetic Rats", *Pharmazie*,2006; Vol. 61, No. 10, pp. 874-877.
7. Arulselvan P, Subramanian S P,"Beneficial Effects of *Murraya Koenigii* Leaves on Antioxidant Defense System and Ultra Structural Changes of Pancreatic Beta cells in Experimental Diabetes in Rats", *hems. Biol. Interact.*,2000;Vol. 165, No. 2, pp.155-164.
8. Ningappa, M. B., Dinesha, R., and Srinivas, L. Antioxidant and free radical scavenging activities of polyphenols-enriched curry leaf (*Murraya Koenigii* L.) extracts. *Food Chemistry*,2008; 106, 720–728.
9. Ghani A, *Medicinal Plants of Bangladesh: Chemical Constituents and Uses*, 2nd Edition, Dhaka, Asiatic Society of Bangladesh, 2003; pp. 309-310.
10. Iyer D and Uma D P, "Phyto-Pharmacology of *Murraya Koenigii*", *Pharmacognosy Reviews*, 2008; Vol. 2, pp.180-184.
11. Syam S, Abdul A B, Sukari M A, Mohan S,Abdelwahab S I, Wah T S, "The Growth Suppressing Effects of Girinimbine on Hepg2 Involve Induction of Apoptosis and Cell Cycle Arrest", *Molecules*, Vol. 16, No. 8, pp. 7155-7170.
12. Arpita, Srivastava, Sachin Jain, Vidhi Gautam, Neeraj Srivastava, R.K.Sharma and K. Shrman,*Murraya Koenigii* (curry leaves) - a review.Science Monitor, An International journal of pharmaceutical sciences 2011;Vol - 4, Issue - 4, Jul-Sept 2013
13. Ramsewak, R. S., Nair, M. G., Strasburg, G. M., De Witt, D. L., and Nitiss, J. L..Biologically active carbazole alkaloids from *Murraya Koenigii*. *Journal of Agricultural and Food Chemistry*, 1999; 47, 444–447.
14. Khanum, F., Anilakumar, K. R., Sudarshana Krishna, K. R., Viswanathan, K. R., and Santhanam, K.. Anticarcinogenic effects of curry leave indimethylhydrazine-treated rats. *Plant Foods for Human Nutrition*, 2000; 55,347–355.
15. Prescott, L. M., Harley, J. P., and Klein, D. A. *Introduction to microbiology* (5th ed.). The Benjamin Cummins Publishing Co. Inc 1996; (pp. 681–684)
16. O.Batista, A. Duarte, J. Nascimento, and S M.F. Imones, *J. of Nat.Product.*1994, 57: 858-861.
17. M. Goji, K.Asres, N.Gemeda, K .Yirsaw. *Ethio Pharma J*, 2006; 24:130-5.
18. Narasimhan NS, Paradkar MV, Kelkar SL.Alkaloids of *Murraya Koenigii*, Structures ofMahanine, koenine, koenigine and koenidine. *Indian JChem* 1970; 8:473-476.275
19. Narasimhan NS, Paradkar MV, Chitguppi VP,Kelkar SL. Alkaloids of *Murraya Koenigii*, Structuresof mahanimbine, koenimbine, mahanine, koenine, koenigine and koenidine. *Indian J Chem* 1975; 13:993-995.
20. Adebayo AC, Avoola OF, Iwalewa EO, Akindahunsi AA, Omisore NO, Cadewunmi CO et al.Anti-trichomonal, biochemical and toxicologicalactivities of methanolic extract and some carbazolealkaloids isolated from the leaves of *Murraya Koenigii*growing in Nigeria. *Phytomedicine* 2006; 13(4): 246-54.
21. Tachibana Y, Kikuzaki H, Lajis NH, Nakatani N.Comparison of antioxidative properties of carbazole alkaloids from *Murraya Koenigii* leaves. *Asian Journal of Pharmacy and Life Science*,JAgric Food Chem 2003; 51(22): 6461-7.
22. Adebajo AC, Reisch J: Minor furocoumarins of *Murraya Koenigii*. *Fitoterapia* 2000; 71(3): 334-7.
23. Sumit Gupta, Padmaa M Paarakh, Usha Gavani.Isolation of Phytoconstituents from the leaves of *Murraya Koenigii* Linn. *Journal of PharmacyResearch* 2009, 2(8), 1313-1314.
24. Maths MV, Balasubramaniam P. *curry leaves*. *Br Dent J* 2004; 197:519.
25. Nevin TA, Bibby GB. The effect of water soluble chlorophyll on pure cultures of microorganisms, commonly found in the oral cavity. *J Dent Res* 1951;30:469.
26. Argal MS, Kumar S, Choudhary HS, Thakkar RM, Verma SK, Seniya C. The efficacy of *Muiraya Koenigii* leaf extract on some bacterial and fungal strain by disk diffusion method. *J Chem Pharm Res* 2011; 3:697-704.
27. Kishore N, Dubey NK, Tripathi RD, Singh SK.Fungi toxic activity of leaves of some higher plants.*Natl. Acad. Sci. Lett* 1982; 5(1): 9.
28. Sunitha Jagalur Doddanna, Shilpa Patel, Madhusudan Astekar Sundarrao, Ravindra Setru Veerabhadrapa Antimicrobial activity of plant extracts on *Candida albicans*: An *in vitro* study, *Indian Journal of Dental Research*, 2013, 24(4), 401-405