



Anti-inflammatory Activity, Total Flavonoids and Tannin Content from the Ethanolic Extract of *Ageratum conyzoides* Linn. Leaf

Md. Musfizur Hassan¹, A.F.M Shahid-Ud-Daula², Ismet Ara Jahan³, Ishrat Nimmi³, Tarek Adnan², Abdullah-Al-Mansur³ and Hemayet Hossain^{3*}

¹ Department of Pharmacy, Jahangirnagar University, Savar, Dhaka-1342, Bangladesh

² Pharmacy Department, Noakhali Science and Technology, Noakhali-3814, Bangladesh

³ BCSIR Laboratories, Dhaka, Bangladesh Council of Scientific and Industrial Research, Dr. Quadrat-E-Khuda Road, Dhaka-1205, Bangladesh

Received on: 06/03/2012

Accepted on: 23/04/2012

ABSTRACT

The crude ethanolic extract of the leaves of *Ageratum conyzoides* Linn. (Family: Asteraceae) was evaluated for its possible anti-inflammatory activity as well as total flavonoids and tannins content growing in Bangladesh. The anti-inflammatory activity was studied using carrageenan and histamine-induced rat paw edema test at different doses (200 and 400 mg/kg body weight) of the ethanol extract. At the dose of 400 mg/kg body weight, the extract showed a significant anti-inflammatory activity both in the carrageenan and histamine-induced oedema test models in rats showing 59.15% and 57.76% reduction in the paw volume comparable ($P < 0.01$) to that produced by the standard drug indomethacin (61.27% and 63.35%) at 4h respectively. The percentage inhibition of the oedema paw volume by the 400 mg/kg body weight of the extract was also statistically significant ($P < 0.05$; $P < 0.01$) compared favorably with the indomethacin treated animals at 1, 2 and 3 h in both models. The total flavonoids and tannins content were calculated as quite high in ethanolic extract (281.88 mg/g of quercetin equivalent and 86.47 mg of gallic acid equivalent respectively). Acute toxicity test showed that the plant might be safe for pharmacological uses. Therefore, the obtained results tend to suggest the acute anti-inflammatory activity as well as total flavonoids and tannins content from the ethanolic extract of the leaves of *Ageratum conyzoides* and thus provide the scientific basis for the traditional uses of this plant as a remedy for pain and inflammations.

Key Words: *Ageratum conyzoides*, Anti-inflammatory, Carrageenan, Histamine, Total flavonoids, Total tannins.

INTRODUCTION

Ageratum conyzoides (*A. conyzoides*) Linn. (Asteraceae) is an annual herbaceous plant with long history of traditional medicinal and agricultural uses in several countries of the world. *A. conyzoides* is one of such plants with folkloric reputation for use as antimalarial agent. The plant has been known since ancient times for its curative properties and has been utilized for the treatment of various ailments, such as burns and wounds, infectious diseases, arthritis, fever¹. Specifically, it has been shown to have activity against larvae of the mosquito *Aedes aegypti*, hepatoprotective and

radioprotective effect^{2,3}. *A. conyzoides* is rich in polyoxygenated flavonoids, 21 of them have been reported in the whole plant. Among them there are 14 polymethoxylated flavones. These polyhydroxyflavones include quercetin, kaempferol and their glycosides⁴. Oral administration of ethanol extract of *A. conyzoides* has been reported to exhibit antibacterial, wound healing and gastroprotective activity^{5,6}.

Chronic inflammatory diseases remain one of the world's major health problems^{7,8}. Inflammation is the response of living tissues to injury. It involves a complex array of enzyme activation, mediator

release, extravasations of fluid, cell migration, tissue breakdown and repair^{9, 10}. Non-steroidal anti-inflammatory drugs (NSAID) are among the most commonly prescribed drugs due to their consistent effectiveness in the treatment of pain, fever, inflammation and rheumatic disorders. However, their use is associated with adverse effects at the level of digestive tract, ranging from dyspeptic symptoms, gastrointestinal erosions and peptic ulcers to more serious complications, such as over bleeding or perforation¹¹. Therefore to overcome the toxicity of NSAID, the development of new anti-inflammatory drugs is still necessary and the natural product such as medicinal plants could lead in discovering new anti-inflammatory drugs with less undesirable effects¹². Now-a-days attention is being focused on the investigation of the efficacy of plant-based drugs used in the traditional medicine because they are cheap, have little side effects and according to WHO, about 80% of the world population still rely mainly on herbal remedies¹³.

Since no literature is currently available to substantiate anti-inflammatory activity as well as flavonoids and tannins content from the ethanolic extract of the leaves of *Ageratum conyzoides* growing in Bangladesh, the present study was designed to provide scientific evidence for its use as a traditional folk remedy by investigating the anti-inflammatory activity as well as total flavonoids and tannins content from the ethanol extract that also confirm its use as a remedy for pain and inflammations.

MATERIALS AND METHODS

Collection and Identification of Plant Materials

The plant (leaves) *A. conyzoides* was collected at November, 2010 from Noakhali, northeast district of Bangladesh and was identified by Bangladesh National Herbarium, Mirpur, Dhaka. (Accession number-DACB-39526).

Preparation of Ethanolic Extract

The leaves of *A. conyzoides* were freed from any of the foreign materials. Then the plant materials were chopped and air-dried under shed temperature followed by drying in an electric oven at 40° C. The dried plant materials were then ground into powder. About 500g of powdered material was taken in a clean, flat-bottomed glass container and soaked in 1.2 liters of 80% ethanol. The container with its contents was sealed and kept for a period of 4 days accompanying occasional shaking and stirring. The ethanolic extract was filtered by Buchner funnel and the filtrate was concentrated with rotary evaporator at bath temperature not exceeding 40° to have gummy concentrate of

greenish black extract (Yield approx. 13.86%).

Test for Different Chemical Groups

The crude ethanolic extract was tested for its different chemical groups as alkaloids, flavonoids, gums, reducing sugars, saponins, steroids and tannins¹⁴. In each test 10% (w/v) solution of the extract in ethanol was taken.

Experimental Animals and Drug

For the screening of in vivo anti-inflammatory activity male rats of Wister strain weighing 175-202 g were used. The animals were housed under standard Laboratory (at Pharmacology Laboratory of BCSIR, Chittagong) conditions maintained at 25±1°C and under 12/12 h light/double cycle and feed with Balanced Trusty Chunts and water ad libitum. All experimental protocols were in compliance with BCSIR Ethics Committee on Research in Animals as well as internationally accepted principles for laboratory animal use and care.

The standard drug Indomethacin was used for this study and purchased from Square Pharmaceuticals Ltd, Bangladesh.

Chemicals

Tannic acid, quercetin, carrageenan, folin-ciocalteu phenol reagent and histamine were obtained from Sigma Chemical Co. (St. Louis, MO, USA). Tween 80, aluminium chloride, potassium acetate and sodium carbonate were of analytical grade and purchased from Merck (Darmstadt, Germany).

Acute Toxicity Test

The acute toxicity of *A. conyzoides* ethanolic extract was determined in rats according to the method of Hilaly *et al*¹⁵ with slight modifications. Rats fasted for 16h were randomly divided into groups of five rats per group. Graded doses of the extract (200, 400, 800, 1600 and 3200 mg/kg p.o.) were separately administered to the rats in each of the groups by means of bulbed steel needle. All the animals were then allowed free access to food and water and observed over a period of 48h for signs of acute toxicity. The number of deaths within this period was recorded.

Anti-inflammatory Activity

Carrageenan-induced oedema test

Carrageenan induced rat hind paw edema was used as the animal model of acute inflammation according to the method of Lanhers *et al*¹⁶. In this experiment, the rats were divided into four groups of five animals each. Group I (control) received 2% Tween 80 in normal saline (2 ml/kg). Group II (Positive control) received 10 mg/kg body wt. of

Indomethacin orally. Group III and IV received 200 and 400 mg/kg body wt. of the extract orally respectively. Acute inflammation was induced in all the four groups by sub plantar injection of 0.05 ml of its suspension of Carrageenan with 2% Tween 80 in normal saline in the right Paw of the rats 30 minutes after the oral administration of the tested materials. The paw volume was measured with a micrometer screw gauge at 1, 2, 3 and 4h after the administration of the drug and the extract. The percentage inhibition of inflammatory effect of the extract was calculated using the following expression:

$$\text{Percentage inhibition of inflammation} = [(V_c - V_t) / V_c] \times 100$$

Where V_c is the average degree of inflammation by the control group and V_t is the average degree of inflammation by the test group (Table-2).

Histamine-induced oedema test

Using the method of Perianayagam *et al*¹⁷, the paw oedema was produced by sub-plantar administration of 0.1% freshly prepared solution of histamine into the right hind paw of the rats. In this experiment, twenty rats were divided into four groups of five animals each. Group I (control) received 2% Tween 80 in normal saline (2 ml/kg). Group II (Positive control) received 10 mg/kg body wt. of Indomethacin orally. Group III and IV received 200 and 400 mg/kg body wt. of the extract orally respectively. Acute inflammation was induced in all the four groups by sub plantar injection of 0.1 ml of Histamine with 2% Tween 80 in normal saline in the right hind paw of the rats 1h after the oral administration of the tested materials. The paw volume was measured with a micrometer screw gauge at 1, 2, 3 and 4h after the administration of the drug and the extract. The percentage inhibition of inflammatory effect of the extract was calculated using the same formula for carrageenan-induced paw oedema.

Total flavonoids content determination

Aluminium chloride colorimetric method was used for determination of total flavonoids concentration of the ethanol extract¹⁸. The extract (0.5 ml, 1:10 g ml⁻¹) in ethanol were separately mixed with 1.5 ml of methanol, 0.1 ml of 10% aluminum chloride, 0.1 ml of 1 M potassium acetate and 2.8 ml of distilled water. It was allowed to stand for 30 min at room temperature and the absorbance of the reaction mixture was measured at 415 nm with a double beam Analykjena UV/Visible spectrophotometer (Model 205, Jena, Germany). Total flavonoids content was determined as mg of Quercetin equivalent per gram using the equation obtained

from a standard Quercetin calibration curve $y=4.7385x + 0.0355$; $R^2 = 0.9993$.

Total tannins content determination

The tannins were determined using the Folin-ciocalteu Phenol reagent as reported by Amorim *et al*¹⁹. Briefly, 0.1 ml of the sample extract was added with 7.5 ml of distilled water and added 0.5 ml of Folin-ciocalteu Phenol reagent, 1 ml of 35% sodium carbonate solution and dilute to 10 ml with distilled water. The mixture was shaken well, kept at room temperature for 30 min and absorbance was measured at 725 nm. Blank was prepared with water instead of the sample. A set of standard solutions of tannic acid is read against a blank. The results of tannins were expressed in terms of tannic acid in mg/g of dry extract. Total tannin content was determined as mg of tannic acid equivalent per gram using the equation obtained from a standard tannic acid calibration curve $y=4.5692x-0.2538$, $R^2=0.9953$.

Statistical Analysis

Data were presented as mean \pm Standard deviation (S.D). Statistical analysis for animal experiment was carried out using one-way ANOVA followed by Dunnet's multiple comparisons using SPSS Data Editor for Windows, Version 11.5.0 (SPSS Inc., U.S.A.). The results obtained were compared with the control group. p values < 0.05 were considered to be statistically significant (p denotes probability).

RESULTS

Phytochemical Screening

Results of different chemical tests on the ethanolic extract of *A. conyzoides* leaves showed the presence of alkaloid, reducing sugars, saponin, gums, steroid, tannins and significantly presence of flavonoid (Table 1).

Acute Toxicity Test

In acute toxicity study, oral administration of graded doses (200, 400, 800, 1600 and 3200 mg/kg p.o.) of the ethanol extract of *A. conyzoides* to rats did not produce any significant changes in behaviour, breathing, cutaneous effects, sensory nervous system responses or gastrointestinal effects during the observation period. No mortality or any toxic reaction was recorded in any group after 48h of administering the extract to the animals. *A. conyzoides* was safe upto a dose level of 3200 mg/kg of body weight.

Anti-inflammatory Activity

Carrageenan-induced paw oedema

The anti inflammation effect of the ethanolic extract of the leaves of *A. conyzoides* using carrageenan induced oedema tests is expressed in (Table-2). In this test, the positive control (Indomethacin) significantly ($p<0.05$; $p<0.01$) decreased the paw edema at 1h to 4h after carrageenan injection compared to saline with inhibition 50.49% to 59.15%. A maximum oedema paw volume of 1.42 ± 0.07 mm was observed in the control rats, 4 h after the carrageenan injection. Rats with the extract at 400 mg/kg body weight significantly decreased ($p<0.05$; $p<0.01$) the carrageenan-induced oedema paw volume from 1h to 4h compared to the standard drug indomethacin at a dose of 10 mg/kg body weight. The inhibition percentage of the oedema paw volume by the 400 mg/kg body weight of the extract was also found statistically significant when it was compared with the indomethacin treated animals at 1, 2, 3 and 4 h. The highest reduction in the paw volume by the 400 mg/kg body weight was 59.15% was comparable to that of the indomethacin (61.27%) at 4 h.

Histamine-induced paw oedema

Table 3 showed the anti-inflammation effect of the ethanolic extract of *A. conyzoides* leaves using histamine-induced paw oedema tests. In the histamine-induced oedema test, a maximum oedema paw volume of 1.61 ± 0.08 mm was observed in the control rats, 4 h after the histamine injection. Rats pre-treated with the extract at 400 mg/kg body weight significantly decreased ($p<0.05$; $p<0.01$) the histamine-induced oedema paw volume from 1h to 4 h compared to the standard drug indomethacin at a dose of 10 mg/kg body weight. The percentage inhibition of the oedema paw volume by the 400 mg/kg body weight of the extract was also statistically significant ($p<0.05$; $p<0.01$) compared favorably with the indomethacin treated animals at 1, 2, 3 and 4 h. The maximum reduction in the paw volume by the 400 mg/kg body weight was 57.76% compared to the indomethacin (63.35%) at 4 h.

Total flavonoids content

The total flavonoids content was calculated as significant in ethanolic extract of *A. conyzoides* 281.88 mg/g of quercetin equivalent per gm of dry extract (Table-4).

Total tannin content

The total tannin content was calculated as quite high in ethanolic extract 86.47 mg/g of tannic acid equivalent (Table-5).

DISCUSSION

The anti-inflammatory activity was studied using two established method namely carrageenan and histamine-induced rat paw edema test at different doses (200 and 400 mg/kg body weight) of the ethanol extract of *A. conyzoides* leaf.

Carrageenan-induced oedema involves the synthesis or release of mediators at the injured site. These mediators include prostaglandins, especially the E series, histamine, bradykinins, leucotrienes and serotonin, all of which also cause pain and fever²⁰. Inhibitions of these mediators from reaching the injured site or from bringing out their pharmacological effects normally ameliorate the inflammation and other symptoms. Development of oedema induced by carrageenan is commonly correlated with early exudative stage of inflammation²¹. Carrageenan oedema is a multimediated phenomenon that liberates diversity of mediators. It is believed to be biphasic; the first phase (1h) involves the release of serotonin and histamine while the second phase (over 1h) is mediated by prostaglandins, the cyclooxygenase products, and the continuity between the two phases is provided by kinins¹⁷. Since carrageenan-induced inflammation model is a significant predictive test for anti-inflammatory agents acting by the mediators of acute inflammation^{22, 23}, the results of this study are an indication that *A. conyzoides* can be effective in acute inflammatory disorders.

The extract also exhibited pronounced reduction in the oedema produced by histamine. This result tends to suggest that the anti-inflammatory activity of the extract is possibly backed by its anti-histamine activity. The antihistaminic effect of the extract increased with increase in the dose of the extract. Histamine is an important inflammation mediator, potent vasodilator substance and also increases the vascular permeability^{24, 25, 26}. Since the extract effectively suppressed the oedema produced by histamine, it showed that the extract exhibited anti-inflammatory actions by inhibiting the synthesis, release or action of inflammatory mediators such as histamine, serotonin and prostaglandins. . This study has shown that the ethanol extract of the leaves of *A. conyzoides* possessed a significant anti-oedematogenic effect ($P<0.01$) on paw oedema induced by carrageenan and histamine compared favorably with the standard drug (indomethacin) in treated rats.

The anti-inflammatory activity of the ethanolic extract of *A. conyzoides* may also be proved due to the presence of flavonoids in a significant amount (281.88 mg quercetin equivalent per g of dry extract). Flavonoids (or bioflavonoids) are naturally occurring compounds, containing in

vascular plants. These compounds have been considered to possess anti-inflammatory properties, both in vitro and in vivo²⁷. Numerous studies have proposed that flavonoids act through a variety of mechanisms to prevent and attenuate inflammatory responses and serve as possible cardioprotective, neuroprotective and chemopreventive agents²⁸. Phytochemically, the leaves of *A. conyzoides* have been also reported to yield tannins. The total tannin amount was calculated as quite high in ethanolic crude extract (86.47 mg/g of tannic acid equivalent) (table 5). Tannins are important compounds known to be potent cyclooxygenase-1 inhibitors and with anti-phlogistic activity²⁹. The mechanisms of anti-inflammatory activity may be related to the anti-phlogistic action of the tannins. Non-steroidal anti-inflammatory drugs (NSAID) such as indomethacin used in this study are known to inhibit cyclooxygenase enzymes I and II which are implicated in the production of inflammation-mediating agent prostaglandin E₂ (PGE₂) from arachidonic acid^{30, 31}. Therefore, the pattern of anti-inflammatory activity exhibited by this extract was similar to that of indomethacin.

CONCLUSION

Since the plant extract reduced significantly the formation of oedema induced by carrageenan and histamine, the leaves of *A. conyzoides* exhibited acute anti-inflammatory activity. The potential of the extract of *A. conyzoides* as acute anti-inflammatory agent may be due to the presence of phytoconstituents like polyhydroxyflavones include quercetin, kaempferol and their glycosides, tannins, phenolics⁴ and might be responsible for its activity. Again, no mortality was recorded in the acute toxicity test; it showed that the plant might be safe for use. Therefore, it can be revealed that the ethanolic extract of *A. conyzoides* leaves possess acute anti-inflammatory activity and justify its use as a traditional folk remedy for inflammation, pain etc. However, a more extensive study is necessary to determine the exact mechanism(s) of action of the extract and its active compound(s).

Table-1: Results of different group tests of ethanolic extract of *A. conyzoides* leaves.

Plant Extract	Alkaloid	Reducing Sugars	Tannins	Gums	Flavonoids	Saponin	Steroid
EE	+	+	+	+	++	+	+

EE: Ethanol extract of *A. conyzoides*; +: Positive result; ++: significantly positive

Table-2: Effect of ethanol extract of *A. conyzoides* leaves and indomethacin on carrageenan-induced oedema paw volume in male Wistar rats.

Treatment Groups	Doses (mg/kg body weight)	Right hind paw volume (mm)			
		1 h	2 h	3 h	4 h
Control	2 ml/kg (2% tween 80 in normal saline)	1.01±0.08	1.25±0.06	1.36±0.09	1.42±0.07
Positive Control (Indomethacin)	10	0.45±0.06* (55.44)	0.52±0.05** (58.40)	0.54±0.08* (60.29)	0.55±0.06** (61.27)
Extract	200	0.75±0.05* (25.74)	0.76±0.09* (39.20)	0.82±0.06** (39.70)	0.85±0.07** (40.14)
Extract	400	0.50 ± 0.07* (50.49)	0.58 ± 0.09** (53.60)	0.56 ± 0.03* (58.82)	0.58 ± 0.06** (59.15)

Values in brackets denote percentage inhibition of the oedema paw volume.

Values are expressed as mean±SD; Values are calculated as compared to control using one way-ANOVA followed by Dunnet's Test; * indicates $P < 0.05$; ** indicates $P < 0.01$ vs. control; $n = 5$.

Table-3: Effect of ethanol extract of *A. conyzoides* leaf and indomethacin (standard drug) on histamine-induced oedema paw volume in male Wistar rats.

Treatment Groups	Doses (mg/kg body weight)	Right hind paw volume (mm)			
		1 h	2 h	3 h	4 h
Control	2 ml/kg (2% tween 80 in normal saline)	1.07±0.07	1.29±0.09	1.38±0.09	1.61±0.08
Positive Control (Indomethacin)	10	0.47±0.05* (56.07)	0.55±0.07* (57.36)	0.56±0.05** (59.42)	0.59±0.08** (63.35)
Extract	200	0.79±0.09* (26.17)	0.86±0.4* (33.33)	0.91±0.07* (34.05)	0.95±0.05* (40.99)
Extract	400	0.51 ± 0.04* (52.33)	0.59 ± 0.07* (54.26)	0.61 ± 0.06** (55.79)	0.68 ± 0.08** (57.76)

Values in brackets denote percentage inhibition of the oedema paw volume.

Values are expressed as mean±SD; Values are calculated as compared to control using one way-ANOVA followed by Dunnet's Test; * indicates $P < 0.05$; ** indicates $P < 0.01$ vs. control; $n = 5$.

Table-4: Total flavonoids content of ethanol extract of *A. conyzoides* leaf.

Extract	Avg. absorbance at 415 nm	Total flavonoid content
		mg of quercetin equivalent (QE) per gm of dry extract
Ethanol extract of <i>A. conyzoides</i> leaf	1.502±0.06	281.88±0.07

The values are expressed as mean ± standard deviation ($n=3$).

Table-5: Total tannin content of ethanol extract of *A. conyzoides* leaf.

Extract	Avg. absorbance at 725 nm	Total tannin content
		mg of tannic acid equivalent (TAE) per gm of dry extract
Ethanol extract of <i>A. conyzoides</i> leaf	0.314±0.08	86.47±0.094

The values are expressed as mean ± standard deviation ($n=3$).

REFERENCES

- Jansen RK. The systematics of *Acmella* (Asteraceae- Heliantheae). Syst. Bot. Monogr, 1985; 8:110-115.
- Pandey V, Agrawal V, Raghavendra K, Dash AP. Strong larvicidal activity of three species of *Spilanthes* (Akarkara) against malaria (*Anopheles stephensi* Liston, *Anopheles culicifacies*, Species C) and filaria vector (*Culex quinquefasciatus* Say). Parasitol. Res, 2007; 102:171-174.
- Prachayasittikul S, Suphamong S, Worachartcheewan A, Lawung R, Ruchirawat S, Prachayasittikul V. Bioactive Metabolites from *Spilanthes acmella* Murr. Molecules, 2009; 14:850-867.
- Nakatani N, Nagashima M. Pungent alkamides from *Spilanthes acmella* L. var. *oleracea* Clarke. Biosci Biotechnol Biochem, 1992, 56:759-762.
- Sittie AA, Lemmich E, Olsen CE, Hviid L, Broegger Christensen S. Alkamides from *Phyllanthus fraternus*. Planta Med, 1998; 64:192-193.
- Majumder M, Shukla AK, Arunachalam A. Nutrient release and fungal succession during decomposition of weed residues in a shifting cultivation system. Communications in Biometry and Crop Science, 2008; 3(1):45-59.

7. Yesilada EO, Ustun, Sezik E, Takishi Y, Ono Y, Honda G. Inhibitory effects of Turkish folk remedies on inflammatory cytokines: interleukin-1 α , interleukin-1 β and tumor necrosis factor- α . J Ethnopharmacol, 1997; 58: 59-73.
8. Li RW, Myers SP, Leach DN, Lin GD, Leach G. A cross-cultural study: anti-inflammatory activity of Australian and Chinese plants. J Ethnopharmacol, 2003; 85: 25-32.
9. Vane JR, Bolting RM. New insights into the mode of action of anti-inflammatory drugs. Inflamm Res, 1995; 44(1):1-10.
10. Perianayagam JB, S.K. Sharma SK, Pillai KK. Anti-inflammatory activity of *Trichodesma indicum* root extract in experimental animals. J Ethnopharmacol, 2006; 104:410-414.
11. Corrado B, Marco T, Colucci R, Fornai M, Antonioli L, Ghisu N, Tacca MD. Role of coxibs in the strategies for gastrointestinal protection in patients requiring chronic non-steroidal anti-inflammatory therapy. Pharm Res, 2009; 59: 90-100.
12. Halliwell B, Cross CE, Gutteridge JMC. Free radicals, antioxidants and human disease: where are we now? J Lab Clin Med, 1992; 119:598 – 620
13. Muthu C, Ayyanar M, Raja N, Ignacimuthu S. Medicinal plants used by Traditional healers in Kancheepuram District of Tamil Nadu, India. J Ethnobi Ethnomed, 2006; 2:43.
14. Evans WC. Trease and Evan's Pharmacognosy. 13th ed., University Press, Cambridge, 1989, p. 546.
15. Hilaly JE, Israili ZH, Lyoussi B. Acute and chronic toxicological studies of *Ajuga iva* in experimental animals. J. Ethnopharmacol., 2004; 91: 43-30.
16. Lanhers MC, Fleurentin J, Dorfman P, Motrier F, Pelt JM. Analgesic, antipyretic and anti-inflammatory properties of *Euphorbia hirta*. Planta Medica, 1991; 57:225-231
17. Perianayagam JB, S.K. Sharma SK, Pillai KK. Anti-inflammatory activity of *Trichodesma indicum* root extract in experimental animals. J Ethnopharmacol, 2006; 104:410-414.
18. Chang C, Yang M, Wen H, Chern J. Estimation of total flavonoid content in propolis by two complementary colorimetric methods. J. Food Drug Analysis, 2002; 10: 178-182.
19. Amorim ELC, Nascimento JE, Monteiro JM, Peixoto Sobrinho, Araujo TAS, Albuquerque UAP. A simple and accurate procedure for the determination of tannin and flavonoid levels and some applications in ethnobotany and ethnopharmacology. Funct. Ecosyst. Commun, 2008; 2, 88-94.
20. Asongalem EA, Foyet HS, Ekoo S, Dimo T, Kamtchouing P. Anti-inflammatory, lack of central analgesia and antipyretic properties of *Acanthus montanus* (Ness) T. Anderson. J. Ethnopharmacol, 2004; 95: 63-68.
21. Ozaki Y. Anti-inflammatory effects of *Curcuma xanthorrhiza* Roxb, and its active principle. Chemical and Pharmaceutical Bulletin, 1990; 38:1045-1048.
22. Mossai JS, Rafatullah S, Gala AM, Al-Yahya MA. Pharmacological studies of *Rhus retinorrhoea*. International Journal of Pharmacognosy, 1995; 33:242-246.
23. Sawadogo WR, Boly R, Lompo M, Some N. Anti-inflammatory, analgesic and antipyretic activities of *Dicliptera verticillata*. International journal of Pharmacology, 2006; 2:267-273.
24. Cuman, RKN, Bersani-Amadio CA, Fortes ZB. Influence of type 2 diabetes on the inflammatory response in rat. Inflammation Res, 2001; 50: 460-465.
25. Linardi A., Costa SKP, DeSilva GR, Antunes E. Involvement of kinins, mast cells, and sensory neurons in the plasma exudation and paw edema induced by staphylococcal enterotoxin B in the mouse. Euro. J. Pharmacol., 2002; 399: 235-242.
26. Vasudevan M., Gunman KK, Parle M. Antinociceptive and anti-inflammatory effects of *Thespesia populnea* bark extract. J. Ethnopharmacol., 2007; 109: 264-270.
27. Gomes A, Fernandes E, Lima JL, Mira L, Corvo ML. Molecular mechanisms of anti-inflammatory activity mediated by flavonoids. Curr Med Chem., 2008; 15(16):1586-605.
28. Pan MH, Lai CS, Ho CT. Anti-inflammatory activity of natural dietary flavonoids Food Funct., 2010; 1:15-31.
29. Wagner H. Search for new plant constituents with potential anti-phlogistic and anti-allergic activity. Planta Medica, 1989; 55: 235-241.
30. Moody JO, Robert VA, Connolly JD, Houghton PJ. Anti-inflammatory activities of the methanol extracts and an isolated furanoditerpene constituent of

- Sphenocentrum jollyanum* Pierre (Menispermaceae). J. Ethnopharmacol., 2006; 104: 87-91.
31. Dhara AK, Suba V, Sen T, Pal S, Chaudhuri AKN. 2000. Preliminary studies on the anti-inflammatory and analgesic activity of the methanol fraction of the root extract of *Tragia involucrata* Linn. J. Ethnopharmacol., 2000; 72: 265-268.
-

***Corresponding Author:** Md. Hemayet Hossain, M. Pharm.,
Senior Scientific Officer,
Chemical Research Division, BCSIR Laboratories, Dhaka, Bangladesh Council of
Scientific and Industrial Research (BCSIR), Dhaka-1205, Bangladesh.
E-mail ID : hemayethossain02@yahoo.com
Tel.No.: 088-01728805884